OPEN ACCESS Freely available online

Anatomy & Physiology: Current Research

Commentary

Functional Anatomy of the Pelvic Floor in Health and Disease

Hana Novotna*

Department of Biomedical Sciences, Charles Medical College, Brno, South Moravian, Czech Republic,

DESCRIPTION

The The pelvic floor is a highly specialized anatomical and functional structure that plays an indispensable role in maintaining continence, supporting pelvic organs, facilitating childbirth and contributing to sexual function. Composed of muscles, ligaments, fascia, connective tissues and neural components, the pelvic floor acts as a dynamic support system rather than a static anatomical barrier. Despite its critical physiological importance, disorders of the pelvic floor remain underdiagnosed and undertreated worldwide, affecting millions of individuals across all age groups. Recent advances in anatomical imaging, biomechanics and clinical research have considerably enhanced our understanding of pelvic floor function, providing new opportunities for prevention, early diagnosis and personalized management of pelvic floor disorders. The functional anatomy of the pelvic floor is remarkably complex. The levator ani muscle group including the puborectalis, pubococcygeus and iliococcygeus muscles forms the principal muscular component of the pelvic diaphragm. Together with the coccygeus muscle, these structures provide continuous support for the bladder, uterus or prostate, rectum and surrounding pelvic organs. Their coordinated contraction and relaxation regulate urinary and fecal continence while accommodating physiological events such as micturition, defecation, pregnancy and childbirth. This intricate balance is maintained through integrated communication between the central nervous system, peripheral nerves and local connective tissue architecture.

One of the most significant advances in recent years has been the appreciation of the pelvic floor as a dynamic biomechanical system. Rather than functioning independently, pelvic floor muscles operate in close coordination with the diaphragm, abdominal wall and deep spinal stabilizing muscles to regulate intra abdominal pressure and maintain postural stability. This concept has expanded the understanding of pelvic floor physiology beyond urogenital function alone, emphasizing its contribution to core stability, respiratory mechanics and movement coordination. Such insights have influenced rehabilitation strategies that integrate breathing exercises, posture correction and neuromuscular training into pelvic floor therapy. Pregnancy and vaginal childbirth remain among the most important physiological events influencing pelvic floor integrity. Mechanical stretching, muscle trauma, connective tissue injury and nerve damage during delivery may compromise pelvic floor support, increasing the risk of urinary incontinence, pelvic organ prolapse and fecal incontinence later in life. However, aging, obesity, chronic constipation, heavy physical labor, neurological disorders and previous pelvic surgeries also contribute significantly to pelvic floor dysfunction. Importantly, pelvic floor disorders are not limited to women; men frequently experience pelvic floor dysfunction following prostate surgery, neurological disease, chronic pelvic pain syndromes, or traumatic injuries.

Modern diagnostic technologies have transformed the evaluation of pelvic floor anatomy. High resolution magnetic resonance imaging, three dimensional and four dimensional ultrasound, dynamic fluoroscopic studies and electromyography now provide detailed assessments of muscular integrity, connective tissue support and functional movement during physiological activities. These imaging techniques allow clinicians to identify subtle anatomical defects that were previously difficult to detect through conventional clinical examination alone. Improved visualization has also enhanced surgical planning and facilitated objective monitoring of treatment outcomes.

Current research increasingly recognizes that pelvic floor dysfunction is influenced by multiple interacting biological systems. Connective tissue remodeling, hormonal regulation, genetic susceptibility, neuromuscular coordination and inflammatory processes all contribute to disease development. Age related declines in collagen quality and muscle strength reduce tissue resilience, while estrogen deficiency following menopause may further weaken pelvic support structures. Understanding these multifactorial mechanisms is essential for developing targeted preventive and therapeutic strategies rather than relying solely on symptom based management.

Conservative treatment remains the first line approach for many pelvic floor disorders. Pelvic floor muscle training, biofeedback therapy, electrical stimulation, behavioral interventions and lifestyle modifications have demonstrated substantial benefits in improving muscle strength, continence and quality of life.

Correspondence to: Hana Novotná, Department of Biomedical Sciences, Charles Medical College, Brno, South Moravian, Czech Republic, E-mail: hana.novotna@cmc.cz

Received: 03-Jan-2025, Manuscript No. APCR-25-43553; **Editor assigned:** 06-Jan-2025, PreQC No. APCR-25-43553 (PQ); **Reviewed:** 20-Jan-2025, QC No. APCR-25-43553; **Revised:** 27-Jan-2025, Manuscript No. APCR-25-43553(R); **Published:** 03-Feb-2025, DOI: 10.35248/2161-0940.25.15.524

Citation: Novotná H (2025). Functional Anatomy of the Pelvic Floor in Health and Disease. Anat Physiol. 15:524.

Copyright: © 2025 Novotná H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.