

Genomic Adaptation and Functional Diversity of Fungi in Deep Subsurface Rock Habitats

Marcus Leclerc*

Department of Subsurface Microbial Genomics, Nouvelle Université de Grenoble, Grenoble, France

DESCRIPTION

Fungi inhabiting deep subsurface rock formations represent one of the least accessible yet biologically significant components of terrestrial microbial life. These organisms persist in environments characterized by extreme nutrient limitation, absence of sunlight, restricted water movement, high mineral density, and long-term geological stability. Unlike surface ecosystems where environmental variability is frequent, deep rock habitats impose chronic energy scarcity and minimal external inputs. Despite such constraints, fungal communities have been detected in fractured rocks, deep boreholes, subterranean caves, and mining shafts extending several kilometers below the Earth's surface. Genomic studies of these fungi have begun to reveal how life persists under conditions that severely restrict biological activity. One of the central challenges in subsurface environments is the scarcity of organic carbon. Without photosynthetic input, fungi must rely on alternative energy sources derived from chemical reactions within rock matrices or trace organic compounds transported through groundwater systems. Genomic analyses of deep subsurface fungi show enrichment of genes involved in chemolithotrophic interactions, enabling them to exploit inorganic chemical gradients indirectly through symbiotic or associative relationships with bacteria. These interactions may provide access to metabolites produced by bacterial communities capable of oxidizing minerals such as iron, sulfur, or manganese.

Energy conservation strategies are a dominant feature of subsurface fungal genomes. Many species exhibit reduced metabolic rates and streamlined pathways that minimize energy expenditure. Genes associated with high-energy metabolic processes are often present but tightly regulated, allowing activation only under favorable conditions. Comparative genomic studies indicate that subsurface fungi frequently possess fewer redundant metabolic pathways compared with surface-dwelling relatives, suggesting evolutionary pressure toward efficiency rather than metabolic breadth. Transport systems play a fundamental role in nutrient acquisition under conditions of extreme scarcity. Genome sequencing reveals expansion of high-affinity transporter genes responsible for uptake of trace organic

molecules, amino acids, and inorganic ions. These transporters are capable of functioning at extremely low substrate concentrations, reflecting adaptation to environments where nutrients are present only in minute quantities. Variability in transporter gene families among subsurface fungal populations suggests localized adaptation to distinct geological substrates.

Cellular stress management systems are also highly developed in deep rock fungi. Although temperature fluctuations are often limited at depth, other forms of stress such as oxidative imbalance and osmotic variation can occur due to mineral-water interactions. Genomic data indicate enrichment of genes involved in redox regulation, detoxification, and maintenance of intracellular stability. These systems allow fungi to survive prolonged periods of environmental stagnation.

Deoxyribonucleic Acid (DNA) repair mechanisms are particularly important in subsurface habitats where cells may remain dormant for extended periods. Over time, exposure to background radiation and chemical stress can lead to genomic damage. Fungal genomes from deep environments frequently contain expanded sets of genes involved in repair pathways, including double-strand break repair, base excision repair, and recombination-based correction systems. These mechanisms help preserve genomic integrity over long timescales. Water availability in subsurface rock environments is highly variable. Although groundwater may be present in fractures, distribution is often uneven. Fungi adapted to these habitats show genetic modifications associated with desiccation resistance and osmotic adjustment. Genes involved in the synthesis of protective sugars and stress-related proteins are commonly observed across multiple subsurface lineages.

Another defining feature of deep subsurface fungi is their interaction with mineral surfaces. Many species exhibit strong attachment to rock substrates, forming biofilm-like structures that facilitate nutrient exchange and stability. Genomic studies identify genes associated with adhesion proteins, extracellular matrix production, and mineral binding. These traits enable fungi to maintain physical association with rock surfaces where limited nutrients may accumulate. Microbial interactions are a central component of subsurface ecosystems. Fungi coexist with

Correspondence to: Marcus Leclerc, Department of Subsurface Microbial Genomics, Nouvelle Université de Grenoble, Grenoble, France, E-mail: marcus.leclerc.subsurfacefungi@univ-grenoble.fr

Received: 01-Jun-2026, Manuscript No. FGB-26-42487; **Editor assigned:** 03-Jun-2026, PreQC No. FGB-26-42487 (PQ); **Reviewed:** 17-Jun-2026, QC No. FGB-26-42487; **Revised:** 24-Jun-2026, Manuscript No. FGB-26-42487 (R); **Published:** 01-Jul-2026, DOI: 10.35248/2165-8056.26.16.313

Citation: Leclerc M (2026). Genomic Adaptation and Functional Diversity of Fungi in Deep Subsurface Rock Habitats. *Fung Genom Biol.* 16:313.

Copyright: © 2026 Leclerc M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

bacteria and archaea within rock fractures and pore spaces. Genomic evidence suggests that these organisms engage in metabolic exchange networks where byproducts of one species serve as substrates for another. Genes associated with interspecies signaling, metabolite transport, and cooperative resource utilization are frequently observed in subsurface fungal genomes.

Horizontal gene transfer may also contribute to adaptation in subsurface ecosystems. Although less frequent than in bacterial communities, genetic exchange between fungi and other microorganisms has been detected in some studies. These transfers may involve genes associated with metabolism, stress response, or environmental sensing, providing new functional capabilities. Long-read sequencing technologies have significantly improved the study of subsurface fungal genomes. Many earlier investigations were limited by fragmented

assemblies and contamination from co-occurring microorganisms. Modern sequencing approaches enable reconstruction of complete genomes from low-biomass samples, revealing previously unknown genetic diversity.

CONCLUSION

Overall, genomic studies of deep subsurface fungi reveal a unique combination of metabolic efficiency, stress tolerance, and ecological specialization. These organisms demonstrate that life can persist in environments with minimal energy input by relying on tightly regulated genetic systems and cooperative microbial interactions. Continued exploration of subsurface fungal genomics will expand understanding of life in one of Earth's most inaccessible and stable ecosystems.