

Genomic Basis of Environmental Adaptation in Marine Fungi Across Salinity Gradients

Henrietta Sokolov*

Department of Marine Microbial Genomics, Odessa Oceanographic University, Odessa, Ukraine

DESCRIPTION

Marine environments host a wide array of fungal life forms that inhabit coastal waters, deep sea sediments, estuarine systems, mangrove ecosystems, and marine-associated animal and plant hosts. Although fungi were once considered rare in oceanic habitats, modern molecular approaches have revealed that they are widely distributed and genetically diverse. Marine fungi occupy ecological roles ranging from decomposition of organic matter to symbiotic interactions with algae and marine invertebrates. Among the many environmental factors shaping their evolution, salinity stands out as a major selective force influencing genomic architecture and functional adaptation.

Salinity levels in marine ecosystems vary significantly depending on geographic location, freshwater input, evaporation rates, and ocean circulation patterns. Estuarine zones experience fluctuating salinity conditions, while open ocean environments maintain relatively stable salt concentrations. Deep-sea habitats present additional pressures such as high hydrostatic pressure and limited nutrient availability. Marine fungi inhabiting these environments must therefore possess genetic systems capable of responding to multiple environmental stressors simultaneously.

One of the primary physiological challenges associated with high salinity is osmotic stress. Elevated salt concentrations can lead to dehydration of fungal cells and disruption of cellular homeostasis. Genome-wide studies have identified numerous genes associated with osmotic regulation in marine fungi, including those involved in ion transport, compatible solute synthesis, and membrane stabilization. These genetic systems allow fungi to maintain internal balance despite external fluctuations in salt concentration.

Ion transport proteins play a central role in salinity adaptation. These proteins regulate the movement of sodium, potassium, calcium, and chloride ions across cellular membranes. Comparative genomic analyses reveal expansion of transporter gene families in marine fungal species compared with terrestrial relatives. Such expansions provide enhanced capacity to control intracellular ion levels and prevent toxic accumulation under high-salt conditions.

Compatible solutes are small organic molecules that help stabilize proteins and cellular structures without interfering with normal biochemical processes. Marine fungi frequently accumulate compounds such as glycerol, trehalose, and proline derivatives to counteract osmotic stress. Genomic investigations have identified biosynthetic pathways responsible for producing these molecules, with variation in gene copy number and regulation across different marine environments.

Membrane composition is another important factor in salinity tolerance. Cellular membranes must maintain flexibility and functionality despite changes in external conditions. Marine fungal genomes show variation in genes involved in lipid biosynthesis, particularly those regulating unsaturated fatty acid production. These modifications help maintain membrane fluidity under high-salinity and temperature-variable conditions.

Symbiotic interactions represent another important aspect of marine fungal biology. Many species form associations with algae, sponges, corals, and marine plants. These relationships often involve nutrient exchange, chemical signaling, and metabolic cooperation. Genomic studies have identified gene clusters associated with host recognition, attachment, and metabolic integration. Variation in these genes suggests adaptation to specific host organisms and ecological niches.

Secondary metabolite production is particularly prominent in marine fungi. These organisms produce a wide range of bioactive compounds, many of which have ecological roles such as defense, competition, or communication. Genome sequencing has revealed extensive biosynthetic gene clusters responsible for producing polyketides, terpenoids, and alkaloid-like compounds. Differences in these clusters across species reflect adaptation to distinct ecological pressures within marine environments.

Population genomic studies have shown that marine fungal communities exhibit both high diversity and strong environmental structuring. Salinity gradients, temperature differences, and nutrient availability contribute to genetic differentiation among populations. In estuarine environments, where salinity fluctuates frequently, fungal populations often

Correspondence to: Henrietta Sokolov, Department of Marine Microbial Genomics, Odessa Oceanographic University, Odessa, Ukraine, E-mail: henrietta.sokolov.marinegen@oou-ua.edu

Received: 27-Feb-2026, Manuscript No. FGB-26-42484; **Editor assigned:** 02-Mar-2026, PreQC No. FGB-26-42484 (PQ); **Reviewed:** 16-Mar-2026, QC No. FGB-26-42484; **Revised:** 23-Mar-2026, Manuscript No. FGB-26-42484 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2165-8056.26.16.310

Citation: Sokolov H (2026). Genomic Basis of Environmental Adaptation in Marine Fungi Across Salinity Gradients. Fung Genom Biol. 16:310.

Copyright: © 2026 Sokolov H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

display greater genetic variability compared with those in more stable oceanic regions.

Dispersal mechanisms play a significant role in shaping marine fungal population structure. Spores and other propagules can be transported by ocean currents, marine organisms, and atmospheric processes. Genomic analyses suggest that while long-distance dispersal is possible, local adaptation often results in distinct genetic signatures across different marine habitats.

Horizontal gene transfer has been observed in marine fungal genomes, although it occurs less frequently than in bacterial systems. Genetic exchange between marine microorganisms may introduce novel metabolic capabilities, particularly those related to nutrient utilization and stress response. Such events contribute to genomic innovation and ecological flexibility in certain lineages.

Biofilm formation is another important ecological strategy used by marine fungi. Biofilms provide structural protection and enhance nutrient acquisition in challenging environments. Genomic studies identify genes associated with adhesion, extracellular matrix production, and community organization.

Variation in these genes contributes to differences in biofilm-forming capacity among species.

Marine fungi also interact with microbial communities consisting of bacteria, archaea, and other eukaryotes. These interactions can involve competition for resources, metabolic cooperation, or signaling exchanges. Genomic analyses reveal genes associated with antimicrobial compound production and interspecies communication, highlighting the complexity of microbial networks in marine ecosystems.

CONCLUSION

The study of marine fungal genomics continues to reveal how organisms adapt to one of the most variable and chemically complex environments on Earth. Through genome-wide analysis of stress tolerance, metabolic diversity, population structure, and ecological interaction, researchers gain deeper insight into the evolutionary processes shaping marine fungal life. Continued exploration of these systems will further enhance understanding of fungal adaptation and ecological function within global ocean ecosystem.