

Genomic Mechanisms Underlying Fungal Adaptation in Volcanic Soil Environments

Tobias van der Merwe*

Department of Microbial Evolutionary Genomics, Cape Highlands University, Cape Town, South Africa

DESCRIPTION

Volcanic soils represent dynamic environments formed through continuous geological activity, mineral deposition, and progressive weathering of basaltic and andesitic materials. These soils differ markedly from stable terrestrial substrates due to their elevated mineral content, fluctuating temperature gradients near geothermal zones, and irregular nutrient composition. Fungal communities inhabiting such landscapes demonstrate distinct genomic characteristics that reflect long-term adaptation to chemically active and physically unstable conditions.

Fungi colonizing volcanic soils are exposed to high concentrations of iron, sulfur, magnesium, and trace metals released from volcanic parent material. These elements can influence cellular physiology by altering enzyme activity and disrupting membrane stability. Genomic analyses of volcanic soil fungi reveal expanded gene families involved in ion regulation, metal detoxification, and oxidative stress management. These genetic features allow organisms to persist in conditions that would otherwise limit cellular function.

One notable genomic trend is the enrichment of transporter proteins responsible for maintaining intracellular ion balance. These proteins regulate uptake and efflux of metal ions, preventing toxic accumulation. Comparative genome sequencing shows that volcanic soil fungi often possess multiple copies of genes encoding Adenosine Triphosphate (ATP)-dependent transporters and membrane-associated ion channels. Such redundancy likely provides flexibility when environmental metal concentrations fluctuate following volcanic eruptions or geothermal shifts. Another important genomic adaptation involves oxidative stress response systems. Volcanic soils frequently contain reactive chemical species that promote formation of free radicals within fungal cells. To counteract this, fungi exhibit expanded antioxidant gene networks, including enzymes that neutralize reactive oxygen compounds. These systems contribute to cellular stability under chemically reactive conditions and reduce molecular damage during metabolic processes.

Carbon and nitrogen availability in volcanic soils is often irregular due to limited organic matter accumulation. Fungi inhabiting these regions rely on flexible metabolic pathways to utilize diverse nutrient sources. Genomic studies highlight the presence of genes associated with decomposition of recalcitrant organic compounds, including lignin-like structures and volcanic organic residues deposited from surrounding ecosystems. This metabolic versatility enables survival in nutrient-poor environments.

Temperature variability in volcanic regions also shapes fungal genomic evolution. Near geothermal zones, soil temperatures may fluctuate significantly over short distances. Fungal species in these habitats show genetic modifications in heat-shock protein families and membrane stabilization pathways. These proteins assist in maintaining protein structure and cellular integrity during thermal fluctuations. Variation in these gene families across populations suggests localized adaptation to microclimatic conditions. Horizontal gene transfer has been proposed as a contributing factor in fungal adaptation to volcanic soils. Genetic material exchanged between bacteria, archaea, and fungi in these environments may introduce novel metabolic capabilities. Although such events are relatively rare, genomic evidence suggests that certain metabolic pathways in volcanic fungi may have originated through cross-domain genetic exchange, particularly those involved in sulfur metabolism and metal transformation.

Population genomic analyses indicate that fungal communities in volcanic soils are often highly structured geographically. Isolated volcanic systems can lead to genetic divergence among populations due to limited dispersal and strong environmental selection pressures. Over time, this isolation contributes to speciation events and the emergence of lineages uniquely adapted to local geochemical conditions. Reproductive strategies also influence genomic diversity in these fungi. Some volcanic soil fungi rely primarily on asexual reproduction, enabling rapid colonization of newly formed substrates after volcanic activity. Others engage in sexual reproduction, which increases genetic variation and may enhance adaptive potential in unpredictable environments. Genomic comparisons reveal that reproductive

Correspondence to: Tobias van der Merwe, Department of Microbial Evolutionary Genomics, Cape Highlands University, Cape Town, South Africa, E-mail: tobias.vandermerwe.fungi@chu-genmail.za

Received: 01-Jun-2026, Manuscript No. FGB-26-42485; **Editor assigned:** 03-Jun-2026, PreQC No. FGB-26-42485 (PQ); **Reviewed:** 17-Jun-2026, QC No. FGB-26-42485; **Revised:** 24-Jun-2026, Manuscript No. FGB-26-42485 (R); **Published:** 01-Jul-2026, DOI: 10.35248/2165-8056.26.16.311

Citation: van der Merwe T (2026). Genomic Mechanisms Underlying Fungal Adaptation in Volcanic Soil Environments. Fung Genom Biol. 16:311.

Copyright: © 2026 van der Merwe T. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

mode correlates with the degree of genetic variability observed within populations.

Secondary metabolism plays a significant role in ecological interactions within volcanic soils. Many fungi produce chemical compounds that inhibit competing microorganisms or facilitate survival under stress. Genome sequencing reveals diverse biosynthetic gene clusters responsible for production of pigments, organic acids, and antimicrobial compounds. Variation in these clusters suggests adaptation to local microbial communities and environmental pressures. Volcanic soils also support complex microbial consortia, including bacteria, archaea, algae, and other fungi. Genomic studies show that fungal species in these environments possess genes associated with interspecies communication and competitive interactions. These genetic systems allow fungi to coexist or dominate within highly competitive microbial networks. Ecologically, fungi in volcanic soils contribute to early-stage soil formation and nutrient cycling. By decomposing organic material and interacting with mineral substrates, they assist in transforming newly formed volcanic deposits into biologically active soils. Genomic insights into these processes help clarify how life establishes itself in geologically young environments.

CONCLUSION

The study of fungal genomes in volcanic soils continues to expand understanding of adaptation to chemically dynamic and physically unstable environments. Genetic systems involved in ion regulation, stress response, metabolic flexibility, and ecological interaction collectively shape survival strategies. Ongoing genomic research will further reveal how fungi persist and diversify in regions shaped by volcanic activity.

REFERENCE

1. Arnold BJ, Huang IT, Hanage WP. Horizontal gene transfer and adaptive evolution in bacteria. *Nat Rev Microbiol.* 2022;20(4):206-218.
2. Burns BL, Rhoads DD. Meningococcal urethritis: Old and new. *J Clin Microbiol.* 2022;60(11):e0057522.
3. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* 2015;25(7):1043-1055.
4. Rodrigues CM, Jolley KA, Smith A, Cameron JC, Feavers IM, Maiden MC. Meningococcal Deduced Vaccine Antigen Reactivity (MenDeVAR) index: A rapid and accessible tool that exploits genomic data in public health and clinical microbiology applications. *J Clin Microbiol.* 2020;59(1):02161-03120.
5. Darriba D, Posada D, Kozlov AM, Stamatakis A, Morel B, Flouri T. ModelTest-NG: A new and scalable tool for the selection of DNA and protein evolutionary models. *Mol Biol Evol.* 2020;37(1):291-294.
6. Stojijkovic I, Hwa V, de Saint Martin L, O'Gaora P, Nassif X, Heffron F, et al. The neisseria meningitidis haemoglobin receptor: Its role in iron utilization and virulence. *Mol Microbiol.* 1995;15(3):531-541.
7. Ladhani SN, Giuliani MM, Biolchi A, Pizza M, Beebeejaun K, Lucidarme J, et al. Effectiveness of meningococcal B vaccine against endemic hypervirulent neisseria meningitidis W strain, England. *Emerg Infect Dis.* 2016;22(2):309.
8. Harrison OB, Bennett JS, Derrick JP, Maiden MC, Bayliss CD. Distribution and diversity of the haemoglobin-haptoglobin iron-acquisition systems in pathogenic and non-pathogenic neisseria. *Microbiology.* 2013;159(Pt9):1920-1930.
9. Brooks A, Lucidarme J, Campbell H, Campbell L, Fifer H, Gray S, et al. Detection of the united States neisseria meningitidis urethritis clade in the united Kingdom, August and December 2019-emergence of multiple antibiotic resistance calls for vigilance. *Eurosurveillance.* 2020;25(15):2000375.
10. Yoder OC, Turgeon BG. Fungal genomics and pathogenicity. *Curr Opin Plant Biol.* 2001;4(4):315-321.