

Green Processes for Seafood Waste Management towards Bio-Economy

V. Venugopal^{*}

Food Technology Division, Bhabha Atomic Research Centre, Mumbai 40085, India

ABSTRACT

The global seafood industry generates significant amounts of waste, consisting of low value by-catch and processing discards, which include shell, head, bone intestine, fin, skin, etc. besides voluminous amounts of nutrient-rich wastewater as process effluents. These discards and effluents, if not properly valorized, are responsible for heavy losses of valuable nutrients as well as for serious environmental hazards. The problems can be addressed by valorization of the waste by environmentally savvy green processes. These generally involve bioconversion based on microbial fermentation or action of externally added enzymes, which release the ingredients from the food waste matrices. The released nutrients, biomaterials, industrial chemicals and also bio-energy are isolated by appropriate green processes. Cultivation of microalgae in seafood waste generates algal biomass, as single cell proteins (SCP), which are rich sources of proteins, oil, polysaccharides, bio-energy and others. A bio-refinery approach for multiple components recovery is economically viable that can contribute towards a sustainable bio-economy.

Keywords: Seafood discards; Process effluents; Cleaner production; Valorization; Green technology; Bio-refinery

INTRODUCTION

Food security is very much dependent on the sustainable availability of nutritious and safe food. Fisheries and aquaculture, which are major components of international food trade, significantly contribute to nutrition, food security and possibly to poverty alleviation [1]. However, burgeoning world population, climate change, ocean acidification, global warming and the current pandemic are showing strains on the availability of marine resources. In the year 2018, an amount of 178.5 Million Tons (MT), consisting of 96.4 MT of capture fisheries and 82.1 MT of aquacultured items were harvested [2]. The popular seafood items belong to finfish (pelagic, anchoveta, pollock, tuna, herring, mackerel, whiting, and others), and shellfish, which can be crustaceans such as shrimp, krill, crab, lobster, and, mollusks, consisting of bivalves (mainly mussels, oysters, clams, and scallops), cephalopods (squid and cuttlefish), and gastropods (mainly abalone, snails). Unfortunately, the fish stocks that are within biologically sustainable levels have decreased from 90 percent in 1974 to 65.8 percent in 2017 [2]. Furthermore, more than a quarter of marine fish stock is over-fished and the rest increasingly fished at their maximum capacity [3]. In addition, all the available catch does not fetch value as food. A sizeable portion of the harvest, designated as by-catch, although rich in nutrients, have poor consumer acceptance due to unappealing appearance, smaller size and other reasons. Besides, commercial processing of valuable fishery items including shellfish results in discards

consisting of shell, head, bones intestine, fin, skin etc., as high as 50%, depending on the raw material. It has been estimated that on an average, about 40% of seafood produced globally is wasted. Annually about 6 to 8 million tons of shell waste from crab, shrimp and lobster are produced globally [4].On dry weight basis, the finfish wastes have approximately 60% proteins, 19 to 20% fat, and 20 to 21% ash. Wet crustacean shells contain 20 to 40% protein, 20% calcium carbonate and other minerals including magnesium salts and 15 to 20% chitin. Fishery wastes are rich in nutrients including functional proteins, enzymes, gelatin, collagen, peptides, carotenoids, oil rich in long chain omega n-3 Polyunsaturated Fatty Acids (omega-3 PUFA), minerals and others such as chitin, chitosan, glucosamine, etc. Loss of these nutrients denies nutrition to millions all over the world [5,6]. Besides, washing, boiling, filleting, marination and other operations consume as high as 10 to 40 m³ freshwater per ton of the raw material; the process effluents contain significant quantity of proteins and other nitrogenous compounds, Fats, Oils, and Grease (FOG), in soluble, colloidal, or particulate forms as Total Suspended Solids (TSS), which are responsible for high Biological Oxygen Demand (BOD) and Chemical Oxygen Demand (COD) [7]. Disposal of by-catch through ocean dumping can damage the oceanic ecosystem, while landfill causes formation of hazardous gases such as methane, ammonia and hydrogen sulfide. The environmental issues of seafood processing include high consumption of water, generation of effluent streams, consumption of energy and the generation of discards [8,9]. In

Correspondence to: Venugopal V, Former Scientific Officer, Food Technology Division, Bhabha Atomic Research Centre, Mumbai 40085, India, E-mail: vvenugopalmenon@gmail.com

Received date: October 06, 2021; Accepted date: October 20, 2021; Published date: October 27, 2021

Citation: Venugopal V (2021) Green Processes for Seafood Waste Management towards Bio-Economy. Poult Fish Wildl Sci. 9:231.

Copyright: © 2021 Venugopal V. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Venugopal V

addition, shortage of drinking water, eutrophication (growth of unwanted biota), biotic depletion, algal blooms, and extensive siltation of corals are other environmental hazards [7]. The above situation warrants that the seafood industry should make sufficient efforts not only to maintain sustainability of the resources but also to control the industry-related environmental hazards. It has been recognized that if global fisheries were optimally managed, it would generate an additional US \$50 billion as income [3].

There is urgent need to significantly reduce global food waste and post-harvest food losses along the production and supply chains to achieve the sustainable development goal target 12.3 of the United Nations by the year 2030 [2]. It is well recognized that a circular economy can strengthen resource efficiency and reduce the use of additional fossil carbon. In a circular economy, the value of products, materials and resources is maintained for as long as possible and the generation of waste is minimized. Bioeconomy, like circular economy, has similar concepts, which aim at improved resources and eco-efficiency, lower Greenhouse Gases (GHGs) emission, reduce the demand of fossil carbon, and valorize waste and side streams. Bio-economy relies on biotechnology for valorization of waste biomass including waste streams into value added products such as nutritious food, feed, bio-based food supplements, functional ingredients, healthy animal feeds and bioenergy. Biotechnological approaches avoid or make lower uses of toxic and harsh chemicals. Circular bio-economy, which is defined as an intersection of circular economy and bio-economy, aims at utilization and up gradation of organic wastes, thereby significantly resolving food-related global issues [10]. Depletion of fishery resources, their incomplete utilization, generation of nutrient-rich waste biomass and effluents as well as their role in environmental degradation are the major seafood-related global issues. Many of these issues can be significantly addressed by circular bio-economy processes. These essentially rely on the concepts of 'reduce, reuse and re-cycle' for the recovery of renewable biological resources including food, food supplements, functional ingredients, healthy animal feed, and bioenergy from food waste biomass and process effluents [11-13]. These approaches are compatible with the European Bio-economy Strategy (2012), which reconciles food security with the sustainable use of renewable resources, while ensuring environmental protection [10].

It is therefore important that the seafood discards and effluents are properly treated by biotechnological processes to achieve the objectives of bio-economy. Conventional processes for the valorization of seafood waste and waste streams are based on chemical and physical methods, which have several limitations and, therefore cannot be used indiscriminately [14]. For example, acid and alkali treatments of crustacean shell waste for chitin extraction require large volumes of freshwater to wash off alkali and acid, releasing corrosive waste water. Similarly, the chemical production of Chitosan Oligosaccharides (COSs) is environmentally hazardous and also difficult to control the degree of polymerization and acetylation. Solvent extraction of lipids from fishery waste can cause oxidation of unsaturated fatty acids [13-15]. Therefore cleaner production methods are required for waste treatment. Cleaner production is defined as continuous application of an integrated preventive, environmental strategy applied to products, processes and service [8]. These biotechnology-based green processes employ processing pathways use less harsh chemicals to achieve the objectives of bio-economy. They are environmentally friendly, cost effective and safe and have potential to convert

OPEN OACCESS Freely available online

food waste and wastewater into valuable raw materials thereby satisfying the objectives of sustainable seafood processing and 'water-energy-food nexus [11,15-19]. Figure 1 depicts potentials for the recovery of valuable ingredients from bioprocessing of seafood discards and effluents. This article briefly points out potentials for environmentally friendly green processes for bioprocessing of seafood discards and effluents that can have achieve the objectives of bio-economy



BIOCONVERSION PROCESSES

Valorization of seafood waste essentially depends on its initial bioconversion to release the components from the food matrices followed by extraction of ingredients from the treated materials. The choice of the green processes depend on the nature of the food matrices and the chemistry of the targeted compounds [6,17,18,20,21]. For example, iso-electric pH solubilization precipitation, which is a pH shift process and weak acid-induced gelation are plausible green methods for the recovery of proteins from fishery discards [22].

Bioconversion by microorganisms

Growth of microorganisms in food waste helps bioconversion, providing safe and effective release of components from the food matrices [18,21,23]. Fermentation by suitable microorganisms (aerobic, anaerobic, or facultative bacteria, fungi, mycelium, or microalgae) results in the production of hydrolytic enzymes such as proteases, chitinases and lipases, and others depending on the microorganism. Proteases replace alkali in the conventional treatment for proteolysis, assisting deproteination and demineralization of the waste. Lactic Acid Bacteria (LAB) dependent bioconversions make otherwise inedible foods edible, nutritional, less toxicity, preserve, and decrease cooking time. LABfermentation breaks down sugar to lactic acid, resulting in pH as low as 3.5; the low pH helps optimal activity of enzymes such as acid proteases, which release proteins bound to lipids, carbohydrates, minerals and carotenoids in the waste. The acid also reacts with calcium carbonate, the main mineral component to form calcium lactate resulting demineralization of the shells and also controls growth of contaminant microorganisms [14]. Chitinases catalyze the cleavage of the β -1,4-O-glycosidic bonds in chitin. Lipases

Venugopal V

function as triacylglycerol hydrolases and also catalyze the synthesis of ester compounds. The fermentation processes can be batch or fed batch types operated under solid, submerged, liquid, or anaerobic conditions. In the commonly used fed-batch process the culture medium is added continuously or in pulses to reach the optimal volume in the reactor. The advantages of fed-batch over the conventional batch operation are higher productivity and decrease in fermentation time. Solid state fermentation of fish offal in presence of sawdust and wood shavings in equal proportions gave a nutritive fertilizer [24]. Microbe-assisted aerobic bioprocess of fishery waste including aquaculture solid waste resulted in bioconversion of nitrogen into ammonium ions [25].

Microbial fermentation processes are highly desirable due to easy handling, simplicity, rapidity, controllability through optimization of process parameters, and negligible requirements of organic solvents. Fermentation is safe, environmental-friendly and low energy consuming. Bio-extraction of chitin is emerging as a green, cleaner, eco-friendly and economical process. Fermentation increases the quality of protein hydrolysates, oil, chitin and produces antioxidant compounds [17,26]. Successful pilot production of chitin using LAB fermentation process has been reported in the early 2000s. The epiphytic Lactobacillus acidophilus isolated from shrimp waste was used to ferment shrimp waste on a pilot scale. The pH of the fermentation broth decreased to a value as low as 3.99 after 12 hr. Besides, the high protease activity of the LAB helped quick removal of minerals and protein. The residue of the fermented shrimp waste contains less than 1% minerals and can be easily transformed into chitin by a mere bleaching treatment [27]. Fermentation by Bacillus megaterium and B. subtilis recovered chitin along with protein hydrolysate from prawn shell waste [28]. Chitosan is a versatile compound having diverse uses. Production of high viscosity chitosan from biologically purified chitin isolated by microbial fermentation has been reported [29]. Protein hydrolytes and calcium lactate are by-products that are useful for animal feed and calcium supplements [4]. Fish viscera may contain 19 to 21 lipids; up to 85% of the lipids could be recovered by fermentation [30]. Several microbial bioconversion processes for seafood waste valorization have been discussed recently [6].

The conversion of food waste including seafood discards into algal biomass resource has attracted attention as a green and economically viable process to recover functionally active compounds. This makes use of cultivation of microalgae such as chlorella, spirulina, dunaliella, diatoms, and cyanobacteria (blue green algae) in the nutrient rich waste in open ponds, under heterotrophic conditions or in closed photo-bioreactors. Growth of these organisms allows degradation of organic contents, and unused food with the removal of CO₂, NH₃-N, CO₂, and H₂S, thereby reducing environmental pollution. The cultivated algal biomass referred as single cell proteins (SCP) can contain up to 60% proteins, good amounts of oil, and also polysaccharides, minerals, carotenoids, and other biomaterials. These ingredients can be extracted from SCP by suitable means. The algal technology has good scope to develop proteins, oil, bioactive peptides, plant growth stimulants, aqua-feeds, food additives, cosmeceuticals, and others [31,32]. Cultivation under stringent nitrogen limitations stimulates the algae to produce lipids as high as 75% rich in n-3 PUFA. Microalgae have the key advantage to produce third generation biofuel (bioethanol, biobutanol, and biogas), because of its rapid growth associated with high lipid production [33]. Chlorella vulgaris accumulated a lipid content of $32.15 \pm 1.45\%$ when cultivated in seafood effluent [34]. Algal products are nutritious and generally safe to human, animals and plants. For instance, spirulina, having 60 to 71% proteins, 13 to 16 carbohydrates and 6-7% fat, is emerging as an effective nutrient to improve poultry productivity [35]. Table 1 gives examples of microbial bioconversions for resource recovery from seafood processing waste.

 Table 1: Examples of resource recovery by microbial bioconversions of seafood processing waste.

Seafood discards	Microorganism	Products	
Shell waste	Lactic acid bacteria	Calcium, carotenoids	
peptides	IDF	IDF	
Shell waste	S. marcescens	Chitinases, proteases,	
pigment	IDF	IDF	
Shrimp waste	L. acidophilus	Chitin, protein hydrolyzate	
Shrimp head	Lactic acid fermentation	Chitin and aquafeed	
Shrimp waste	P. acidolactici	Carotenoids	
Shrimp waste	B. cereus and E. acetylicum.	Chitin, protein	
Prawn shell waste	B. megaterium, B. subtilis	Chitin, protein hydrolysate	
Shrimp waste	Solid state fermentation by L. brevis and R. oligosporus	Chitin	
Crab shell waste	L. paracasei, and S. marcescens	Chitin	
Fish waste	Mixed microorganisms	Liquid fertilizer	
Seafood effluents	Microalgae	Single cell protein [SCP]	
Mussel processing effluents	S. zooepidemicus fermentation	Hyaluronic acid	
Aquaculture solid waste	Aerobic microbial bioconversion.	Liquid fertilizer	
Tuna waste	L. plantarum, B. licheniformis	Aquafeed	
Fresh water fish viscera	P. acidilactici	Oil	
Grass fish bone	Proteolysis and LAB [L. mesenteroides] fermentation	Calcium supplement	
Tuna condensate	Candida rugosa and .L. futsaii	Glutamic acid, gamma aminobutyric acid	

Bioconversion by enzymes

Enzymes can be used as additional processing aids to conventional processes Bioconversions using exogenous enzymes cause direct releases of the components from the food matrices. Hydrolases such as carbohydrases, proteases, and lipases are popular enzymes used for this purpose. The advantages of enzymes are their low energy requirements, safety, and low cost [36]. Chitinases have gained recent biotechnological attention due to their ability to transform chitin in biological waste COSs having wide agricultural, industrial or medical applications. Chitinase Chit42 was expressed in a heterologous system. The enzyme produced small partially acetylated COSs, which have enormous biotechnological potentials in agriculture, medicine and food. Chitosanases are glycosyl hydrolases that catalyse the endohydrolysis of β -1,4-glycosidic bonds of partially acetylated chitosan to release COSs [37-39]. Enzymatic hydrolysis and bacterial fermentation of heads and other discards of several fish species, followed by green extraction processes, recovered Fish Protein Hydrolysates (FPHs), peptones and PUFArich oils. The FPHs, having a degree of hydrolysis of more than 13% and soluble protein concentrations greater than 27 g/L, had good antioxidant and antihypertensive activities. The peptones were viable alternatives to expensive commercial ones [40]. The enzyme, alcalase extracted 26% high quality oil from salmon heads and frames [41]. A few examples on the uses of enzymes for the recovery of various ingredients from diverse fishery wastes are pointed out in Table 2.

 Table 2: Examples of resource recovery by enzymatic bioconversions of seafood discards.

Seafood discards	Enzyme	Components
Shrimp heads	Brevibacilluys sp. alkaline protease	Chitin
Crab and shrimp shells	Deproteination by crab protease	Chitin
Shrimp shell	Proteases	Protein, chitin, carotenoprotein
Shrimp shell	Chitinases, chitosanases and chitin deacetylases	Better defined chitosan and chitosan oligomers
Lobster waste, shrimp waste	Papain, bacterial protease, alcalase, pancreatin	Astaxanthin
Crude chitin	Chitinase	Chitin oligomers, N-acetyl-D-glucosamine
Fin, head, scales	Collagenase, trypsin	Collagen
Fish waste	Proteases	Fish protein hydrolysate
Salmon frame	Proteases	Oil, peptides
Salmon heads	Proteolysis, lipase	PUFA-rich oil
Fish oil	Candida rugosa lipase	PUFA-rich oil
Tuna head	Alcalase	Deodourized oil
Fish bone	Intestine crude proteinase	Bone oligo-phospho- peptide
Fish oil	Candida rugosa lipase	PUFA-rich oil

GREEN PROCESSES FOR RESOURCE RECOVERY

The compounds released by microbial or enzymatic bioconversion processes can be isolated and concentrated by coupling the bioconversion processes with suitable biotechnology-based green innovative processes, which are generally mild, cost-effective, safe, and environmental friendly. They mostly involve reduced process time along with enhanced yields of products. These processes include ultrafiltration, nano-filtration pressurized liquid, subcritical, super-critical fluid, enzyme-mediated, microwave and ultrasound-assisted extractions, and others [18,20,42,43]. LAB fermentation followed by green extraction processes can help sequential or simultaneous recovery of astaxanthin, hydrolyzed protein and chitin from crustacean waste [44]. Green methods are available for fish oil extraction; one of the promising methods is supercritical CO₂ extraction [45]. A sulfated polysaccharide fraction having anticoagulant activity was isolated from the gonads of abalone by enzymatic hydrolysis followed by ion-exchange and gel-filtration chromatography [46]. Bioactive compounds from lobster processing by-products can be extracted using microwave, ultrasonic, and supercritical fluid extraction [47]. Pre-treatment by ultrasonication and microwave radiation improved extraction of chitin by fermentation or chemical routes [38]. Hydrothermal conversion (including hydrothermal liquefaction and hydrothermal

OPEN OACCESS Freely available online

carbonization), is a cost-effective and environmentally friendly thermo-chemical technology for the utilization of biomass as sources of biochar and bio-oil [48] Hydrothermal treatment followed by enzymatic hydrolysis recovered about 85% protein from tilapia scales, giving a gelatin hydrolysate containing peptides in the range of 200-2000 Da. The peptide fraction had stable Angiotensin I-Converting Enzyme (ACE) inhibitory activity [49]. Membrane bioreactors integrate reaction vessels with membrane separation units for producing materials such as peptides, COSs and PUFA from seafood discards [50]. Advancements of waste valorization by biotechnological conversion processes have been discussed [51]. Figure 2 depicts general process for extraction of various ingredients from seafood waste biomass through bioconversion and green methods.



BIOPROCESSES FOR EFFLUENT TREATMENT

The voluminous amount of process effluents is a burden to the seafood industry, as pointed out earlier. The major advantages of effluent treatment are alleviation of environmental hazards, conservation of water, and recovery of commercially valuable ingredients. Biological treatments of effluents make use of aerobic, anaerobic or facultative microorganisms including bacteria, fungi, and protozoa to degrade organic matter present in the waste water. Fermentation of seafood wastewater by organisms such as Saccharomyces sp. and Lactobacillus plantarum in presence of molasses can give a stable product that could be used as feed. Membrane processes are novel cost-effective. These have been discussed in detail [9]. The aquaculture industry is one of the fastest growing sectors in food production and consumes volumes of water for fish farming operations. Discharge of the untreated waste water is highly harmful to the environment. The treatment options for aquaculture effluents include sedimentation, filtration, and mechanical separation, chemical and biological treatments. Electrochemical Oxidation (EO) has potential to treat aquaculture wastewater for removal of multiple pollutants. The EO process has also good bactericidal effect and can remove antibiotics in the effluents [52]. Industrial level treatments of effluents and opportunities, challenges, and prospects of recovery of resources particularly by biological technologies have been discussed in detail [9,53]. Table 3 summarizes some of the green processes for effluent treatment.

Venugopal V

 Table 3: Some green processes for the treatment of seafood process
 effluents.

Process	Advantages
Dissolved Air Flotation (DAF)	Reduce sedimentation time of suspended solids, BOD and COD
Activated sludge consisting of mixed microorganisms	Degrades organic materials in presence of dissolved O ₂ . Removes nutrients and heavy metals. Reduction in BOD, COD, Saves water.
Anaerobic digestion	Involves hydrolysis, liquefaction, and fermentation in the absence of O ₂ . Gives methane for energy use.
Membrane processes: microfiltration (MF), ultrafiltration (UF), nanofiltration (NF), reverse osmosis (RO), forward osmosis (FO). Can be integrated with biological methods	Can efficiently separate organic matter consisting of proteins, bacteria, viruses, etc. The COD and NH ₃ -N removal efficiency of the system can be as high as 95%.
Electrochemical oxidation	Removes proteins, lipids etc. Potential to treat aquaculture wastewater
Cultivation of microalgae in systems such as photobioreactors to give SCP	SCP is source of various biomaterials. Lipids of SCP can be converted into biofuels.
Flocculation by electro-flocculation or by flocculants such as chitosan, carrageenan, alginate, carboxymethyl cellulose, etc.	Sedimentation of proteins and other materials, which can be recovered by membrane processes. Useful for aquaculture water treatment.
Microbial fuel cell	COD reduction, source of energy.

COMMERCIAL USES OF ISOLATED PRODUCTS

The ingredients isolated from seafood waste biomass can have diverse commercial uses [22,23]. The effluents can also be subjected to multiple processes for the isolation of biomaterials including proteins, peptides, amino acids, oil, and others [6,9]. Proteins including collagen and also gelatin isolated from seafood discards as well as effluents invariably retain their natural functional properties including solubility, water-holding, emulsion capacity, antioxidant activities, among others. They can be used as additives including emulsifiers, and texturizers. Protein hydrolysates can function as bio-stimulants for improvement of plant growth, chlorophyll synthesis and as sources of peptides for use as antioxidants, antimicrobials and other functional additives [22]. Lipids are rich in PUFA and therefore can be food additives. Fish oil, when subjected to ozonization and trans-esterification can be source of bio-energy, while carotenoids are natural antioxidants and food colorants [9]. Chitin and other polysaccharides can have diverse industrial and pharmaceutical applications. Applications of ingredients isolated from various seafood discards and effluents have been discussed [6,9].

BIOREFINERY APPROACH FOR COMMERCIAL LEVEL PRODUCTION

Most individual extraction processes of components from seafood waste and effluents, as pointed out earlier, may have limitations

OPEN OACCESS Freely available online

with respect to their economical viabilities. Therefore, recovery of multiple compounds through a cascade of processes based on an integrated bio-refinery approach (defined as 'sustainable processing of biomass into a spectrum of bio-based products) has received recent attention. A successful seafood bio-refinery can have lower operation costs and lower energy consumption, along with high productivity. The bio-refinery allows zero discharge of nutrients from both solid wastes and effluents [54]. The European Commission has opined that these bio-refineries should adopt a cascading approach that favors the highest volume of value added and resource efficient products [10]. Bio-refinery can convert fish waste material into value added biological products such as biofuels, industrial chemicals, animal and fish feed, human food, neutraceuticals and organic fertilizer, etc., [55]. A typical example is the shell refinery where crustacean can be subjected to fermentation, sequential enzymatic, and other processes to recover chitin, proteins including collagen and gelatin, protein hydrolyzates, peptides, PUFA-rich lipids, carotenoids, calcium carbonate, chitin, and chitin oligosaccharides, sulfated and amino-polysaccharides and N-acetylglucosamine [4,56]. Algal biotechnology can be a promising platform for bio-refining of seafood discards on a commercial scale [57]. Cultivation of oleaginous microorganisms including microalgae, yeasts, and bacteria helps valorization of organic wastes for energy production [58,59]. Wastewater treatment incorporating microalgae culture could be greatly developed in the future to achieve a greener environment [34]. The bio-refinery approach can also be extended to make better use of low-cost bycatch fish. Successful pilot plant studies in valorization of seafood waste biomass and effluents including using algal technology have been summarized recently [6,9]. Table 4 gives some examples of bio-refinery based seafood waste treatment. Advances in genetics, biotechnology, process chemistry, and enzyme engineering can further improve the bioconversion and bio-refinery strategy, which can further achieve the objectives of bio-economy [54].

Table 4: A few examples of seafood waste bio-refinery for multiple products having various uses.

Bio-refinery	Products having various uses
Algal bio-refinery, Cultivation of alga, <i>Haematococcus pluvialis</i>	Astaxanthin, SCP
Lactic fermentation	Astaxanthin, hydrolyzed protein and chitin
Sequential treatment of crustacean shells	Chitin, proteins, lipids, carotenoids and CaCO ₃ .
Demineralization and enzymatic degradation of N-acetylglucosamine	Chitin monomers
Sequential enzymatic, acid-alkaline extraction of shrimp cephalothorax	Chitin, chitosan, protein and astaxanthin
Integrated autolysis of shrimp head	Chitin, protein hydrolyzate, sulfated glycosaminoglycans
Sequential treatment of marine cartilage	Chondroitin sulfate, fish meal
Oil extraction, ethanol transesterification, n-3 PUFA concentration	Fish proteins, Glycerol, Lquid bio- fuel
Hydrolysis of fish waste by proteolytic enzymes	Food, Feed, Fertilizer ingredients
Coupled alcalase-mediated hydrolysis and bacterial fermentation	Gelatin, oils, FPH, bioactive peptides, and fish peptones

OPEN ACCESS Freely available online

DISCUSSION AND CONCLUSION

The major problems facing the seafood industry are huge losses of nutrients through the by-catch, process waste biomass and effluents, and the environmental hazards caused by these wastes and effluents including those from aquaculture them. The current situation of stagnating oceanic resources and its adverse impacts on food security demands sustainable and cost effective secondary processing of both seafood waste biomass and process effluents. Green processes, can economically recover valuable resources and bio-energy from the waste biomass, without adverse impacts on the environment. Bio-refinery and algal technology are approaches that can have commercial potentials. The advantages of these processes include total utilization of the fishery harvest not only as food but also as sources of diverse industrial compounds, conservation of water and mitigation of seafood-related environmental hazards. Proper management of seafood discards can ultimately lead to achieve the goals of water-energy-food nexus and seafood sustainability towards bio-economy. Future developments in marine biotechnology can further help realize these goals.

REFERENCES

- Béné C, Arthur R, Norbury H, Allison EH, Beveridge M, Bush S, et al. Contribution of fisheries and aquaculture to food security and poverty reduction: assessing the current evidence. World Dev. 2016;79:177-196.
- 2. FAO. The state of world fisheries and aquaculture 2020. Sustainability in action. Food and Agriculture Organization, Rome. 2020.
- 3. OECD Green Growth Studies. Green growth in the blue economy: Integrating fisheries, aquaculture and the environment, In; Green growth Fish Aquacult. 2015
- Yan N, Chen X. Sustainability: Don't waste seafood waste. Nat. 2015; 524[7564]:155–157.
- Mutalipassi M, Esposito R, Ruocco N, Viel T, Costantini M, Zupo V. Bioactive compounds of nutraceutical value from fishery and aquaculture discards. Foods. 2021;10[7]:1495
- 6. Venugopal, V. Valorization of Seafood processing discards: Bioconversion and bio-refinery approaches, 2021;5:611835.
- 7. Hall M, Delaporte SJ, Phillips A, Beveridge MJ, O'Keefe M, The World Fish Center, Penang, Blue Frontiers. Man Environ Cost Aqu. 2011:92.
- 8. B. Matcon CJ, Christiansen K, Hummrlmose. United Nations Environment Program (UNEP) and Danish Environmental Protection Agency (DEP), Cleaner Production Assessment in Fish Processing. 2000.
- Venugopal V, Sasidharan A. Seafood industry effluents: environmental hazards, treatment and resource recovery. J Environ Cheml Eng. 2020:104758.
- Carus M, Dammer L. The circular bioeconomy-concepts, opportunities, and limitations. Ind Biotech. 2018;14(2):83-91.
- Mao X, Guo N, Sun J, Xue C. Comprehensive utilization of shrimp waste based on biotechnological methods: A review. J Clean Prod. 2017;143:814-823.
- 12.Adegoke SC, Tahergorabi R. Utilization of seafood-processing byproducts for the development of value-added food products. Valoriz Agri Food Wast By-Prod 2021:537-559.
- 13.Alkaya E, Demirer GN. Minimizing and adding value to seafood processing wastes. Food Bioprod Proce. 2016;100:195-202.
- 14. Baiano A. Recovery of biomolecules from food wastes-A review. Molecules. 2014;19(9):14821–14842.

- 15.Yadav M, Goswami P, Paritosh K, Kumar M, Pareek N, Vivekanand V. Seafood waste: A source for preparation of commercially employable chitin/chitosan materials. Biores Biopro. 2019;6(1):1-20.
- 16. Arbia W, Arbia L, Adour L, Amrane A. Chitin extraction from crustacean shells using biological methods-a review. Food Tech Biotech. 2013;51(1):12-25.
- 17. Kaur S, Dhillon GS. Recent trends in biological extraction of chitin from marine shell wastes: A review. Criti Rev Biotech. 2015;35(1):44-61.
- 18.Bruno SF, Ekorong FJ, Karkal SS, Cathrine MS, Kudre TG. Green and innovative techniques for recovery of valuable compounds from seafood by-products and discards: A review. Trends Food Sci Tech. 2019;85:10-22.
- 19.Udugama IA, Petersen LA, Falco FC, Junicke H, Mitic A, Alsina XF, et al. Resource recovery from waste streams in a water-energy-food nexus perspective: Toward more sustainable food processing. Food Bioprod Proces. 2020;119:133-147.
- 20.Martin AM. Fisheries waste biomass: Biconversion alternatives. In Bioconversion of Waste Materials to Industrial Products. Springer. 1998:449-479.
- Dessie W, Luo X, Wang M, Feng L, Liao Y, Wang Z, et al. Current advances on waste biomass transformation into value-added products. Appl Microbiol Biotech. 2020;104(11):4757-4770.
- 22.Sasidharan A, Venugopal V. Proteins and co-products from seafood processing discards: Their recovery, functional properties and applications. Waste Biomass Valoriz. 2020;11(11):5647-5663.
- 23.Shavandi A, Hou Y, Carne A, McConnell M, Bekhit AE. Marine waste utilization as a source of functional and health compounds. Adv Food Nutri Res. 2019;87:187-254.
- 24.Kim JK. Scaled-up bioconversion of fish waste to liquid fertilizer using a 5 L ribbon-type reactor. J Environ Manag. 2011;92(10):2441-2446.
- 25.Khiari Z, Kaluthota S, Savidov N. Aerobic bioconversion of aquaculture solid waste into liquid fertilizer: Effects of bioprocess parameters on kinetics of nitrogen mineralization. Aquacul. 2019;500:492-499.
- 26.Marti-Quijal FJ, Remize F, Meca G, Ferrer E, Ruiz MJ, Barba FJ. Fermentation in fish and by-products processing: An overview of current research and future prospects. Curr Opin Food Sci. 2020;31:9-16.
- Duan S, Li L, Zhuang Z, Wu W, Hong S, Zhou J. Improved production of chitin from shrimp waste by fermentation with epiphytic lactic acid bacteria. Carbo Polym. 2012;89(4):1283-1288.
- 28.Mathew GM, Huang CC, Sindhu R, Binod P, Sirohi R, Awsathi MK, et al. Enzymatic approaches in the bioprocessing of shellfish wastes. 3 Biotech. 2021;11(8):1-3.
- 29.Ploydee E, Chaiyanan S. Production of high viscosity chitosan from biologically purified chitin isolated by microbial fermentation and deproteinization. Int J Polym Sci. 2014;2014.
- 30.Rai AK, Swapna HC, Bhaskar N, Halami PM, Sachindra NM. Effect of fermentation ensilaging on recovery of oil from fresh water fish viscera. Enzy Micro Tech. 2010;46(1):9-13.
- 31. Mohan SV, Hemalatha M, Chakraborty D, Chatterjee S, Ranadheer P, Kona R. Algal biorefinery models with self-sustainable closed loop approach: Trends and prospective for blue-bioeconomy. Biores Tech. 2020;295:122128.
- 32.Nigam S, Sinha S, Srivastava A, Srivastava A. Cultivation and production techniques of marine algae, In Ed. Kim SK Encyclopedia of Marine Biotechnology. 2020.
- 33.Kumar A. Current and future perspective of microalgae for simultaneous wastewater treatment and feedstock for biofuels production. Chem Afri. 2021:1-27.

OPEN CACCESS Freely available online

Venugopal V

- 34.Nguyen TD, Nguyen DH, Lim JW, Chang CK, Leong HY, Tran TN, et al. Investigation of the relationship between bacteria growth and lipid production cultivating of microalgae Chlorella vulgaris in seafood wastewater. Energies. 2019;12(12):2282.
- 35.Deepika DS, Sowmya KL, Kiran KR. Efficacy of microalgal biomass in poultry nutrition. Poult Fish Wildl Sci. 2021;9(5):1-3.
- 36.Venugopal V. Enzymes from seafood processing waste and their applications in seafood processing. Adv in Food NutrI Res. 2016;78:47-69.
- 37. Schmitz C, González Auza L, Koberidze D, Rasche S, Fischer R, Bortesi L. Conversion of chitin to defined chitosan oligomers: current status and future prospects. Marine Drug. 2019;17(8):452.
- 38.Kaczmarek MB, Struszczyk-Swita K, Li X, Szczlsna-Antczak M, Daroch M. Enzymatic modifications of chitin, chitosan, and chitooligosaccharides. Front Bioeng Biotech. 2019;7:243.
- 39.Kidibule PE, Santos-Moriano P, Jiménez-Ortega E, Ramírez-Escudero M, Limón MC, Remacha M, et al. Use of chitin and chitosan to produce new chitooligosaccharides by chitinase Chit42: enzymatic activity and structural basis of protein specificity. Microb Cell Factor. 2018;17(1):1-3.
- 40.Vázquez JA, Meduíña A, Durán AI, Nogueira M, Fernández-Compás A, Pérez-Martín RI, et al. Production of valuable compounds and bioactive metabolites from by-products of fish discards using chemical processing, enzymatic hydrolysis, and bacterial fermentation. Marine Drugs. 2019;17(3):139.
- 41. Liu Y, Ramakrishnan VV, Dave D. Enzymatic hydrolysis of farmed Atlantic salmon by-products: Investigation of operational parameters on extracted oil yield and quality. Process Biochem. 2021;100:10-19.
- 42.Coppola D, Lauritano C, Palma Esposito F, Riccio G, Rizzo C, de Pascale D. Fish waste: From problem to valuable resource. Marine Drugs. 2021;19(2):116.
- 43.Freitas AC, Rodrigues D, Rocha-Santos TA, Gomes AM, Duarte AC. Marine biotechnology advances towards applications in new functional foods. Biotech Adv. 2012;30(6):1506-1515.
- 44.Routray W, Dave D, Cheema SK, Ramakrishnan VV, Pohling J. Biorefinery approach and environment-friendly extraction for sustainable production of astaxanthin from marine wastes. Critic Rev Biotech. 2019;39(4):469-488.
- 45.Ivanovs K, Blumberga D. Extraction of fish oil using green extraction methods: A short review. Energy Procedia. 2017;128:477-483.
- 46.Zhao J, Yang J, Song S, Zhou D, Qiao W, Zhu C, et al. Anticoagulant activity and structural characterization of polysaccharide from abalone (Haliotis discus hannai Ino) gonad. Molecules. 2016;21(6):697.

- 47. Nguyen TT, Barber AR, Corbin K, Zhang W. Lobster processing byproducts as valuable bioresource of marine functional ingredients, nutraceuticals, and pharmaceuticals. Bioresour Bioprocess. 2017;4(1):1-9.
- 48.Usman M, Chen H, Chen K, Ren S, Clark JH, Fan J, et al. Characterization and utilization of aqueous products from hydrothermal conversion of biomass for bio-oil and hydro-char production: A review. Green Chem. 2019;21(7):1553-1572.
- 49.Zhang Y, Tu D, Shen Q, Dai Z. Fish scale valorization by hydrothermal pretreatment followed by enzymatic hydrolysis for gelatin hydrolysate production. Molecules. 2019;24(16):2998.
- 50.Kim SK, Senevirathne M. Membrane bioreactor technology for the development of functional materials from sea-food processing wastes and their potential health benefits. Membranes. 2011 Dec;1(4):327-344.
- Liguori R, Amore A, Faraco V. Waste valorization by biotechnological conversion into added value products. Appl Microbio Biotech. 2013;97(14):6129-47.
- 52.Lang Z, Zhou M, Zhang Q, Yin X, Li Y. Comprehensive treatment of marine aquaculture wastewater by a cost-effective flow-through electrooxidation process. Sci Total Environ. 2020;722:137812.
- 53.Puyol D, Batstone DJ, Hülsen T, Astals S, Peces M, Krömer JO. Resource recovery from wastewater by biological technologies: opportunities, challenges, and prospects. Front Microbio. 2017;7:2106.
- 54.Maina S, Kachrimanidou V, Koutinas A. A roadmap towards a circular and sustainable bioeconomy through waste valorization. Cur Opin Green Sustain Chem. 2017;8:18-23.
- 55.Sahu BB, Barik NK, Paikaray A, Agnibesh A, Mohapatra S, Jayasankar P. Fish waste bio-refinery products: Its application in organic farming. Int J Environ Agric Biotech. 2016;1(4):238605.
- 56.Hülsey MJ. Shell biorefinery: A comprehensive introduction. Green Energy Environ. 2018;3(4):318-327.
- 57. Koyande AK, Show PL, Guo R, Tang B, Ogino C, Chang JS. Bioprocessing of algal bio-refinery: A review on current advances and future perspectives. Bioenginee. 2019;10(1):574-592.
- 58. Cho HU, Park JM. Biodiesel production by various oleaginous microorganisms from organic wastes. Biores Tech. 2018;256:502-508.
- 59.Banu JR, Kavitha S, Gunasekaran M, Kumar G. Microalgae based biorefinery promoting circular bioeconomy-techno economic and lifecycle analysis. Biores Tech. 2020;302:122822.