

Idiopathic Osteoporosis: How Physical Activity Can Reduce the Risk of Developing the Disease

Jiawen Lawless*

Department of Orthodontics, Peking University School and Hospital of Stomatology, Munich, Germany

DESCRIPTION

Osteoporosis is a condition characterized by the loss of bone mass and deterioration of bone tissue, resulting in increased bone fragility and susceptibility to fractures. It is a common disease, affecting an estimated 200 million people worldwide. Osteoporosis is often associated with aging, but there is a type of the disease that affects younger individuals known as idiopathic osteoporosis. Idiopathic osteoporosis is a rare form of the disease that occurs in individuals under the age of 50, with no apparent cause or risk factors. Unlike postmenopausal osteoporosis, which is often caused by hormonal changes and nutritional deficiencies, the cause of idiopathic osteoporosis is not yet fully understood. This has made it difficult to diagnose and treat effectively, leaving many patients at risk of developing serious complications. The symptoms of idiopathic osteoporosis are similar to those of other forms of the disease, including bone pain, fractures, and decreased height. However, because the disease affects younger individuals, it can be especially devastating, as it can significantly impact their quality of life and ability to work and participate in activities. Despite the challenges associated with diagnosing and treating idiopathic osteoporosis, recent advances in medical research are shedding light on the disease and providing new hope for patients. For example, genetic studies have identified several genes that may be associated with idiopathic osteoporosis, including the LRP5 and SOST genes. These genes are involved in regulating bone formation and density, and mutations in these genes may be responsible for the development of the disease. Other research has focused on the role of lifestyle factors in the development of idiopathic osteoporosis. For example, studies have found that individuals who engage in high-impact physical activities, such as running or jumping, have a lower risk of developing the disease. This is thought to be because these activities stimulate bone formation and help to maintain bone density. Conversely, individuals who engage in sedentary lifestyles, such as sitting for

long periods or watching television for extended periods, may be at higher risk of developing the disease. Despite these recent advancements, there is still much that remains unknown about idiopathic osteoporosis. For example, while genetic mutations are thought to play a role in the development of the disease, it is not yet clear why some individuals with these mutations develop the disease while others do not. Similarly, while lifestyle factors are known to influence the risk of developing the disease, it is not clear why some individuals who engage in high-impact physical activities still develop the disease. Despite these uncertainties, there are several steps that individuals can take to reduce their risk of developing idiopathic osteoporosis. These include engaging in regular physical activity, maintaining a healthy diet rich in calcium and vitamin D, avoiding smoking and excessive alcohol consumption, and getting regular bone density screenings. For individuals who have already been diagnosed with idiopathic osteoporosis, there are several treatment options available. These include medications that can help to slow or even reverse bone loss, as well as lifestyle changes that can help to improve bone health. For example, physical therapy can be used to help individuals improve their balance and reduce the risk of falls, while dietary changes can help to ensure that the body has the nutrients it needs to maintain bone density.

CONCLUSION

Idiopathic osteoporosis is a rare but serious disease that can significantly impact the lives of affected individuals. While the cause of the disease is not yet fully understood, recent advances in medical research are providing new insights into the disease and offering new hope for patients. By taking steps to maintain bone health, such as engaging in regular physical activity and eating a healthy diet, individuals can reduce their risk of developing the disease. For those who have already been diagnosed with idiopathic osteoporosis, there are several treatment options available that can help to slow or even reverse bone loss and improve bone health.

Correspondence to: Jiawen Lawless, Department of Orthodontics, Peking University School and Hospital of Stomatology, Munich, Germany, E-mail: jia.lawless@newcastle.ac.de

Received: 02-May-2023, Manuscript No. JOPA-23-23419; **Editor assigned:** 04-May-2023, PreQC No. JOPA-23-23419 (PQ); **Reviewed:** 18-May-2023, QC No. JOPA-23-23419; **Revised:** 22-May-2023, Manuscript No. JOPA-23-23419 (R); **Published:** 02-Jun-2023, DOI: 10.35841/2329-9509.23.11.356

Citation: Lawless J (2023) Idiopathic Osteoporosis: How Physical Activity Can Reduce the Risk of Developing the Disease. J Osteopor Phys Act. 11:356.

Copyright: © 2023 Lawless J. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.