

Indications of Hydrocarbon Prospects in the Lower Benue Trough from Aeromagnetic Data

Choko Chukwuemeka*, Chukwuemeka Ngozi Ehirim, Joseph Onukansi Ebeniro

Department of Physics, University of Port Harcourt, Choba, Nigeria

ABSTRACT

The hydrocarbon prospect of the lower Benue trough has been investigated using aeromagnetic data. Objectively, the study was meant to evaluate structure, lithology and sedimentary thickness that could have direct bearing to hydrocarbon formation and due to the proximity of the study area to the hydrocarbon-rich Niger delta to the south and the discovery of hydrocarbon in Kolmani River II well in Gongola basin, Chad and Niger republics to the northeast that share similar geologic settings. Maps of magnetic parameters were generated using Oasis montaj, Surfer 13 and Arc GIS softwares for subsequent qualitative and quantitative interpretations. Results of the study reveal low and high magnetic anomalies that correspond to basement valleys with thick sedimentary cover and uplifted basement areas with thin sedimentary cover, respectively. Low lineament density, Low magnetic anomalies were delineated in Akwa, Agwu, okposi and Nkalagu areas with 3318.9 m sedimentary thickness which is adequate for the thermal maturation of potential source rocks and hydrocarbon formation. A dominant NNE-SSW structural trend believed to be the continental extension of oceanic fracture system was mapped. This controls sediment deposition, hydrocarbon migration and trap mechanisms that could have direct bearings to the formation of hydrocarbon plays in the trough. Geologic map correlated with the residual magnetic map reveal that areas of low magnetic anomalies with thick sediments corresponds to parts of the Mamu and Nkporo shale formations and Ajali sandstones which are known petroleum systems in the lower trough. With appropriate and favorable juxtaposition of these geologic elements and tectonic structures, prospective hydrocarbon systems could have been formed for possible exploitation in the lower Benue trough. Akwa, Agwu, okposi and Nkalagu areas were thus recommended for detailed seismic surveys.

Keywords: Remote sensing; Hydrocarbon; Aero magnetic; Lower Benue trough

INTRODUCTION

The lower Benue trough has much been studied in the recent past for probable petroleum accumulations. Petroleum exploration in the trough dates back to the early 1930s, following the results of geochemical and geophysical analysis, mapping of surface seeps, sediment alterations and other indications of hydrocarbon presence [1]. It is believed that the trough may have reasonable quantity of hydrocarbon accumulations concealed in deeply buried Cretaceous sediments that satisfy the basic requirement of a hydrocarbon system such as source, reservoir, migration, seal and trap. This is due its proximity to the

hydrocarbon-rich Niger delta to the south and the discovery of hydrocarbon in Kolmani River II well in Gongola basin, Chad and Niger republics to the northeast that share similar geologic settings.

The Niger delta and lower Benue trough are genetically related in the generation and accumulation of hydrocarbons in both basins. According to Ekweozor, et al. [2], the southern boundary of Anambra Basin coincides with the Northern boundary of the Niger Delta. Petroleum accumulations in the northern portion of the Niger Delta contains light waxy oils sourced mainly from land plant derived Organic Matter (OM) disseminated throughout

Correspondence to: Choko Chukwuemeka, Department of Physics, University of Port Harcourt, Choba, Nigeria, Tel: +23408032658241; E-mail: chokochukwuemeka@gmail.com

Received: 23-Dec-2022, Manuscript No. JGG-22-21152; **Editor assigned:** 26-Dec-2022, PreQC No. JGG-22-21152 (PQ); **Reviewed:** 09-Jan-2023, QC No. JGG-22-21152; **Revised:** 16-Jan-2023, Manuscript No. JGG-22-21152 (R); **Published:** 24-Jan-2023, DOI: 10.35248/2381-8719.23.12.1063.

Citation: Chukwuemeka C, Ehirim CN, Ebeniro JO (2023) Indications of Hydrocarbon Prospects in the Lower Benue Trough from Aeromagnetic Data. J Geol Geophys. 12:1063.

Copyright: © 2023 Chukwuemeka C, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the Eocene source rock units, which could possibly be the scenario in the adjacent trough [2,3].

A Several geophysical exploration methods such as gravity, magnetic and seismic have variously been deployed in the search for petroleum in the trough [4]. Of these, the airborne magnetic study has remained the most used geophysical method in the search for petroleum and reconnaissance studies [5-7]. This is because of its cost effectiveness, fast, spatial coverage and resolution. The method maps the thickness, lateral extent and structural texture of the sediments overlying the magnetic basement, which when interpreted could give clues to probable petroleum presence.

The magnetic method depends on the magnetic properties of the rock, depth and geometry of the magnetic source and the external field. Rock magnetism depends on the type and quantity of magnetic mineral in the rock. This determines the magnitudes of the induced and remnant magnetism that varies for the different rock types. Of all the magnetic minerals, magnetite is the most magnetic and of relative abundance such that the magnetism exhibited by a rock under the influence of the earth field is a function of the magnetite mineral content of the rock [8]. Among the different rock types, sedimentary rocks are the least magnetic, igneous the most magnetic and metamorphic is intermediate.

Sedimentary rocks have characteristic low magnetic content or susceptibility and therefore, exhibit uniform magnetic character in contrast to the basement areas. The observed high magnetic anomalies in sedimentary terrains are mostly due to intra-sedimentary intrusive igneous and metamorphic rocks and shallow underlying magnetic basement. Magnetic susceptibility is a fundamental property in magnetic method used for petroleum exploration. Changes in the magnetic susceptibility cause small variations in the magnetic fields of rocks that may indicate the presence of traps and petroleum accumulation as well as intrusives, lava flows and igneous plugs that could provide the geothermal energy needed for the maturation of petroleum source rocks in the sediments [9,10].

The magnetic method rarely maps hydrocarbon directly in the subsurface. This is because the fundamental magnetic property or susceptibility of the rock is not directly sensitive to the presence of hydrocarbons. The analysis of magnetic anomalies gives clues to probable traps due to structure and stratigraphy associated with hydrocarbons accumulation. However, biochemical processes in hydrocarbon bearing sediments result in the alteration of the sediments and accumulation of magnetic minerals diagenetically in the overlying sediment strata by petroleum seepage through faults and micro fractures. This causes local weak magnetic anomalies or aureoles above the hydrocarbon zone, which when isolated from the stronger regional field serves as Direct Indicators of Hydrocarbons (DIH). Temperature gradients, compaction and changes in atmospheric pressure help drive this upward migration [11].

The study area is located at the southeastern part of Nigeria in the lower Benue trough geologic zone, underlain by early to late Cretaceous sediments. It lies between latitude 6° 00' N–7° 00' N and longitude 7° 00' E–8° 00' E, comprising of Nsukka,

Igumale, Udi and Nkalagu areas and covering a total area of 12,000 square kilometers (Figure 1).




Figure 1: Location map of the study area. Note: ● Reference town, ■ Cenozoic, ■ Cretaceous, ■ Volcanic, ■ Basement complex.

The area is characterized by undulating landforms with mixed biomes. Rainfall is at its peak in June and low in November with average daily temperature of 32°C [12,13]. The present study is aimed at mapping and delineating structural and lithological features and estimate depth to magnetic basement (thickness of sedimentary pile) from aeromagnetic data for possible indications of petroleum accumulations in the lower Benue trough.

MATERIAL AND METHODS

Geology of the study area

The study area falls within the Abakaliki and Anambra basins in the lower Benue trough (Figure 2). The lower Benue trough is underlain by thick pile of sedimentary rocks with intra-sedimentary igneous intrusions on the Precambrian basement. The Precambrian Basement Complex is made up of essentially granitic and magmatic rocks which outcrop in the eastern portion of the study area [14]. The stratigraphic sequences of the lower Benue trough range in age from the Aptian to mid Eocene. It is comprised of thick sequence of rocks from earliest to latest such as Asu River Group, Eze-Aku, Awgu, Nkporo Group, Mamu, Ajali, Nsukka, Imo and Bende-Ameki Formations.




Figure 2: Geology map of the study area. Note: ■ Ameki/Nanka formations (Eocene), ■ Imo formation (Thanetian), ■ Nsukka formation (Danian), ■ Ajali sandstone (Maastrichtian), ■ Mamu formation (Maastrichtian), ■ Nkporo group (Campanian), ■ Awga group (Coniacian), ■ Eze-Aku group (Cenomanian/Turonian), ■ Asu-river group (Aptian/Albian).

Two petroleum systems Nkporo-Mamu and Mamu-Ajali were identified, while the overlying Nsukka and Imo shale Formations provides the regional seals [15,16]: Nkporo-Enugu and Mamu shale Formations exhibit characteristics of potential hydrocarbon source rocks for a series of oil/gas and condensate shows found within Ajali reservoir sandstones [17,18].

Method of study

Four sheets of high-resolution aeromagnetic data over Nsukka, Igumale, Udi and Nkalagu areas were obtained from the Nigeria Geological Survey Agency (NGSA) [19]. The data was acquired with nominal flight height of 76 m along N-S flight lines with 500 m inter line spacing and total magnetic field intensity B (nT) was measured.

The data sheets were subsequently assembled and knitted together using Oasis Montaj and Sulfer 13 softwares to produce the Total Magnetic Intensity (TMI) map of the study area. Data were further processed by the removal of regional magnetic effects through the application of filtering (regional-residual separation) to generate a residual magnetic map of the study. Thereafter vertical and horizontal derivative maps were generated from the residual magnetic intensity grid using Oasis montaj software to enhance shallow geological features and identify geologic boundaries in the study area. Source Parameter Imaging (SPI) for depth to magnetic basement was carried out on the residual maps to determine the thickness of the sedimentary pile.

Subsequently, the filtered images (residual gradients) were then used for the extraction of lineaments which were then merged together, traced and measured clockwise from the North automated by Arc GIS program. The mapped lineaments were statistically analyzed and plotted in the form of rose diagrams, using rockworks software program. Finally, these maps were analyzed visually on the basis of amplitude and shape as well as identification of anomalous boundaries, volcanic zones, lineaments and other regional structures that could be akin to those that are associated with petroleum accumulations.

RESULTS

The results obtained from aeromagnetic data were processed and transferred into information through variety of maps. The maps were characterized for magnetic features, basement lineaments and depth to the magnetic source for indications of hydrocarbon in the lower Benue trough.

Total Magnetic Intensity (TMI)

The TMI map of the study area is characterized by low and medium to high magnetic anomalies with intensity values ranging from -51.05 nT to 128.45 nT, respectively (Figure 3). The magnetic lows vary from -51.05 nT to 8.23 nT and dominate in Enugu Ezike, Nsukka and Awgu, while medium to high magnetic anomalies varies from 13.23 nT to 128.45 nT mapped at Igumale, Adani, Enugu, Nkalagu, Okposi, Udi and Ukehe areas. The magnetic lows lie parallel and bounded by the medium to high magnetic intensities and both trends approximately E-W. Low and medium to high magnetic

intensities are related to local and regional magnetic features in the study.




Figure 3: Total Magnetic Intensity (TMI) map of the study area.

Residual Magnetic Intensity (RMI)

The Residual Magnetic Intensity (RMI) map exhibits characteristics low and medium to high magnetic anomalous signatures with intensities varying from -52.80 nT to 54.11 nT (Figure 4). The elongated magnetic lows have intensities varying from -52 nT to 2.73 nT dominate Nsukka, Awgu and Awka areas, while the medium to high ridge-like magnetic anomalies with intensities varying from 4.28 nT to 54.11 nT are mapped in Enugu, Okposi, Udi and Ukehe. The RMI in contrast to the TMI are dominated by local magnetic features with low magnetic intensities. The low and medium to high magnetic anomalies are parallel to each other and trends dominantly in the NE-SW with minor occurrences in the E-W trend.




Figure 4: Residual Magnetic Intensity (RMI) map of the study area.

These broad areas of low and medium to high magnetic intensities corresponds to sagged and uplifted parts of the underlying magnetic basement, respectively. The alternation between low and high magnetic intensities with steep gradients and elongated anomalous features indicate the presence of faults or contacts that divides the study area into blocks with different magnetization contrasts.

Source Parameter Imaging (SPI)

Source Parameter Imaging (SPI) map displays the depth to magnetic sources in the study area (Figure 5). The map shows significant depth variations across the area with depths varying

from -95.5 m to -3318.8 m. The Nsukka, Ukehe, Enugu, Udi, Igumale, Eha-Emufu and Awka exhibit characteristic shallow to medium depths varying from -95.5 m to -561.7 m. Deeper depths to magnetic sources varying from -604.49 m to -3318.9 m, occur mostly in the SE of the study.




Figure 5: Source Parameter Imaging (SPI) map of the study area.

These depths variations are attributed to the tectonic upwarp and downwarp of the magnetic basement and/or magnetic sources, respectively, in the area. The downwarped Awgu, Awka, Nkalagu and Okposi areas are characterized by thick sediment cover than the upwarped Eha-Emufu, Enugu, Igumale, Nsukka, Udi and Ukehe areas with thin sediment cover.

Magnetic lineament

The magnetic lineament map shows the distribution of lineaments in rocks with different magnetic susceptibilities (Figure 6). The lineament distributions are widespread and densely distributed in Udi, Okposi and W and NE of Enugu and Nkalagu areas, respectively and sparsely distributed in the other parts of the study area. They are characterized by both short and long linears, crisscrossing and intersecting each other and forming fault closures capable of forming traps for probable hydrocarbon accumulations.




Figure 6: Magnetic Lineament map of the study area. Note: ★ Town, — Lineament, □ Study area.

Lineament orientation analysis (Figure 7), revealed that the study area is dominated mostly by NE-SW and NNE-SSW lineament orientations, with minor occurrences of NW-SE, N-S and E-W lineament orientations.




Figure 7: Magnetic Rose Diagram of the study area.

The magnetic lineament density map varies from low to high density and correlates with the distribution of lineaments in the study (Figure 8). High lineament densities were delineated in Udi, and W and NE of Igumale and Eha-Amufu areas, respectively, while lows were mapped in Adani, Enugu-Ezike, Awka, Awgu, Enugu, Nsukka, Okposi, Nkalagu and Ukehe areas. These lineament density distributions are reflections of varying degrees of basement tectonics and faulting in the study area. The high lineament density indicates more deformed than the less deformed in low lineament density areas in the study [20].




Figure 8: Magnetic lineament density map of the study area. Note: ★ Town, □ Lineament density ≤ 0.24, ■ ≤ 0.47, ■ ≤ 0.71, ■ ≤ 0.95, ■ ≤ 1.18.

DISCUSSION

The maps of magnetic parameters (TMI, RMI, and SPI) and Lineaments, exhibit characteristics colour gradations and magnetic anomalous distributions caused by subtle contrasts in magnetic susceptibilities of the underlying rocks, reflecting magnetic heterogeneity, structural, lithological and depths to anomalous source variations in the study area. These maps were visually analyzed and evaluated in terms of structure, lithology and contacts for possible indications of hydrocarbon prospect in the lower Benue trough.

The Total Magnetic Intensity (TMI) map is characterized by elliptical magnetic lows and ridge-like magnetic highs trending in the E-W direction. These magnetic signatures suggest anomalous magnetic features mostly of regional origin in the study. However, Residual Magnetic Intensity (RMI) map exhibits distinct elliptical to isolated hemispherical magnetic lows

bounded by ridge-like magnetic highs and both trends mostly in the NE-SW with minor E-W trend in contrast to the E-W trend delineated in the TMI map. This apparent discrepancy in structural orientation is attributed to magnetic susceptibility variations of local magnetic sources at shallow depths.

Low magnetic anomalies were delineated in Adani, Enugu-Ezike, Nsukka, Awka, Awgu, Enugu and Nkalagu areas while magnetic highs were delineated in Eha Amufu, Okposi, Igumale, Udi and Ukehe. Analysis of the SPI data revealed that sediment thickness vary from 95.5 m in the shallow basement depths to 3318.8 m in the large basement depths. This result is in agreement with the estimate of 3300 m of sedimentary thickness by Kogbe, et al. [21], and the works of Daniel, et al. and Onyewuchi, et al. [22,23] in the lower trough. Results show that the magnetic highs are characterized by shallow basement depths in contrast to the large basement depths in the magnetic low areas. The boundaries of these magnetic lows and highs are suspected to be linear belts of fracture zones or lithologic contacts running parallel to the NESW trend in line with the predominant tectonic trends of the trough [24,25].

Magnetic lineaments are widespread but densely distributed in the magnetic highs than low areas in the study area. The lineaments occur predominantly in the NESW and NNE-SSW lineament orientations, with minor occurrences in the NW-SE, E-W and N-S lineament orientations. Magnetic lineaments density analysis reveals variable lineament density across the study area. Lineament density varies from low to high densities and correlate with the spatial distribution of lineaments in the study. According to Hung, *et al.* [20], high lineament density corresponds to the most deformed areas and low lineament density for less deformed areas. Therefore, high lineament densities in Enugu, Udi, Ukehe, Eha-Emufu and Igumale suggests that these areas are more deformed than the low lineament density areas of thick sediment cover in Adani, Enugu-Ezike, Awka, Agwu, Nkalagu and Okposi areas.

According to Ossai, et al. and Nwosu, et al. [26,27], sediment thickness greater than 2000 m is favorable for the thermal maturation of potential source rocks and hydrocarbon formation. Shallow basements depths (-95.5 m to -561.7 m) with high magnetic intensities and high lineament densities were mapped in Enugu, Udi, Ukehe, Eha-Emufu and Igumale and suggests structurally deformed areas with less thick sediment cover unfavourable for hydrocarbon formation but with good prospect for mineral deposits and groundwater resource. Large basement depths (-604.49 m to -3318.9 m) with low magnetic intensities and low lineament densities were mapped in Adani, Enugu-Ezike, to the SW of the study between Awka and Awgu as well as to the NE of Okposi towards Nkalagu. This suggests structurally less deformed areas with thick sedimentary cover and comprised of rocks poor in magnetic content that are associated with potential source rocks and reservoirs favorable for hydrocarbon formation and accumulation in the lower trough.

Correlation of residual magnetic map with the geologic map revealed that areas of low magnetic anomalies with thick sedimentary cover correspond to parts of the Mamu and Nkporo shale formations and Ajali Sandstones. According to

Uzoegbu, et al., Petters, et al., Ekweozor, et al. and Uzoegbu, et al. [15,17,18,28], Nkporo and Mamu shale Formations exhibit characteristics of potential hydrocarbon source rocks for a series of oil/gas and condensate shows found within the Ajali reservoir sandstones. The interbedded marine sandstone facies in the Nkporo and Mamu shales could provide additional reservoirs in the formations [29,30]. The shales and claystones of the Nsukka and Imo Formations may provide regional seals to these petroleum systems in the trough.

CONCLUSION

Magnetic lineaments in the present study are widespread but densely distributed in the magnetic highs than lows area. The lineaments trends predominantly in the NE-SW and NNE-SSW, with minor occurrences in the NW-SE, E-W and N-S lineament orientations.

Geologic map correlated with the residual magnetic map reveal that areas of low magnetic anomalies with thick sediments corresponds to parts of the Mamu and Nkporo shale formations and Ajali sandstones which are known petroleum systems in the lower trough. With appropriate and favourable juxtaposition of these geologic elements and tectonic structures, prospective hydrocarbon systems could have been formed for possible exploitation in the lower Benue trough.

Areas with shallow magnetic anomalous sources, high magnetic intensities and high lineament densities mapped in Enugu, Udi, Ukehe, Eha-Emufu and Igumale are structurally deformed areas with less thick sediment cover unfavourable for hydrocarbon formation but suspected to be good prospect for mineral deposits and groundwater resource. Deep lying magnetic anomalous sources with low magnetic intensities and low lineament densities mapped within SW of the study between Awka and Awgu as well as to the NE of Okposi towards Nkalagu and Adani, Enugu-Ezike, with structurally less deformed areas and thick sedimentary cover, comprising of rocks poor in magnetic content that are related to potential source rocks and reservoirs favorable for hydrocarbon formation and accumulation in the lower trough were thus recommended for detailed seismic surveys.

REFERENCES

1. Kulke H. Gebrüder Borntraeger, Berlin. 1995:146-172.
2. Ekweozor CM, Daukoru EM. Northern delta depobelt portion of the Akata-Agbada (1) petroleum system, Niger Delta, Nigeria. In: The petroleum system. Am Assoc Pet Geol Memoir. 1994;60:599-613.
3. Bustin RM. Sedimentology and characteristics of dispersed organic matter in Tertiary Niger Delta: Origin of source rocks in a deltaic environment. AAPG Bull. 1988;72(3):277-298.
4. Ugwu GZ, Ezema PO. Forward and inverse modelling of aeromagnetic anomalies over Abakaliki and Nkalagu areas of the Lower Benue Trough, Nigeria. Int Res J Geol Min. 2012;2(8): 222-229.
5. Oladele S, Ojo B. Basement architecture in part of the Niger Delta from aeromagnetic data and its implication for hydrocarbon prospectivity. Pac J Sci Technol. 2013;14(2):512-20.
6. Mohamed HS, Senosy MM, Zaher MA. Interactive interpretation of airborne gravity, magnetic, and drill-hole data within the crustal

- framework of the northern Western Desert, Egypt. *J Appl Geophys.* 2016;134:291-302.
7. Di Massa D, Fedi M, Florio G, Vitale A, Viezzoli A, Kaminski V. Joint interpretation of AEM and aeromagnetic data acquired over the Drybones kimberlite, NWT (Canada). *J Appl Geophys.* 2018;158:48-56.
 8. Gao X, Xiong S, Yu C, Zhang D, Wu C. The estimation of magnetite prospective resources based on aeromagnetic data: A case study of Qihe Area, Shandong Province, China. *Remote Sens.* 2021;13(6):1216.
 9. Montgomery CW. *Environmental geology.* McGraw-Hill Companies. 2000:421.
 10. Obi DA, Okereke CS, Obei BC, George AM. Aeromagnetic Modeling of subsurface intrusives and its implication on hydrocarbon evaluation of the Lower Benue Trough Nigeria. *Eur J Sci Res.* 2010;47(3):347-61.
 11. Eventov L. Applications of magnetic methods in oil and gas exploration. *Lead Edge.* 1997;16(5):489-492.
 12. Reyment RA. *Aspects of Geology of Nigeria.* Ibadan University Press, Ibadan, Nigeria. 1965.
 13. Short KC, Stäuble AJ. Outline of geology of Niger Delta. *AAPG Bull.* 1967;51(5):761-779.
 14. Ofoegbu CO, Onuoha KM. A review of geophysical investigations in the Benue Trough. In: *The Benue trough structure and evolution.* 1990:171-201.
 15. Uzoegbu UM, Ekeleme IA, Uchebo UA. Oil generation capacity of maastrichtian coals from the Anambra basin, SE Nigeria. *Int J Eng Sci.* 2014;3:33-46.
 16. Whiteman AJ. *Nigeria: Its Petroleum Geology, Resources and Potential: Volume 1.* Springer Science and Business Media. 1982.
 17. Petters SW, Ekweozor CM. Petroleum geology of Benue trough and southeastern Chad basin, Nigeria. *AAPG Bull.* 1982;66(8): 1141-1149.
 18. Ekweozor CM, Gormly JR. Petroleum geochemistry of late Cretaceous and early Tertiary shales penetrated by the Akukwa-II well in the Anambra Basin, Southern Nigeria. *J Petrol Geol.* 1983;6(2): 207-215.
 19. Nigerian Geological Survey Agency (NGSA). 2011.
 20. Hung LQ, Dinh NQ, Batelaan O, Tam VT, Lagrou D. Remote sensing and GIS-based analysis of cave development in the Suoimuoi catchment (Son La-NW Vietnam). *J Cave Karst Stud.* 2002;64(1): 23-33.
 21. Kogbe CA. The Cretaceous and Paleogene sediments of southern Nigeria. In: *Geology of Nigeria.* 1989;2:325-334.
 22. Daniel NO, Mirianrita NO, Emmanuel O. A case study of aeromagnetic data interpretation of Nsukka area, Enugu State, Nigeria, for hydrocarbon exploration. *Int J Phys Sci.* 2015;10(17): 503-519.
 23. Onyewuchi RA, Ugwu SA. Geological interpretation of the high resolution aeromagnetic data over Okigwe-Udi Area, Anambra Basin, Nigeria, Using 3-D Euler Deconvolution and 2-D Spectral Inversion Methods. *J Geog Envir Earth Sci Int.* 2015;10(1):1-22.
 24. Burke KC, Dessauvagie TF, Whiteman AJ. Geological history of the Benue valley and adjacent areas. In *African Geology.* 1972;1(8):188-205.
 25. Okereke CN, Ananaba SE. Deep crustal lineament inferred from aeromagnetic anomalies over the Niger Delta, Nigeria. *J Min Geol.* 2006;4(2):127-131.
 26. Ossai NM. Interpretation of aeromagnetic data around Nsukka Area, South Eastern Nigeria. University of Nigeria (Unpublished). 2015.
 27. Nwosu IE. Structural interpretation of part of Anambra Basin Nigeria using aeromagnetic and landsat ETM data. *Int J Adv Sci Res Eng.* 2018;4(9):49-59.
 28. Uzoegbu MU, Ukaegbu VU, Wehner H. Geochemical effects of petroleum generation potential from basal Campanian source rocks in the Anambra basin, SE Nigeria. *Adv Appl Sci Res.* 2013;5(5): 278-285.
 29. Murat RC. Stratigraphy and paleogeography of the cretaceous and lower tertiary in Southern Nigeria. In: *African Geology.* 1972;1(1): 251-266.
 30. Nwachukwu SO. The tectonic evolution of the the southern portion of the Benue Trough, Nigeria. *Geol Mag.* 1972;109(5):411-419.