

# Editorial

# Indices of Codon Usage Bias

#### Arif Uddin\*

Department of Zoology, Moinul Hoque Choudhury Memorial Science College, Hailakndi, India

## Introduction

Most of the commonly used indices of codon usage bias are Relative synonymous codon usage (RSCU), Effective Number of Codon (ENC), Codon bias Index (CBI) and Intrinsic codon bias Index (ICDI). The parameters and algorithms for each index are described below:

### Relative Synonymous Codon Usage (RSCU)

The relative synonymous codon usage is the ratio of the observed frequency of a codon to the expected frequency of a codon if all the synonymous codons for a particular amino acid are used equally [1]. If the RSCU value of a codon is greater than one that the codon is frequently used than expected whereas if the RSCU value of a codon is less than one that the codon is less frequently used than expected. If the RSCU value of a codon is <0.6, it represents the under-represented while if the RSCU value of a codon >1.6, it represents over-represented. RSCU is calculated as:

$$RSCU_{ij} = \frac{X_{ij}}{\frac{1}{ni}\sum_{j=1}^{ni} X_{ij}}$$

Where,  $X_{ij}$  is the frequency of occurrence of the j<sup>th</sup> codon for i<sup>th</sup> amino acid (any  $X_{ij}$  with a value of zero is arbitrarily assigned a value of 0.5) and  $n_i$  is the number of codons for the i<sup>th</sup> amino acid (i<sup>th</sup> codon family).

## Effective Number of Codon (ENC)

ENC is a widely used measure of codon usage bias and its value is dependent upon the nucleotide composition of a gene. Its value ranges from 20 to 61 and higher value means lower codon usage bias and viceversa. If the value is 20, it signifies only one codon is used for each amino acid and if the value is 61 it denotes all the codons are equally likely to code for the same amino acid. ENC is measured as:

$$ENC = 2 + \frac{9}{F_2} + \frac{1}{F_3} + \frac{5}{F_4} + \frac{3}{F_6}$$

Where  $F_k$  (k=2, 3, 4, 6) is the mean of  $F_k$  values for the k-fold degenerate amino acids [2].

# Codon Bias Index (CBI)

The codon bias index is also the measure of codon usage bias, based on the degree of preferred codons used in a gene, like to the frequency of optimal codons. Its value ranges from 0 to 1. The 0 indicates random choice of codon whereas 1 signifies only the preferred codons used in a gene while less than zero denotes non-preferred codons used more in the gene [3]. CBI can be calculated as:

$$CBI = \frac{N_{opt} - N_{ran}}{N_{tot} - N_{ran}}$$

#### Intrinsic Codon Bias Index (ICDI)

The ICDI is a parameter of codon usage bias and it is not based on optimal codons. The value of ICDI ranges from 0 to 1 where 0 signify for equal usage and 1 for extremely high-biased genes. The ICDI value greater than 0.5 indicates high bias whereas its value lowers than 0.3 means little bias. The ICDI is computed based on  $S_a$  values for each of the 18 amino acids with k-fold degeneracy [4].

$$S_a = \frac{1}{K_a(K_a - 1)} \sum_{c \in C_a} (r_{ac} - 1)^2$$

Where  $r_{ac}$  is the relative synonymous codon usage and  $k_a$  is the degeneracy of amino acid in the sequence. The value of the index is then computed as:

$$ICDI = \sum_{a \in A} F_a S_a$$

The ICDI gives equal weight to all amino acids included, that is, all values of Fa are 1/18.

#### References

- Sharp PM, Li WH (1986) An evolutionary perspective on synonymous codon usage in unicellular organisms. J Mol Evol 24: 28-38.
- Wright F (1990) The 'effective number of codons' used in a gene. Gene 87: 23-29.
- Bennetzen JL, Hall B (1982) Codon selection in yeast. J Biol Chem 257: 3026-3031.
- Freire-Picos MA, Gonzalez-Siso MI, Rodriguez-Belmonte E, Rodriguez-Torres AM, Ramil E, et al. (1994) Codon usage in *Kluyveromyces lactis* and in yeast cytochrome c-encoding genes. Gene 139: 43-49.

\*Corresponding author: Uddin A, Department of Zoology, Moinul Hoque Choudhury Memorial Science College, Algapur, Hailakndi-788150, India, Tel: +91-9613554108; E-mail: arif.uddin29@gmail.com

Received June 28, 2017; Accepted June 29, 2017; Published June 30, 2017

Citation: Uddin A (2017) Indices of Codon Usage Bias. J Proteomics Bioinform 10: e34. doi: 10.4172/jpb.1000e34

**Copyright:** © 2017 Uddin A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.