Immunogenetics: Open Access

OPEN ACCESS Freely available online

Commentary

Inherited Cytokine Gene Variants and their Association with Individual Differences in Autoimmune Disease Susceptibility

Adrian Velasco*

Department of Molecular Immunology and Immunogenetics, Baltic School of Life Sciences, University of Riga, Riga, Latvia

DESCRIPTION

The human immune system relies on continuous communication between numerous cell populations that coordinate defense against infectious organisms while preserving healthy tissues. This communication occurs through signaling proteins known as cytokines, which regulate cellular activation, differentiation, migration, proliferation, and programmed cell death. The genes responsible for cytokine production exhibit inherited variation that influences the quantity, timing, and balance of these signaling molecules. Such genetic diversity contributes to differences in immune activity among individuals and has become an important subject in immunogenetics because of its relationship with autoimmune diseases.

Cytokines consist of several functional groups, including interleukins, interferons, tumor necrosis factors, transforming growth factors, and chemokines. Each family performs distinct biological functions while interacting through highly coordinated signaling networks. A temporary increase in cytokine production assists the elimination of infectious organisms and supports tissue repair following injury. However, prolonged production or inadequate regulation may result in continuous inflammation capable of damaging healthy organs.

Single nucleotide polymorphisms represent the most frequently studied form of inherited variation within cytokine genes. These alterations involve substitution of one nucleotide for another at a specific genomic position. Although many polymorphisms have little biological effect, others occur within promoter regions, coding sequences, untranslated regions, or regulatory elements that directly influence gene activity. Variants affecting transcription factor binding sites may increase or decrease cytokine synthesis following immune stimulation. Such differences become especially important when repeated inflammatory responses occur over many years.

Interleukin-6 serves as an informative example of cytokine gene variation. This cytokine participates in acute inflammatory responses, B-cell maturation, antibody production, and differentiation of helper T cells. Elevated interleukin-6 concentrations have been documented in several autoimmune diseases, including rheumatoid arthritis, systemic lupus erythematosus, juvenile idiopathic arthritis, and inflammatory bowel disorders. Certain inherited promoter variants influence the amount of interleukin-6 produced during immune activation. Individuals carrying high-expression variants may experience stronger inflammatory reactions, potentially increasing susceptibility to chronic autoimmune conditions under appropriate environmental circumstances.

Interleukin-10 illustrates the importance of anti-inflammatory cytokines in maintaining immune tolerance. Unlike pro-inflammatory mediators, interleukin-10 limits excessive immune activation by reducing cytokine production, suppressing antigen presentation, and regulating macrophage activity. Variants associated with reduced interleukin-10 production have been linked with increased inflammatory activity in several autoimmune diseases. Lower availability of this regulatory cytokine may permit prolonged activation of autoreactive lymphocytes, contributing to tissue injury during chronic inflammation.

Interleukin-17 has attracted increasing attention because of its relationship with chronic inflammatory disorders. Produced mainly by T helper 17 cells, this cytokine stimulates neutrophil recruitment and promotes inflammatory mediator production by epithelial cells, fibroblasts, and endothelial cells. Variants affecting pathways controlling interleukin-17 production have been associated with psoriasis, ankylosing spondylitis, multiple sclerosis, and several other immune-mediated disorders. These findings demonstrate how inherited regulation of specific lymphocyte populations contributes to disease susceptibility.

Cytokine genes rarely function independently. Instead, multiple inherited variants interact simultaneously to influence immune responses. One individual may possess several polymorphisms associated with increased production of inflammatory cytokines while carrying additional variants affecting regulatory pathways. The combined effect of these inherited characteristics

Correspondence to: Adrian Velasco, Department of Molecular Immunology and Immunogenetics, Baltic School of Life Sciences, University of Riga, Riga, Latvia, E-mail: adrian.velasco@bsls-riga.lv

Received: 24-Feb-2026, Manuscript No. IGOA-26-43862; **Editor assigned:** 26-Feb-2026, PreQC No. IGOA-26-43862(PQ); **Reviewed:** 12-Mar-2026, QC No. IGOA-26-43862; **Revised:** 19-Mar-2026, Manuscript No. IGOA-26-43862 (R); **Published:** 26-Mar-2026, DOI: 10.35248/IGOA.26.11.291

Citation: Velasco A (2026). Inherited Cytokine Gene Variants and their Association with Individual Differences in Autoimmune Disease Susceptibility. Immunogenet Open Access. 11:291.

Copyright: © 2026 Velasco A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

determines overall immune responsiveness more accurately than examination of any single genetic marker. Modern statistical methods therefore evaluate combinations of variants rather than isolated polymorphisms.

Gene-gene interaction represents an important consideration in immunogenetic analysis. Cytokine genes operate within extensive signaling pathways involving receptors, intracellular signaling molecules, transcription factors, and regulatory proteins. Variation within one component may influence the biological effect of variation elsewhere in the pathway. Consequently, comprehensive genomic studies increasingly examine networks of interacting genes instead of focusing upon individual loci.

Twin studies have provided valuable insight into these relationships. Monozygotic twins possess nearly identical genomic sequences, yet autoimmune diseases frequently occur in only one sibling. Such observations indicate that inherited cytokine variation increases susceptibility without independently determining disease occurrence. Environmental influences, epigenetic regulation, and random biological events contribute additional variability throughout development and adulthood.

Large genome-wide association studies have identified numerous susceptibility regions associated with autoimmune disorders. Many of these regions contain cytokine genes or molecules involved in cytokine signaling pathways. Advances in sequencing technology permit simultaneous analysis of millions of genetic markers across diverse populations, improving statistical confidence while identifying variants with relatively small individual effects. These studies demonstrate that autoimmune susceptibility usually results from the combined influence of numerous inherited variants rather than a single genetic alteration.

CONCLUSION

Future studies will likely combine whole-genome sequencing, transcriptomics, proteomics, metabolomics, and single-cell immune profiling to generate increasingly detailed descriptions of cytokine regulation across diverse populations. Such integrated approaches may identify additional inherited variants influencing immune tolerance and inflammatory control while improving understanding of the biological pathways associated with autoimmune disease susceptibility. Continued international collaboration and carefully designed population studies will remain essential for expanding knowledge within immunogenetics and for explaining why immune responses differ so substantially among individuals despite exposure to similar environmental conditions.