Journal of Cancer Research and Immuno-Oncology

OPEN ACCESS Freely available online

Opinion Article

Innate Lymphoid Cell Functions Shaping Antitumor Immunity across Human Cancer Microenvironments

Natalia Sørensen*

Department of Translational Cancer Immunology, Baltic International University, Copenhagen, Denmark

DESCRIPTION

Cancer develops through continuous interactions among malignant cells, stromal populations, vascular networks, extracellular components, and numerous immune cell subsets that collectively influence disease progression. While considerable attention has traditionally focused on T lymphocytes, B lymphocytes, macrophages, and dendritic cells, increasing scientific interest has been directed toward innate lymphoid cells because of their diverse contributions to tissue immunity and tumor biology. These immune populations do not possess antigen-specific receptors generated through somatic recombination, yet they rapidly respond to environmental signals and coordinate local immune activity through cytokine secretion and communication with neighboring cells. Their activities influence inflammatory regulation, epithelial integrity, tissue repair, vascular organization, and interactions with adaptive immunity. Within cancer immuno-oncology, investigation of innate lymphoid cells has expanded considerably as researchers seek to understand how these populations affect anti-tumor responses across different forms of human malignancy.

Innate lymphoid cells originate from common lymphoid progenitor cells within the bone marrow and subsequently differentiate into several functionally distinct populations. Although these cells lack rearranged antigen receptors, they share developmental relationships with adaptive lymphocytes while displaying characteristics commonly associated with innate immunity. Their rapid responsiveness enables immediate reactions to alterations occurring within tissue microenvironments before adaptive immune responses become fully established. These properties make innate lymphoid cells important contributors to early immune surveillance within healthy tissues and malignant lesions.

Several biologically distinct innate lymphoid cell populations have been identified based on transcriptional programs, cytokine production, and tissue localization. Individual subsets respond to different combinations of cytokines released by epithelial cells, stromal populations, dendritic cells, and macrophages. Rather than functioning independently, these populations communicate extensively with surrounding immune and non-immune cells, producing highly coordinated responses that vary according to anatomical location and pathological conditions. This functional diversity allows innate lymphoid cells to participate in numerous physiological and pathological processes throughout the body.

Within healthy tissues, innate lymphoid cells contribute to maintenance of epithelial integrity and regulation of local immune equilibrium. Mucosal surfaces lining the gastrointestinal tract, respiratory system, and genitourinary organs contain abundant innate lymphoid cell populations because these tissues remain continuously exposed to microorganisms and environmental stimuli. These cells respond rapidly to epithelial injury by producing cytokines that influence tissue repair, antimicrobial defense, and communication with additional immune populations. Their capacity to maintain tissue stability illustrates their importance beyond conventional antimicrobial immunity.

Malignant transformation substantially modifies the local environment encountered by innate lymphoid cells. Cancer cells continuously release cytokines, chemokines, metabolites, extracellular vesicles, and growth factors that alter immune composition within tumors. Fibroblasts, endothelial cells, macrophages, dendritic cells, and extracellular matrix components provide additional signals influencing innate lymphoid cell behavior. Continuous exposure to these changing molecular environments results in considerable functional diversity among innate lymphoid cell populations located within malignant tissues. Modern molecular analyses demonstrate that these cells adapt dynamically according to local biological conditions rather than maintaining fixed functional identities.

Natural killer cells share developmental similarities with innate lymphoid cells but possess distinct biological characteristics related to direct elimination of abnormal cells. Although classified separately, these populations cooperate extensively

Correspondence to: Natalia Sørensen, Department of Translational Cancer Immunology, Baltic International University, Copenhagen, Denmark, E-mail : natalia.sorensen@biu-denmark.edu

Received: 27-Feb-2026, Manuscript No. JCRIO-26-43983; **Editor assigned:** 02-Mar-2026, PreQC No. JCRIO-26-43983(PQ); **Reviewed:** 16-Mar-2026, QC No. JCRIO-26-43983; **Revised:** 23-Mar-2026, Manuscript No. JCRIO-26-43983 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2684-1266.26.12.282

Citation: Sørensen N (2026). Innate Lymphoid Cell Functions Shaping Antitumor Immunity across Human Cancer Microenvironments. J Cancer Res Immunooncol. 11:282.

Copyright: © 2026 Sørensen N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

within malignant tissues. Cytokines released by innate lymphoid cells influence natural killer cell recruitment, activation, and survival. Conversely, natural killer cells produce mediators that regulate neighboring innate lymphoid cell populations. Coordinated activity between these immune subsets strengthens local immune surveillance and contributes to early recognition of transformed cells.

The intestinal microbiome also influences innate lymphoid cell biology through production of metabolites and microbial products entering systemic circulation. Short-chain fatty acids, bile acid derivatives, and additional microbial compounds regulate cytokine production and differentiation of multiple immune populations, including innate lymphoid cells. Differences in microbial composition among individuals may therefore contribute to variation in immune responses observed during cancer progression and treatment. Continued investigation seeks to clarify these relationships across diverse malignancies.

Immunotherapeutic strategies have further stimulated interest in innate lymphoid cells because successful treatment depends upon coordinated interactions among multiple immune populations. While many current therapies primarily target adaptive immunity, accumulating evidence suggests that innate immune regulation significantly influences therapeutic effectiveness. Communication between innate lymphoid cells, dendritic cells, macrophages, natural killer cells, and cytotoxic T lymphocytes contributes to overall immune competence within tumors. Understanding these interactions may improve interpretation of patient-specific treatment responses and guide future therapeutic development.

Liquid biopsy technologies offer additional opportunities for monitoring biological activity associated with innate lymphoid cells. Blood samples containing extracellular vesicles, circulating cytokines, soluble receptors, and nucleic acids may provide complementary information regarding immune regulation without repeated tissue biopsies. Longitudinal assessment during therapy enables investigators to examine evolving immune communication throughout disease management while reducing procedural burden for patients.

CONCLUSION

Future progress in cancer immuno-oncology will increasingly depend upon comprehensive understanding of innate lymphoid cell biology and its integration with broader immune communication networks. These cells participate in tissue surveillance, inflammatory regulation, epithelial maintenance, interaction with adaptive immunity, and coordination of stromal responses throughout malignant tissues. Their remarkable functional diversity reflects continuous adaptation to changing biological environments created by malignant cells and surrounding stromal populations. Continued investigation combining molecular biology, advanced imaging, computational analysis, and clinical research will strengthen current understanding of innate lymphoid cell functions while supporting development of improved biomarkers, refined patient classification methods, and innovative therapeutic strategies directed toward enhancing anti-tumor immunity across a wide spectrum of human cancers.