

Interplay Between Oxidative Phosphorylation Efficiency and Early Embryo Cleavage Kinetics in Human IVF Culture Systems

Sabrina Laurent*

Department of Reproductive Cellular Bioenergetics, Alpine Institute of Medical Sciences, Grenoble, France

DESCRIPTION

Early embryonic development in assisted reproductive treatment is characterized by rapid cell divisions that require precise coordination of energy production and cellular signaling. Among the multiple biological factors influencing embryo viability, mitochondrial function and oxidative phosphorylation efficiency have gained increasing attention. Oxidative phosphorylation represents the primary pathway for adenosine triphosphate production in early embryos, particularly during cleavage stages when energy demand increases substantially. Variability in mitochondrial efficiency within oocytes and early embryos has been associated with differences in cleavage timing, morphological development, and blastocyst formation rates in in vitro fertilization systems.

Cleavage kinetics refers to the timing and pattern of early embryonic cell divisions. Embryologists often assess the intervals between successive cell divisions to evaluate embryo quality in culture. Embryos with synchronized and timely cleavage are generally considered to have higher developmental potential compared with embryos exhibiting irregular or delayed division patterns. Although morphological grading systems remain widely used, time-lapse imaging technology has enabled more detailed analysis of cleavage dynamics and their relationship with metabolic function.

Mitochondrial oxidative phosphorylation depends on the coordinated activity of protein complexes embedded in the inner mitochondrial membrane. These complexes facilitate electron transfer and proton gradient formation, ultimately driving adenosine triphosphate synthesis. Disruption of any component within this system may reduce energy output and increase production of reactive oxygen species. Excess reactive oxygen species can damage Deoxyribonucleic Acid (DNA), proteins, and lipids, thereby affecting embryonic development and cellular viability. Oocyte quality plays a central role in determining mitochondrial efficiency in early embryos. Since mitochondria are maternally inherited, their number, structure, and functional capacity are established during oogenesis. Women with diminished ovarian reserve or advanced reproductive age often

produce oocytes with altered mitochondrial DNA copy number and reduced membrane potential. These abnormalities may contribute to impaired oxidative phosphorylation during early embryogenesis, potentially influencing cleavage kinetics after fertilization.

Culture environment conditions in in vitro fertilization laboratories also affect embryonic energy metabolism. Oxygen concentration, temperature stability, pH buffering, and nutrient composition of culture media all influence mitochondrial activity. Embryos cultured under atmospheric oxygen conditions may experience increased oxidative stress compared with those cultured under reduced oxygen levels that more closely resemble physiological conditions within the reproductive tract. Elevated oxidative stress may impair mitochondrial function and alter cleavage timing. Glucose and pyruvate metabolism represent important energy pathways supporting oxidative phosphorylation in early embryos. Pyruvate is often the preferred substrate during early cleavage stages, while glucose utilization increases during later development. Imbalances in substrate availability may influence mitochondrial efficiency and energy production. Culture media formulations are therefore designed to support metabolic transitions throughout embryonic development. Variations in metabolic substrate composition may contribute to differences in cleavage kinetics observed between laboratory protocols.

Reactive oxygen species production is a natural byproduct of oxidative phosphorylation. However, excessive accumulation of these molecules can lead to oxidative damage and mitochondrial dysfunction. Embryos possess antioxidant defense systems including superoxide dismutase, glutathione peroxidase, and catalase to mitigate oxidative stress. If these defense mechanisms are insufficient, cleavage progression may be delayed or arrested due to cellular damage. Some studies have suggested that embryos with higher antioxidant capacity exhibit more consistent cleavage timing and improved developmental outcomes. Time-lapse incubators have significantly advanced the study of cleavage kinetics by enabling continuous observation of embryo development without repeated exposure to environmental fluctuations. These systems allow precise

Correspondence to: Sabrina Laurent, Department of Reproductive Cellular Bioenergetics, Alpine Institute of Medical Sciences, Grenoble, France, E-mail: sabrina.laurent.aims@mitolabmail.org

Received: 02-Jun-2025, Manuscript No. JFIV-25-41845; **Editor assigned:** 04-Jun-2025, PreQC No. JFIV-25-41845 (PQ); **Reviewed:** 18-Jun-2025, QC No. JFIV-25-41845; **Revised:** 24-Jun-2025, Manuscript No. JFIV-25-41845 (R); **Published:** 02-Jul-2025, DOI: 10.35841/2375-4508.25.13.419

Citation: Laurent S (2025). Interplay Between Oxidative Phosphorylation Efficiency and Early Embryo Cleavage Kinetics in Human IVF Culture Systems. *J Fertil In Vitro IVF World w Reprod Med Gent Stem Cell Biol.* 13:419.

Copyright: © 2025 Laurent S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

measurement of cell division intervals and morphological changes. Researchers have identified associations between specific cleavage patterns and implantation success, suggesting that metabolic efficiency may be reflected in early developmental timing. Embryos with rapid and synchronous cleavage often demonstrate higher blastocyst formation rates.

Mitochondrial DNA integrity is another factor influencing oxidative phosphorylation efficiency. Damage to mitochondrial DNA can impair electron transport chain function and reduce adenosine triphosphate production. Such damage may arise from oxidative stress, environmental exposure, or intrinsic genetic instability. Embryos with compromised mitochondrial DNA may exhibit slower cleavage progression and reduced viability during extended culture. Assisted reproductive technologies involving extended embryo culture and blastocyst selection may indirectly favor embryos with more efficient mitochondrial function. Embryos that successfully reach the blastocyst stage often demonstrate stable energy metabolism and

coordinated cleavage patterns. However, some embryos with slower early cleavage may still achieve successful implantation, indicating that metabolic efficiency is only one component of developmental competence.

CONCLUSION

Current evidence indicates that oxidative phosphorylation efficiency plays a significant role in determining early embryo cleavage kinetics during fertilization culture systems. Mitochondrial function, substrate availability, oxidative stress regulation, and culture conditions collectively influence embryonic energy metabolism. Continued research combining reproductive biology, bioenergetics, and embryology may improve understanding of early developmental processes and support optimization of assisted reproductive laboratory practices.