

Investigation of Ginsenoside Rb₁ from *Acanthopanax koreanum* by Eastern Blotting and ELISA Analyses

Nguyen Huu Tung^{1,3}, Takuhiro Uto¹, Susumu Isoda² and Yukihiro Shoyama^{1*}

¹Faculty of Pharmaceutical Science, Nagasaki International University, 2825-7 Huis Ten Bosch, Sasebo, Nagasaki 859-3298, Japan

²Faculty of Pharmaceutical Science, Showa University, 1-5-8 Hatanodai, Shinagawa, Tokyo 142-8555, Japan

³School of Medicine and Pharmacy, Vietnam National University, Hanoi, 144 Xuan Thuy St., Cau Giay, Hanoi, Vietnam

Abstract

Ginsenosides such as ginsenoside Rb₁ are the principal components in *Panax* spp. and have been found in some other Araliaceae plants. In our survey of ginsenoside from Araliaceous species based on eastern blotting and ELISA methods using anti-ginsenoside Rb₁ monoclonal antibody, ginsenoside Rb₁, one of the principal ginseng saponins, was investigated with concentrations of 0.000016, 0.000014, and 0.000039% dry weight in the leaves, stem, and roots of the well-known medicinal plant *Acanthopanax koreanum* respectively. The obtained results further support potential and promising application of MAb including eastern blotting and ELISA for surveying ginsenoside sources including ginsenoside Rb₁.

Keywords: *Acanthopanax koreanum*; Araliaceae; Ginsenoside; Ginsenoside Rb₁; Monoclonal antibody; Eastern blotting

Introduction

Acanthopanax koreanum which is a shrub belonging to the Araliaceae family and is distributed in Northeast Asia, has been used as a tonic and to treat rheumatism, allergies, hepatitis, and diabetes [1,2]. Phytochemical profile of *A. koreanum* have been documented with several lignans and diterpenes [3,4] along with, especially, various lupane-type triterpene saponins, which are considered as major constituents [5,6].

In the case of qualitative and/or quantitative analytical approaches of ginsenosides, thin layer chromatography (TLC) [7], high performance liquid chromatography (HPLC) [8,9], and liquid chromatography-mass spectrometry (LC-MS) [10] are used frequently. Recently, it becomes evident that an enzyme-linked immunosorbent assay (ELISA) using monoclonal antibody (MAb) against small molecular has been developed for natural products as a highly sensitive, specific, and simple methodology. In this regard, we have succeeded in the preparation of MAb against ginsenoside Rb₁ (G-Rb₁) [11], ginsenoside Rg₁ [12], and G-Re [13], and established a new immunostaining method, eastern blotting for G-Rb₁ and -Rg₁ [14]. Additionally, in our previous study, the combination of ELISA and eastern blotting methods using anti-G-Rb₁ MAb was applied for the identification of G-Rb₁ in *Panax* species and traditional Chinese medicines containing lower concentrations of G-Rb₁ [15]. This paper herein deals with the finding and determination of G-Rb₁ in *A. koreanum* of Araliaceae family by using ELISA and eastern blotting monitoring.

Materials and Methods

Materials

Polyethersulfone (PES) membranes (Immobilon-N) and glass microfiber filter sheets were from Millipore Corporation (Bedford, MA, USA), and bovine (BSA) and human (HSA) serum albumins were from Pierce (Rockford, IL, USA). Peroxidase-labeled anti-mouse IgG was from Organon Teknika Cappel Products (West Chester, PA, USA). Standard G-Rb₁, -Rc, -Rd, -Re and -Rg₁ were from Wako Pure Chemical Industries, Ltd (Osaka, Japan). All chemicals and solvents used were of analytical grade and water was filtered by a Millipore ultrapure water system (Milli-Q Direct, Merck Millipore, Germany).

Sample preparation

A. koreanum samples were collected by Dr. Susumu Isoda from Herbal Garden in Faculty of Pharmaceutical Science, Showa University, and authenticated by one of the authors (YS). Dried samples of the plant (roots, stems, and leaves) (50 mg individually) were powdered then extracted with methanol (5 mL) under sonication five times. They were then filtered and the combined extract was diluted with 20% methanol for ELISA and eastern blotting.

Eastern blotting and double staining

After adaptation with procedures from Tanaka et al. [15], *A. koreanum* extracts were loaded onto two TLC plates and developed with *n*-BuOH-EtOAc-H₂O (15:1:4). Of which, one developed TLC plate was dried and stained with H₂SO₄ and another plate was dried followed by spraying with a blotting solution mixture of isopropanol-methanol-H₂O (1:4:16, v/v/v). Next, the treated TLC plate was placed on a stainless steel plate then covered with a PES membrane sheet. After covering with a glass microfiber filter sheet, it was pressed evenly for 70 s with a 120°C hot. The PES membrane was separated from the plate and dried then dipped in water containing NaIO₄ (10 mg/mL) while stirring at room temperature for 1 h. After washing with water, 50 mM carbonate buffer solution containing BSA was added followed by stirring for 5 h. The PES membrane was then washed twice with PBS containing 0.05% Tween-20 (TPBS) for 5 min then washed with water. It was then immersed in anti-G-Rb₁ MAb and shaken at 4°C overnight. In the next day, the membrane was washed twice with TPBS-water and 1000 times diluted of peroxidase-labeled goat anti-mouse IgG in PBS containing 0.2% gelatin (GPBS) was then added followed by stirring at room temperature for 1 h. The obtained membrane was then rinsed twice with TPBS and water, and treated with 1 mg/mL 4-chloro-1-naphthol-0.03% H₂O₂ in PBS solution freshly prepared before use for

***Corresponding author:** Yukihiro Shoyama PhD, Professor, Faculty of Pharmaceutical Sciences, Nagasaki International University, Nagasaki 859-3298, Japan, Tel: +81956205653; Fax: +81956205653; E-mail: shoyama@niu.ac.jp

Received July 13, 2016; Accepted July 26, 2016; Published July 29, 2016

Citation: Tung NH, Uto T, Isoda S, Shoyama Y (2016) Investigation of Ginsenoside Rb₁ from *Acanthopanax koreanum* by Eastern Blotting and ELISA Analyses. Pharm Anal Chem Open Access 2: 116. doi: [10.4172/2471-2698.1000116](https://doi.org/10.4172/2471-2698.1000116)

Copyright: © 2016 Tung NH, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

10 min at room temperature.

For the staining by anti-G-Rg₁ MAb, the blotted PES membrane was treated in the same procedure as anti-G-Rb₁ MAb except that it was exposed to 2 mg/10 mL 3-amino-9-ethylcarbazole-0.03% H₂O₂ in acetate buffer (0.05 M, pH 5.0) containing 0.5 mL of *N,N*-dimethylformamide.

Competitive ELISA

G-Rb₁ concentrations in *A. koreanum* were analyzed by ELISA [14]. G-Rb₁-HSA (100 μL of 1 μg/mL) was adsorbed onto the wells of a 96-well immunoplate and treated with 300 μL PBS containing 5% skimmed milk (S-PBS) for 1 h to reduce non-specific adsorption. Different concentrations of G-Rb₁ (50 μL) or samples diluted in 20% MeOH were incubated with 50 μL G-Rb₁ MAb solution for 1 h. The sample plates were washed three times with T-PBS and incubated with 100 μL of a 1000-fold dilution of peroxidase-labeled goat anti-mouse IgG for 1 h. Next, the plates were washed again plates with T-PBS and then 100 μL substrate solution (100 mM citrate buffer, pH 4.0, containing 0.003% H₂O₂ and 0.3 mg/mL of 2,2-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS; Wako Pure Chemical, Osaka, Japan) was added to each well followed by incubation for 15 min. The absorbance was recorded on micro-plate reader (Immuno Mini NJ-2300, Nalge Nunc, Roskilde, Denmark) at 405 nm. All reactions were carried out at 37°C.

Results and Discussion

While PES membranes treated with NaIO₄ solution and reactive BSA protein enhanced the fixation of ginsenoside-BSA conjugation, small cross-reactivities against anti-G-Rb₁ MAb like G-Rc (0.024%) and G-Rd (0.02%) were also observed [11]. It is noticeable that only G-Rb₁, G-Rc and G-Rd having protopanaxadiol as an aglycone could be stained and this became evident that the specific reactivity of the sugar part in the ginsenoside molecule against MAb might be modified by NaIO₄ treatment of the ginseng saponin on the membrane. This phenomenon is important for the surveys of saponins having the same aglycone like ginsenosides. Furthermore, we used two MAbs, anti-G-Rb₁ MAb and anti-G-Rg₁ MAb for eastern blotting as previously reported [16]. In this case, two types of ginsenoside having protopanaxadiol and protopanaxatriol as an aglycone can be stained separately. Purple color and blue color indicate ginsenosides having protopanaxatriol and protopanaxadiol as aglycone, respectively as indicated in previous paper [16].

As shown in Figure 1, although H₂SO₄ staining detected clearly all standard ginsenosides without changing color, the TLC profile of *A. koreanum* crude extract revealed complicated spot patterns indicating that ginsenosides are ambiguously determined. However, the evidence of double eastern blotting unambiguously indicated that no ginsenoside having protopanaxatriol as an aglycone was detected in the *A. koreanum* crude extract because of no purple spot appeared. On the other hand, blue spots were observed meaning that protopanaxadiol type ginsenosides are occurred in *A. koreanum*. In addition, it is clear that R_f value on TLC reflects the sugar number in general. Taken together, based on these evidences, it is suggested that *A. koreanum* contains small amount of G-Rb₁ and a more polar ginsenoside which cannot be longer analyzed due to its trace amount (Figure 1).

Next, we analyzed *A. koreanum* leaves crude extract by competitive ELISA using anti-G-Rb₁ MAb (Figure 2) in order to confirm the existence and concentration of G-Rb₁ resulting in 0.000016% dry wt. of G-Rb₁. The roots and stems were also analyzed separately by the same




Figure 1: (A) Eastern blotting profile for the standard ginsenosides and MeOH extract of *A. koreanum* (leaf) (B) TLC plate treated with 10% H₂SO₄.




Figure 2: Calibration curve for G-Rb₁, various concentrations of G-Rb₁ were incubated with MAb in wells pre-coated with G-Rb₁ HSA (1 μg/mL). After washing with PBS containing 0.05% of Tween 20, the wells were incubated again with peroxidase-labeled anti-mouse IgG. Absorbance was measured at 405 nm.

manner finding concentrations of 0.000039% and 0.000014% dry wt., respectively. In the view point of naturally occurring compounds, the concentrations of G-Rb₁ in *A. koreanum* is extremely low, therefore, it has become evident that chromatographic purification and analyses of G-Rb₁ have been unaffordable to date. To the best of our knowledge, occurrence of G-Rb₁ and dammarane-type saponins in the *Acanthopanax* genus had not been reported previously.

In conclusion, it is found that *A. koreanum* might be a new resource of G-Rb₁ and more analytical studies are suggested to accumulate G-Rb₁ from the medicinal materials. This study further support potential and

promising application of MAb such as eastern blotting and ELISA for surveying ginsenoside sources.

Acknowledgements

This work was supported by a Grant-in-Aid from the Japan Society for the Promotion of Science (JSPS). One of the authors (N.H.T) is grateful to the JSPS for a Postdoctoral Research Fellowship at Nagasaki International University.

References

1. Lee YN (1996) Flora of Korea. Kyohaksa, Seoul, pp: 1107-1109.
2. Perry LM, Metzger J (1980) Medicinal plants of East and Southeast Asia. MIT Press, Cambridge, London.
3. Cai XF, Shen G, Dai NT, Kang OH, Kim JA, et al. (2003) Inhibitory effect of TNF- α and IL-8 secretion by pimarane type diterpenoids from *Acanthopanax koreanum*. Chem Pharm Bull 51: 605-607.
4. Cai XF, Lee IS, Dai NT, Shen G, Kang JS, et al. (2004) Inhibitory lignans against NFAT transcription factor from *Acanthopanax koreanum*. Arch Pharm Res 27: 738-741.
5. Choi HS, Kim HJ, Nam SG, Kim IS, Lee KT, et al. (2008) Lupane glycosides from the leaves of *Acanthopanax koreanum*. Chem Pharm Bull 56: 1613-1616.
6. Nhiem NX, Tung NH, Kiem PV, Minh CV, Ding Y, et al. (2009) Lupane triterpene glycosides from leaves of *Acanthopanax koreanum* and their cytotoxic activity. Chem Pharm Bull 57: 986-989.
7. Kitagawa I, Taniyama T, Shibuya H, Noda T, Yoshikawa M (1987) Chemical studies on crude drug processing. V. On the constituents of ginseng radix rubra (2): comparison of the constituents of white ginseng and red ginseng prepared from the same *Panax ginseng* root. Yakugaku Zasshi 107: 495-505.
8. Chan TWD, But PPH, Cheng SW, Kwok IMY, Lau FW, et al. (2000) Differentiation and authentication of *Panax ginseng*, *Panax quinquefolius*, and ginseng products by using HPLC/MS. Anal Chem 72: 1281-1287.
9. Yamaguchi H, Kasai R, Matsuura H, Tanaka O, Fuwa T (1988) High-performance liquid chromatography analysis of acidic saponins of ginseng and related plants. Chem Pharm Bull 36: 3468-3473.
10. Wang X, Sakuma T, Asafu-Adjaya E, Shiu GK (1999) Determination of ginsenosides in plant extracts from *Panax ginseng* and *Panax quinquefolius* L. by LC/MS/MS. Anal Chem 71: 1579-1584.
11. Tanaka H, Fukuda N, Shoyama Y (1999) Formation of monoclonal antibody against a major ginseng component, ginsenoside Rb1 and its characterization. Cytotechnology 29: 115-120.
12. Fukuda N, Tanaka H, Shoyama Y (2000) Formation of monoclonal antibody against a major ginseng component, ginsenoside Rg₁ and its characterization. Monoclonal antibody for a ginseng saponin. Cytotechnology 34: 197-204.
13. Morinaga O, Takuhiro U, Yuan CS, Tanaka H, Shoyama Y (2010) Evaluation of a new eastern blotting technique for the analysis of ginsenoside Re in American ginseng berry pulp extracts. Fitoterapia 81: 284-288.
14. Fukuda N, Tanaka H, Shoyama Y (2000) Application of ELISA, Western blotting and immunoaffinity concentration for survey of ginsenosides in crude drugs of *Panax* species and traditional Chinese herbal medicines. Analyst 125: 1425-1429.
15. Tanaka H, Fukuda N, Shoyama Y (2007) Eastern blotting and immunoaffinity concentration using monoclonal antibody for ginseng saponins in the field of traditional Chinese medicines. J Agric Food Chem 55: 3783-3787.
16. Fukuda N, Tanaka H, Shoyama Y (2001) Double staining of ginsenosides by western blotting using anti-ginsenoside Rb₁ and Rg₁ monoclonal antibodies. Biol Pharm Bull 24: 1157-1160.