

## Isolation and quantification of nanovesicles in isoproterenol-induced myocardial infarcted rats

Aviwe Ntsethe

### Abstract

Myocardial infarction (MI) is one of the main sources of death around the world. The pathogenesis and etiology of MI is as yet indistinct. Cardiovascular troponin is the solitary known heart explicit marker for the determination of MI yet because of the postponed arrival of troponin in the dissemination, a novel cardiovascular biomarker is required in the beginning phases of improvement of MI to lessen MI mortality. Exosomes are accounted for to be exceptionally managed by pressure. In this manner we surveyed the theory that exosome emission is expanded after MI and subsequently fill in as biomarker for MI. The point of this examination was to evaluate exosomes in an isoproterenol (ISO)- actuated MI rodents. Twelve rodents were utilized in this investigation, Group-A (n=6) was the typical control rodents and Group-B (n=6) was the ISO-treated gathering. Group B creatures were infused with isoproterenol for two successive days to prompt MI. Circulatory strain, pulse and body weight were checked in all creatures earlier the ISO infusion and all through the test. After the second ISO-infusion, all creatures were forfeited and blood, heart tissues were gotten. Histopathological examination was acted in heart tissue tests and levels of cardiovascular markers were estimated from the serum. Exosomes were disengaged from the plasma and evaluated by differential ultracentrifugation, nanoparticle following examination, transmission electron magnifying lens and ELISA separately. MI was affirmed by the expansion in BP and heart markers in ISO-induced rats. The centralization of exosomes was raised in plasma of ISO-treated animals. The investigation uncovered that exosomes are likely biomarkers of myocardial dead tissue.

**Keywords:** Isoproterenol, Myocardial infarction, Exosome, Nanoparticle tracking analysis, Cardiac markers

### INTRODUCTION

Cardiovascular diseases (CVD) and their thrombotic complications are the leading cause of morbidity and mortality in the world and mostly in developed countries. The World Health Organisation (WHO) estimated that approximately 17.3 million people die from CVD and its complications worldwide and in African nations CVD constitute 8.8% of all deaths. MI is one of the complications of CVD. MI is commonly known as a heart attack and occurs when there is an imbalance in the oxygen supply and demand, which is caused by injury to the coronary artery, blocking blood flow to the heart thus myocardial cell death. Early diagnosis of myocardial infarcted patients is required to prevent or reduce ischemic injury to the myocardium and subsequently prevent cardiac remodelling. Creatine kinase (CK) and lactate dehydrogenase (LDH) were previously used for diagnosis of myocardial infarcted patients. However, these markers are not cardiac specific markers as their levels increase in the circulation as result of various tissue damage. Thus, they were replaced with cardiac troponin T and troponin I which are cardiac specific markers, these biomarkers have led to early diagnosis of MI. However, there is a relative delay of the release of troponin as in AMI patients, troponin levels raise around 3.5 hours after the onset of chest pain. Earlier biomarkers with high specificity and sensitivity are in high demands to reduce MI mortality. In addition additional alternative markers are needed to reduce MI mortality. Recent studies have focused on exosomes as both potential biomarkers and therapeutic agent. Exosomes are membrane vesicles that differ from others by their size (30-140nm), density and specific composition of molecules. These are the smallest extracellular vesicles,

originating from cellular endosomes and are generated in endosomal vesicles called multivesicular bodies. Exosomes have emerged as important biological signalling entities because they act as a vehicle to transport important molecules such as proteins, lipids and nucleic acids to distant cells. They are secreted by multiple cell types, including smooth muscle cells, endothelial cells, cardiomyocytes and stem cells. The exosomes have unique protein markers which include CD9, CD63 and CD81 amongst others and are important in differentiating exosomes from other extracellular vesicles. Exosomes can be found in most body fluids such as saliva, blood and urine since they are secreted by multiple cells. Studies have shown that exosomal content is regulated by stress and disease conditions. Although research on cardiac exosomes is new, a few publications provide strong evidence that exosomes can exert pathological effects during the cardiac response to stress and different myocardial diseases.

A recent study reported that peripartum cardiomyopathy (PPCM) result to the release of miR-146-enriched exosomes by endothelial cells. These exosomes can be used as a vehicle by cardiomyocytes where miR-146a impedes with the physiological metabolism and contractile ability of the cell, resulting in hypertrophy. Moreover, levels of exosomal miR-146a are higher in plasma of acute PPCM patients as compared to patients with dilated cardiomyopathy and healthy postpartum controls. Furthermore, heart failure therapy in PPCM patients shows decreased circulating exosomal miR-146a to normal levels, suggesting that miR-146a is a strong potential biomarker for diagnosis PPCM. The type of stress to which myocardial tissue is exposed may determine the level of exosome secretion. During MI where the heart is exposed to ischemic stress signals such as inflammation, hypoxia and injury, cardiomyocytes may elevate the secretion of exosomes. miR-1 together with miR-208 are increased in the urine of acute MI patients suggesting that circulating miRNAs released from the injured myocardium can go to distant organs through exosomes as they are stable and protected from degradation by RNases present in the different body fluids. Exosomes secreted by cardiomyocyte transport miRNAs The present study aimed to

isolate and quantify exosomes in isoproterenol-induced MI rats. In this study we assessed the hypothesis that exosome secretion by the injured cardiomyocytes is increased during MI and thereby serve as potential biomarker for MI. We used isoproterenol-induced myocardial infarcted Wistar rats model to quantify exosomes in MI rats. We monitored blood pressure and cardiac marker's level to confirm the development of MI.

## MATERIALS AND METHODS

### Ethics

UKZN Animal Ethics Sub-committee (Clearance Number 082/016PD) approved the experimental protocol used in this study.

### Experimental Animals

All the experiments were carried out with male albino Wistar rats weighing 250-300g, obtained from Biomedical Research Unit, University of KwaZulu-Natal, Westville, Durban, South Africa. They were housed in polypropylene cages (47cm×34cm×20cm) lined with husk and renewed every 24h under a 12:12h light/dark cycle at around 22°C. The rats had free access to tap water and food. The rats were fed on a standard pellet diet. The experiment was carried out according to the Guidelines of the Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Durban, South Africa.

### Drug and Chemicals

Isoproterenol (ISO) was purchased from Sigma Chemical Co., St. Louis, MO, USA. All other chemicals used were of analytical grade.

### Experimental Design

Twelve animals were used in this experiment. They were grouped into two groups, group A (n=6) were the normal control rats and group B (n=6) were the

test (MI) rats. The body weight and blood pressure of the both groups were measured every day. On the 8th day and 9th day Isoproterenol (85mg/kg body weight) was dissolved in ice-cold saline and injected subcutaneously into group B at an interval of 24h for two consecutive days to induce MI. At the end of the experimental period, after 12h of second isoproterenol injection (i.e. on 9th day), all the rats were anesthetized with the high dose of Isofar and were then sacrificed by cervical decapitation. Blood was collected in dry test tubes without anticoagulant for serum and plasma. Heart tissues were excised immediately, rinsed with saline and tissue homogenates were prepared phosphate saline buffer for the estimations/assays of various biochemical parameters. All the enzyme assays were done immediately.

## Estimation of Cardiac Markers

The level of serum cardiac troponin-T, CK and CK-MB and LDH were estimated by electrochemiluminescence immunoassay using a standard kit (Roche Diagnostics, Switzerland).

## Histopathological Examination

Immediately after the heart tissues were dissected from experimental animals, they were washed with saline and cut across the left ventricle into two slices. The heart tissue slices were then fixed in 10% buffered formalin and were embedded in paraffin wax, sectioned at 5µm and stained with haematoxylin and eosin (H&E). After fixation, the heart tissues were dehydrated by immersion into 70% ethanol overnight, followed by 90%, 100% and 100% for an hour. The tissues were then immersed into xylene two times for an hour. Finally, the tissues were put in molten wax (55-60°C) and moved into the pre-heated oven overnight to solidify before sectioning. The heart tissues were sectioned and mounted onto the slides. The slides were cleared using xylene solution and then dehydrated using ethanol (100, 90, 70 and 50%) for 2 minutes in each percentage. The slides were rehydrated using

distilled water until ready for staining. The slides were stained with haematoxylin and rinsed off with distilled water, and stained again with eosin for 2-3 minutes. The residual eosin was rinsed-off with distilled water and slides were dipped into 90% ethanol to remove water. The slides were finally immersed into xylene solution and then left out to dry. They were covered with the coverslips and left overnight. Finally, they were examined under high power microscope (100X) and photomicrographs were taken.

## Exosome Isolation and Quantification

Exosomes were isolated according to the method as described. Plasma (1ml) was diluted with an equal volume of phosphate buffered saline (PBS; pH 7.4). Exosomes were isolated and purified by differential ultracentrifugation using a 30% sucrose cushion. In brief, to remove cells, centrifugation was initially performed at 2000Xg at 4°C for 30 min, followed by 12 000Xg at 4°C for 45 min. To remove the remaining debris, the supernatant was transferred to other centrifuge tubes and centrifuged at 110 000Xg at 4°C for 120 min (Optima™ MAX-XP Ultracentrifuge, fixed angle MLA-55 rotor, Beckman Coulter Inc., Brea, CA, USA). To remove particles that are bigger than 200nm, the pellet was suspended in 2ml of PBS and filtered through a 0.20µm pore filter (Cellulose acetate, GVS™, Europe). The filtrate was centrifuged at 110 000Xg at 4°C for 70 min, the pellet was re-suspended in 2ml PBS (pH 7.4) and centrifuged at 110 000Xg for 70 min at 4°C. To purify the exosomes, the exosome pellet was suspended in 2ml of PBS and subsequently purified using a 30% sucrose cushion and centrifuged at 110 000Xg at 4°C for 75 min. The final pellet was re-suspended in 200µl of PBS and stored at -80°C. The concentration of total exosomes in the circulation was determined by the quantification of total immune-reactive CD63 enzyme-linked immune absorbency assay (ExoELISA™, System Biosciences, Mountain View, CA), as described by Salomon et al. (2014). CD63 is not an exosome specific marker but is

commonly bound to the exosomal membrane and hence the method employed uses isolated and purified exosomes, which contain the CD63 marker. The kit used consists of an exosome specific primary CD63 antibody developed by the manufacturer. Briefly, exosomes were immobilized on microtiter plates for overnight at 37°C using exosome binding buffer supplied the manufacturer (System Biosciences). Plates were washed and incubated at room temperature for 1h with exosome specific primary antibody (CD63), followed by a wash step and incubation with secondary antibody (1:5000) at RT for 1h with agitation. Plates were thereafter washed and incubated with Super-sensitive TMB ELISA substrate at RT for 45 min with agitation. The reaction was terminated using Stop Buffer solution. Absorbance was measured at 450nm. The number of exosomes/ml, (ExoELISA™ kit) was obtained using an exosomal CD63 standard curve that was generated using the calibrated exosome standard that was supplied.

## Nanoparticle Tracking Analysis

The size distribution of exosomes and their concentration were determined using the NS500 equipped with a 405nm laser and sCMOS camera (NanoSight NTA 3.0 Nanoparticle Tracking and Analysis Release, Version Build 0069). Plasma samples were diluted with PBS (1:100) prior to analysis in order to obtain particle distribution of 10 and 100 particles per image (optimal, 50 particles per image) before the analysis with NTA system. Samples were introduced into the sample chamber using the following script: PUMLOAD, REPEATSTART, PRIME, DELAY 10, CAPTURE 60 and REPEAT 5. Videos were recorded at a camera level of 10, camera shutter speed of 20ms and camera gain of 600, these settings were kept constant between samples. Each video was then analysed to give the mean particle size together with the concentration of particles. The size of the exosomes was represented as the mean particle size  $\pm$  SD.

## Transmission Electron Microscopy

Exosomes were applied to a continuous carbon grid and negatively stained with 2% uranyl acetate. The size and morphology of the particles were examined using a JEOL 1010 transmission electron microscope (JEOL, Peabody, MA, USA) at the Electron Microscopy Unit, University of KwaZulu-Natal.

## Statistical Analysis

Statistical analysis was performed using one-way analysis of variance (ANOVA) by SPSS software 14.0 followed by Duncan's multiple range test (DMRT). Results were expressed as mean $\pm$ S.D. from six rats in each group.

## RESULTS

### Blood Pressure, Body Weight and Heart Weight

A significant increase in blood pressure and heart rate ( $P<0.05$ ) in ISO-treated animals when compared to control group. At the end of the experiment, the body weight of group B (ISO-treated animals) was considerable decreased ( $P<0.05$ ) when compared to control group. The heart weight of ISO-treated animals was significantly increased ( $P<0.05$ ) when compared to the control group.

### Cardiac Markers

Rats treated with ISO showed significant ( $P<0.05$ ) increase in the activities of cardiac marker enzymes in serum compared to normal control rats. Figure 3A shows a considerable ( $P<0.05$ ) increase in the levels of serum troponin T (cTnT) in ISO-induced rats as compared to control rats. Figure 3B, 3C and 3D shows a significant increase in the activities of creatine kinase, creatine kinase-MB and lactate dehydrogenase (LDH) ( $P<0.05$ ) in ISO- treated rats when compared to control rats.

### Quantification of Exosomes Using ELISA



ELISA kit was used for the quantification of total exosomal protein. ISO-treated circulatory exosomal protein concentration was significantly increased (476mg protein/ml plasma) as compared with control (350mg protein/ml plasma). ELISA was used to quantify exosomal marker CD63, circulating exosomal protein (CD63) was significantly increased in ISO-treated rat's plasma ( $2.15 \times 10^8$  CD63/ml) as compared to normal control rat's plasma ( $1.10 \times 10^8$  CD63/ml).

## DISCUSSION

Biogenesis, characterization and functions of exosomes are a novel and exciting field of research that has triggered significant interest over the past decade. Exosomes (as well as other types of extracellular vesicles) play a vital role in regulating a broad range of physiological and pathological cellular processes. They may be utilized for therapeutic purposes and act as candidate biomarkers in various diseases. A vascular disease like MI affects a high proportion of the population. Based on the previously published studies we used an ISO model to induce MI in experimental rats in order to characterize and quantify exosomes secreted during MI. ISO is a potent synthetic catecholamine that causes severe stress in the myocardium, resulting in infarctlike necrosis of the heart muscle. ISO induces oxidative stress and results in alterations of cardiac function and ultrastructure in experimental rats. Experimental induction of MI by ISO in animals is a well-established model to study the protective role of various cardio protective agents. Studies have shown that animals develop MI when injected with high doses of catecholamines. Evidence of MI was initially indicated by the significant increase in the blood pressure and heart rate of the ISO-treated rats. The ISO administration caused the leakage of cardiac marker enzymes such as cardiac troponin T, CK, CK-MB and LDH of heart tissues to the circulation as evidence was indicated by the increase in the levels of these markers in circulation of ISO-treated rats. Increased levels of these

enzymes in the circulation may be due to myocardial necrosis, hypoxia and damage of cell membrane. In addition, histological analysis of ISO-treated heart tissue revealed damages in the myocardium.

Hypoxia is a potent stimulator for exosome release by cardiomyocytes. Studies have shown that exosomes secreted by primary cultured cardiomyocytes during hypoxia have a high content of tumor necrosis factor (TNF)- $\alpha$ , a proinflammatory cytokine whose cardiomyocyte expression is induced by hypoxia-inducible factor (HIF)-1 $\alpha$ . The type of stress to which myocardial tissue is exposed may determine the level of exosome secretion. During MI where the heart is exposed to ischemic stress signals such as inflammation, hypoxia and injury, cardiomyocytes are thought to elevate secretion of exosomes, but to the best of our knowledge, no study involve this model has provided evidence of the increased circulating exosome concentration secreted during MI. Our study provided evidence of increased levels of the exosome in the circulation during MI. Exosomes with a mean particle size distribution of (125.4nm) and concentration of ( $1.54 \times 10^8$  total exosomes/ml plasma) were obtained from control rat plasma. However, in the ISO-treated rats there was a significant increase in the quantity of exosomes with a mean particle distribution of (123nm) and concentration of ( $2.88 \times 10^8$  total exosomes/ml plasma) compared to control rat's plasma. Exosomal protein (CD63) was significantly increased in ISO- treated rats as compared to normal control rats (Figure 9), indicating increased levels of exosomes in the circulation during MI. Although exosomes were quantified from the circulation, which may have several sources, we assumed that cardiomyocytes secrete elevated amount of exosomes during MI.

Exosomes are the smallest extracellular vesicles, therefore purification methods of exosomes are limited and have not been validated. In our study, to confirm that the entities isolated from plasma were indeed exosomes, we used different techniques to

verify the particle size. NTA and TEM were used to characterize the exosomes. All particles obtained from TEM analysis were spherical in shape with a particle diameter of about 40-90nm. A slightly different particle size was obtained from NTA, particle diameter of about 90-130nm. The difference in the particle size distribution obtained from NTA may be because NTA uses light scattering and Brownian motion in order to obtain particle size distribution in a solution and large particles contribute more strongly to the light scattering than small particles thus are more visible. Hence, we concluded that the isolated extracellular vesicles were indeed exosomes.

Exosomes cannot only be viewed as potential biomarker candidate, are the key component of paracrine secretion in many cell types and are an important component in cell therapies. Exosomes may have either harmful or beneficial effect, as exosomes secreted from neonatal rat cardiac fibroblasts were found to be enriched with miR-21 which promote cardiac cardiomyocyte hypertrophy and exosomes secreted by Goto-Kakizaki (GK) rat cardiomyocytes were found to be enriched with miR-320 which inhibits myocardial endothelial cell promotion, migration and tube formation. In addition, exosomes secreted by platelets from patients with sepsis-induced endothelial cell apoptosis and cardiac dysfunction through NADPH, NOS and PDI. Several studies have focused only on the pathological role of exosomes as the entities that spread diseases throughout the body, however that may not be the only case. Therefore, it is important to know the actual quantity of exosomes secreted during heart-related diseases. Our study showed evidence that there is a significant increase in the secretion of exosomes during MI. Nevertheless, further studies are still required to verify this study and quantify cardiac specific exosomes secreted during MI.

This work is partly presented at 9th International Conference and Expo on Proteomics and Molecular Medicine, November 13-15, 2017, Paris, France.

## CONCLUSION

Our study has shown an increased exosome concentration secreted during MI. These findings indicate that exosomes are potential biomarkers for MI. However, exosomes were quantified in the circulation in this study, therefore the source is still unverified. Further studies are required to quantify cardiac specific exosomes secreted during MI.