Journal of Cancer Research and Immuno-Oncology

OPEN ACCESS Freely available online

Perspective

Lymphatic Vessel Biology Influencing Immune Surveillance across Diverse Human Malignancies

Emilia Laurent*

Department of Cancer Immunobiology, Western Continental University, Lyon, France

DESCRIPTION

Cancer progression is influenced by an extensive network of biological interactions involving malignant cells, stromal components, immune populations, vascular structures, and extracellular signaling molecules. Among these interconnected systems, the lymphatic vasculature has received increasing scientific attention because of its contribution to tissue fluid regulation, immune cell trafficking, antigen transport, and dissemination of malignant cells. For many years, investigations involving cancer primarily focused on blood vessels because of their importance in supplying oxygen and nutrients to expanding tumors. More recently, however, growing evidence has demonstrated that lymphatic vessels participate in numerous biological processes extending beyond fluid drainage. Their activities influence immune surveillance, communication between peripheral tissues and lymph nodes, cellular migration, and responses to modern immunotherapeutic interventions. As a result, lymphatic biology has become an important area of research within cancer immuno-oncology.

The lymphatic vascular system forms an extensive network distributed throughout nearly every organ of the human body. Unlike the circulatory system, which functions through continuous pumping by the heart, lymphatic vessels collect excess interstitial fluid, proteins, lipids, immune cells, and soluble molecules from surrounding tissues before returning them to the bloodstream. This process maintains tissue homeostasis while preventing excessive accumulation of extracellular fluid. Initial lymphatic capillaries possess specialized structural characteristics that permit efficient uptake of cells and soluble molecules from surrounding tissues. Larger collecting vessels subsequently transport lymph toward regional lymph nodes, where immune surveillance and antigen presentation occur before lymph reenters systemic circulation.

Under physiological conditions, lymphatic vessels contribute significantly to immune regulation. Antigen-presenting cells such as dendritic cells migrate through lymphatic channels after capturing foreign or abnormal proteins within peripheral tissues. These cells travel to draining lymph nodes, where they present processed antigens to naïve T lymphocytes and initiate adaptive immune responses. Simultaneously, soluble antigens transported through lymphatic fluid reach lymphoid organs for further immune evaluation. This coordinated transport system allows continuous monitoring of peripheral tissues while supporting rapid recognition of biological abnormalities.

Malignant transformation modifies normal lymphatic function through persistent communication between cancer cells and surrounding stromal populations. Tumor cells release cytokines, chemokines, extracellular vesicles, and growth factors capable of altering lymphatic endothelial cell behavior. Fibroblasts, macrophages, endothelial cells, and additional immune populations also contribute signaling molecules that reshape lymphatic architecture during disease progression. These molecular interactions influence vessel density, permeability, structural organization, and immune cell trafficking throughout the tumor microenvironment.

Lymphatic endothelial cells actively participate in immune regulation rather than serving solely as passive transport structures. These specialized endothelial cells express adhesion molecules, chemokines, and surface receptors that regulate migration of dendritic cells, lymphocytes, macrophages, and additional immune populations. Through controlled expression of these molecules, lymphatic vessels determine which immune cells enter lymphatic circulation and subsequently reach regional lymph nodes. Changes in endothelial signaling therefore influence initiation and maintenance of anti-tumor immune responses.

Dendritic cells maintain particularly close functional relationships with lymphatic vessels. Following recognition of tumor-associated antigens within peripheral tissues, dendritic cells migrate toward nearby lymphatic capillaries under the influence of chemotactic signals. After entering lymphatic circulation, they travel to draining lymph nodes where antigen presentation stimulates adaptive immunity. Efficiency of this migration process affects activation of cytotoxic T lymphocytes

Correspondence to: Emilia Laurent, Department of Cancer Immunobiology, Western Continental University, Lyon, France, E-mail: emilia.laurent@wcu-france.edu

Received: 27-Feb-2026, Manuscript No. JCRIO-26-43982; **Editor assigned:** 02-Mar-2026, PreQC No. JCRIO-26-43982(PQ); **Reviewed:** 16-Mar-2026, QC No. JCRIO-26-43982; **Revised:** 23-Mar-2026, Manuscript No. JCRIO-26-43982 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2684-1266.26.12.281

Citation: Laurent E (2026). Lymphatic Vessel Biology Influencing Immune Surveillance across Diverse Human Malignancies. J Cancer Res Immunooncol. 11:281.

Copyright: © 2026 Laurent E. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

responsible for recognizing malignant cells. Alterations in lymphatic communication therefore influence subsequent immune responses occurring throughout the body.

Cancer cells frequently utilize lymphatic vessels during early metastatic dissemination. Before entering systemic circulation, malignant cells may invade nearby lymphatic capillaries owing to their relatively permeable structural organization. Once transported through lymphatic channels, these cells reach regional lymph nodes, where some may survive and establish secondary lesions. Presence of malignant cells within lymph nodes often provides valuable clinical information regarding disease stage and potential systemic dissemination. Nevertheless, lymphatic involvement extends beyond metastatic transport because these vessels simultaneously participate in immune activation directed against tumor-associated antigens.

Liquid biopsy approaches may complement tissue-based analyses by detecting extracellular vesicles, cytokines, circulating nucleic acids, and soluble proteins associated with lymphatic activity. Serial sampling during treatment provides opportunities to monitor evolving biological changes without repeated invasive procedures. Integration of blood-based biomarkers with imaging findings and tissue pathology may improve assessment of disease progression and therapeutic effectiveness.

CONCLUSION

Future advances in cancer immuno-oncology will require comprehensive understanding of lymphatic vessel biology and its extensive interactions with immune populations, stromal cells, and malignant tissues. The lymphatic system participates in antigen transport, immune surveillance, cellular migration, tissue fluid regulation, and communication between peripheral tumors and regional lymphoid organs. These interconnected functions influence every stage of disease evolution, from early immune recognition to metastatic dissemination and therapeutic response. Continued investigation combining molecular biology, advanced imaging, computational analysis, and clinical research will strengthen current knowledge regarding lymphatic regulation within the tumor microenvironment. Such efforts may contribute to improved biomarker identification, refined patient stratification, and innovative therapeutic strategies that enhance anti-tumor immunity while improving long-term outcomes across a broad spectrum of human malignancies.