

Macrophage Polarization in Chagas Disease

Nagela Ghabdan Zanluqui¹, Priscilla Fanini Wowk¹ and Phileo Pinge-Filho^{2*}

¹Instituto Carlos Chagas, ICC-Fiocruz-PR, Rua Prof. Algacyr Munhoz Mäder 3775, Bloco C CIC, 81350-010, Curitiba, Paraná, Brazil

²Laboratório de Imunopatologia Experimental, Departamento de Ciências Patológicas Centro de Ciências Biológicas, Universidade Estadual de Londrina, 86051-970, Londrina, Paraná, Brazil

*Corresponding author: Pinge-Filho P, Departamento de Ciências Patológicas, Centro de Ciências Biológicas, Universidade Estadual de Londrina, 86051-970, Londrina, Paraná, Brazil, Tel: + 55 43 3371-4978; Fax: + 55 43 3371-4465; E-mail: pingefilho@uel.br

Received date: February 26, 2015, Accepted date: March 30, 2015, Published date: April 06, 2015

Copyright: © 2015 Zanluqui NG, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Macrophages are terminally differentiated cells of the mononuclear phagocytic system, which play an indispensable role in the maintenance of homeostasis and defense. Macrophages can be phenotypically polarized by the microenvironment to mount specific functional responses. Polarized macrophages can be broadly classified into two main groups: classically activated macrophages (M1), whose prototypical activating stimuli are IFN- γ and LPS, and alternatively activated macrophages (M2), further subdivided in M2a (after exposure to IL-4 or IL-13), M2b (immune complexes in combination with IL-1 β or LPS) and M2c (IL-10, TGF- β or glucocorticoids). M1 exhibit potent microbicidal properties and promote strong IL-12-mediated Th1 responses, while M2 macrophages support Th2-associated effector functions. Here we review the main functions of polarized macrophages in Chagas disease and discuss their potential value in evaluating disease severity.

Keywords: Macrophage polarization; *Trypanosoma cruzi*; Therapies

Introduction

Chagas disease is caused by the protozoan *Trypanosoma cruzi*, which affects approximately eight million individuals in Latin America, out of which 30-40% either already have or will develop cardiomyopathy, digestive mega syndromes, or both [1]. More recently, another major concern has been the emergence of Chagas disease in non-endemic areas such as North America, Europe and the Western Pacific Region, due to immigration of infected individuals [2,3]. The disease is characterized by two clinical phases: a short, acute phase defined by patent parasitemia, and a long, progressive, chronic phase.

The parasite load in the acute phase of *T. cruzi* infection influences the activation of host immune system and the development of Chagas disease pathology in the late chronic phase [4]. Although the exact mechanisms that mediate control of parasites in humans, have not been elucidated [5], it is believed that parasites rely greatly on the function of innate immune cells such as natural killer (NK) cells, neutrophils, and macrophages [6]. *T. cruzi* induces monocytes and macrophages to produce various endogenous mediators, including cytokines, nitric oxide (NO), and prostaglandin E₂ (PGE₂) [7-9].

Monocyte populations are heterogeneous and can polarize depending on the micro-environmental stimuli. Classically activated monocytes differentiate into M1 macrophages after stimulation by microbial agents such as lipopolysaccharides (LPS) and Th1 pro-inflammatory cytokines such as IFN- γ , whereas alternatively activated or M2 macrophages are induced by other stimuli, including transforming growth factor beta (TGF- β), interleukin-4 (IL-4), interleukin-10 (IL-10), and interleukin-13 [10,11].

The two polarized macrophage populations are also functionally different: M1 cells have inflammatory functions, produce high levels of pro-inflammatory cytokines, produce reactive nitrogen and oxygen

intermediates, and have bactericidal activity [12]. M2 cells have immunoregulatory functions, help in clearance of parasite, have increased phagocytic activity, and are involved in matrix remodeling, angiogenesis, wound healing [13], synovitis, and cartilage damage following hemarthrosis [14]. M2 macrophages are further characterized by the functional expression of their alternative activation markers. M2 macrophages have at least three subsets: M2a, induced by IL-4 or IL-13; M2b, induced by immune complexes and agonists of TLRs or IL-1 receptors; and M2c, induced by IL-10 and glucocorticoid hormones [15]. However, the distinct expression patterns of surface markers that clearly define macrophage subsets are still unclear, particularly in the case of human macrophages [16]. Recently, nomenclature and experimental guidelines were proposed to attain consensus between researchers regarding macrophage immunobiology [17].


The role of macrophage polarization in parasitic diseases is far from being well defined. Several pathogens exploit regulatory responses to facilitate immune escape and enhance their own survival in the host. For example, *Leishmania* binds and triggers Fc γ R signaling during entry into the host, resulting in the development of macrophages, which are permissive for its intracellular growth [18]. It is noteworthy that cell activation is critical for the induction of an effective immune response against pathogens or tumors, as inappropriate and sustained activation/polarization of macrophages can lead to tissue damage, immune dysfunction, and disease [18]. Given their capacity to suppress adaptive immune responses, it is important to understand how M1-M2 macrophages contribute to dysfunctional immune responses in infections such as Chagas disease. This review highlights the current understanding of the interplay between *T. cruzi* infection and macrophage polarization.

Macrophage polarization

M1-M2 nomenclature derives from the Th1 and Th2 cytokines: In 1986 Mosmann, Coffman et al. reported that murine T-lymphocytes

could be divided into Th1 and Th2 cells, based on their respective cytokine production profiles: IFN- γ and IL-4 [19]. The concept of macrophage polarization was first introduced in 1992 with the discovery that IL-4 inhibits the respiratory burst of macrophages while enhancing expression of MHC-II and mannose receptors (CD206) on

their cell surface [20]. Since then, two opposite and competing macrophages phenotypes were defined, often referred to as classically activated macrophages (M1 macrophages) and alternatively activated macrophages (M2 macrophages) [21,22].



In fact, macrophages can secrete either IL-12 or IL-10, cross regulatory cytokines crucial for the elicitation of IFN- γ production and development of Th1 cells or IL-4/IL-13 secretion and development of Th2 cells proliferation, respectively [22,23].

This classification was further extended by Mantovani and collaborators [24], M1 polarization included, the classical activation, obtained by stimulation with pathogen-derived LPS alone or in combination with IFN- γ , which improve microbicidal activity and pro-inflammatory mediator secretion, while M2 polarization, mainly associated with tissue repair, was further subdivided in M2a or alternatively activated macrophages; M2b, corresponding to type II activated macrophages; and M2c, which includes heterogeneous macrophage deactivation stimuli (Figure 1). Polarization states and

functional properties of macrophages largely depend on environmental conditions, such as hypoxia, cytokines, pathogen-derived TLR-ligands, and lipid mediators [21].

Some molecular factors underlying macrophage polarization have been identified [25,26]. These include members of the IRF/Stat families, Myc, NF- κ B hetero-and homo-dimers, KLF4, PPAR γ , as well as miRNA, and epigenetic modifications [25-27].

The observation that macrophage functional phenotypes can be manipulated has drawn attention towards macrophages as a potential therapeutic target for cancer therapy [28-30]. Thus, elucidation of the signaling pathways that regulate functional polarization of

macrophage in *T. cruzi* infection will aid in the designing of strategies for the modification of macrophage behavior.

Macrophage Polarization in *T. cruzi* infection

M1-M2 signatures: During the early stages of infection, *T. cruzi* induces an intense inflammatory response, which plays a crucial role

in the pathogenesis of the disease [31]. During this stage the macrophages induce a cascade of cytokines: initially they produce IL-12, which act on NK cells to induce IFN- γ production, which in turn increases the production of IL-12, TNF- α and NO in macrophages, thus contributing to the killing of the parasite [32,33].




Figure 2: Macrophage polarization in the acute phase of *Trypanosoma cruzi* infection. M1 polarization is associated with Th1 cytokine secretion and increased expression of iNOS, NADPH, MHC-class II, CD86, AP-1, and NF- κ B, in addition to that of inflammatory mediators such as TNF- α , IL-6, NO, ROS, IL-12, and IL-1 β , resulting in tissue damage and parasite killing (Left panel). M2 macrophages are related to tissue remodeling, encapsulation of parasite, Th2 cytokine secretion, which improves the expression of CD36, CD163, MR (mannose receptor), AMPK, Arg1, PPARs, and STAT6, in addition to improving IL-10 and TGF- β expression and polyamine secretion (Right panel). *T. cruzi* macrophage infection increases the expression of CD64, CD80, NADPH, COX2, iNOS, NF- κ B, STAT6, CD200, CD206, CD16, and CD32, as well as the secretion of TNF- α , IL-12, IL-1 β , IL-6, NO, and ROS, which promotes parasite elimination, and secretion of IL-10, TGF- β , and PGE₂, critical for immunoregulation and reduction of tissue damage resulting from excessive stimuli (Lower panel).

In fact, spleen or peritoneal macrophages harvested from mice in the acute phase of *T. cruzi* infection release large amounts of NO in the absence of any other stimuli and accumulate high levels of inducible nitric oxide synthase (iNOS) mRNA with secretion of TNF- α , IL-6 and IL-1 β in cultured macrophages [34,35]. This initial inflammatory response leads to M1 polarization.

The protective role of M1 macrophage has been exemplified in mice deficient for components of the IL-12 pathway [36]. Moreover, peritoneal macrophages from IL-12p40 gene knockout mice have a bias toward the M2 profile, spontaneously secreting large amounts of TGF- β and responding to rIFN- γ with weak NO production in *T. cruzi* infected peritoneal macrophages [37]. This host response is associated with the control of acute infection.

At the same time, macrophages and NK cells synthesize regulatory cytokines such as IL-10 and IL-4 to reduce the harmful effects associated with excessive stimulation of the immune system [38]. IL-4 in association with IL-10, modulates IFN- γ production and provides resistance against *T. cruzi* infection [39]. This new environmental change leads to M2 polarization. Thus, *T. cruzi* infection stimulates the activation of both M1 and M2 type macrophages during the early phases (Figure 2).

The persistence of inflammatory responses associated with tissue fibrosis and cell death is a hallmark of chronic Chagas disease. Macrophages present in chagasic hearts as well as those exposed *in vitro* to sera from *T. cruzi*-infected mice polarized toward a pro-inflammatory phenotype (CD64^{hi} CD80^{hi}) with extensive production of TNF- α /IFN- γ [40].

Interestingly, the multi-component DNA-prime/protein-boost vaccine (TcVac2) conferred protection against persistence of parasite and inflammation [41]. The vaccine induced a significant decline in the positive TNF- α /macrophage population in the heart and it was reflected by M2 polarization (CD200⁺, CD206⁺, CD16⁺, CD32⁺, IL-4, and IL-10 producing) upon *in vitro* incubation with sera from TcVac2-vaccinated infected mice [40].

PPAR α and PPAR γ ligands in M2 polarization

Peroxisome proliferator-activated receptors (PPARs) are ligand-dependent nuclear transcription factors. PPAR γ is a member of the nuclear hormone receptor family that has been implicated in inflammation and macrophage activation [42]. The importance of PPAR γ in regulating the M1/M2 phenotypic switch has been confirmed by Amine Bouhleh and collaborators (2007). These authors demonstrated that activation of PPAR γ potentiates the polarization of circulating monocytes into M2 macrophages [38]. 15-Deoxy- Δ 12, 14 prostaglandin J2 81 (15dPGJ2), has high affinity for PPAR γ and can exert its effects by either binding to PPAR γ or through interaction with intracellular targets such as the NF- κ B-signaling pathway or Erk MAP kinase cascade [38].

The role of PPAR γ ligands in *T. cruzi* infection is poorly understood. Hovsepian and collaborators (2011) reported the first evidence that treatment with 15dPGJ2 increases the number of intracellular parasites and inhibits the expression and activity of different inflammatory enzymes such as NOS2, matrix metalloproteinase as well as the mRNA expression of pro-inflammatory cytokines (TNF- α and IL-6) in neonatal mouse cardiomyocytes after *T. cruzi* infection [43].

Different investigations using PPAR α isoform ligands have shown a reduction in the symptoms of inflammation and disease in several models, including allergic airway disease, arthritis, and inflammatory bowel disease [38]. Penas et al. (2011) showed that treatment with either PPAR α or PPAR γ ligands drive M1-to-M2 polarization of macrophages in *T. cruzi*-infected mice [30].

Interestingly, PPAR agonists induce M1 macrophage polarization via cytosolic phospholipase A2 (cPLA2) and cyclooxygenase-2 (COX-2) inhibition, against *Leishmania mexicana* [44]. The potential use of PPAR agonists as modulators of overt inflammatory response during the course of Chagas disease can open a new avenue in the pharmacological approach to improve host health.

Manipulating macrophage metabolism is an attractive approach to controlling *T. cruzi* infection

When interacting with the host cell, *T. cruzi* stimulates the phosphorylation of several tyrosine residues, and this could be involved in the survival, multiplication, and pathogenicity of the parasite. Depending on the strain and developmental form of the parasite, distinct signaling pathways might be induced [42].

A recent study revealed for the first time the participation of mammalian target of mTOR in *T. cruzi* cell invasion [45]. Using an elegant study design, the authors demonstrated that the treatment of HeLa cells with the mTOR inhibitor, rapamycin, reduced lysosomal exocytosis and *T. cruzi* metacyclic trypomastigotes (MT) invasion. Downregulation of phosphatidylinositol 3-kinase and protein kinase C also impaired exocytosis and MT internalization. Thus, mammalian PI3/TOR kinase inhibitors can be a productive start point for anti-trypansomatid drug discovery. The only problem was that *T. cruzi* was relatively insensitive to rapamycin inhibitor compared with some other trypanosomatid pathogens, such as *T. brucei* [43].

However, we know that the AMP-activated protein kinase (AMPK) regulates energy homeostasis and metabolic stress, acting like a cellular energy sensor. AMPK is activated by high AMP and low ATP via a complex mechanism, which involves allosteric regulation, promotion of phosphorylation, and inhibition of dephosphorylation. When AMP/ATP ratio is high, AMPK is activated, it protects the cell by switching off the ATP-consuming pathways (e.g. fatty acid synthesis and sterol synthesis) and switching on alternative pathways for ATP generation (e.g. fatty acid oxidation) [46]. AMPK activation in macrophages is associated with M2 polarization, which suppresses pro-inflammatory responses and promotes anti-inflammatory functional phenotype [30]. Interestingly, when macrophages were stimulated with IL-10 and TGF- β , it resulted in the rapid activation of AMPK, whereas stimulation of macrophages with LPS-induced cytokines resulted in AMPK inactivation. In addition, inhibition of AMPK increased the mRNA levels of TNF- α , IL-6, and COX-2 [30]. AMPK has also shown to inactivate the mammalian target of rapamycin (mTOR) pathway via phosphorylation and activation of the mTOR inhibitor, tuberous sclerosis complex-2 TSC2 [47].

AMPK and mTOR are critical regulators of host cell metabolism making them logical targets for manipulation by invading pathogens such as *T. cruzi*. It would be interesting to determine if *T. cruzi* induces AMPK to generate energy and nutrients for its growth in the host cell. We must remember that inhibiting AMPK or inducing mTOR can provide essential conditions for *T. cruzi* replication [48].

Conclusion

Control of the *T. cruzi* infections is critically dependent on cytokine-mediated macrophage activation leading to intracellular killing of the parasite. M1 polarization is closely linked to the elimination of parasites, and M2 polarization could be effective in preventing the progression of oxidative and inflammatory pathology in Chagas disease. AMPK and mTOR are rational targets for manipulation by invading pathogens such as *T. cruzi*. Manipulation of host metabolism using PPAR agonists seems an attractive approach to controlling *T. cruzi* infection as targeting the host rather than the pathogen can considerably reduce the ability of pathogens to develop drug resistance.

References

1. Rassi A Jr, Rassi A, Marcondes de Rezende J (2012) American trypanosomiasis (Chagas disease). *Infect Dis Clin North Am* 26: 275-291.
2. Schmunis GA, Yadon ZE (2010) Chagas disease: a Latin American health problem becoming a world health problem. *Acta Trop* 115: 14-21.
3. Bonney KM (2014) Chagas disease in the 21st century: a public health success or an emerging threat? *Parasite* 21: 11.
4. Marinho CR, D'Império Lima MR, Grisotto MG, Alvarez JM (1999) Influence of acute-phase parasite load on pathology, parasitism, and activation of the immune system at the late chronic phase of Chagas' disease. *Infect Immun* 67: 308-318.
5. Álvarez JM, Fonseca R, Borges da Silva H, Marinho CR, Bortoluci KR, et al. (2014) Chagas disease: still many unsolved issues. *Mediators Inflamm* 2014: 912965.
6. Andrade DV, Gollob KJ, Dutra WO (2014) Acute chagas disease: new global challenges for an old neglected disease. *PLoS Negl Trop Dis* 8: e3010.
7. Gomes JA, Molica AM, Keesen TS, Morato MJ, de Araujo FF, et al. (2014) Inflammatory mediators from monocytes down-regulate cellular proliferation and enhance cytokines production in patients with polar clinical forms of Chagas disease. *Hum Immunol* 75: 20-8.
8. Borges MM, Kloetzel JK, Andrade HF Jr, Tadokoro CE, Pinge-Filho P, et al. (1998) Prostaglandin and nitric oxide regulate TNF-alpha production during *Trypanosoma cruzi* infection. *Immunol Lett* 63: 1-8.
9. Abel LC, Ferreira LR, Cunha Navarro I, Baron MA, Kalil J, et al. (2014) Induction of IL-12 production in human peripheral monocytes by *Trypanosoma cruzi* is mediated by glycosylphosphatidylinositol-anchored mucin-like glycoproteins and potentiated by IFN-gamma and CD40-CD40L interactions. *Mediators Inflamm* 2014: 345659.
10. Martinez FO, Helming L, Gordon S (2009) Alternative activation of macrophages: an immunologic functional perspective. *Annu Rev Immunol* 27: 451-483.
11. Mosser DM, Edwards JP (2008) Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* 8: 958-969.
12. Benoit M, Desnues B, Mege JL (2008) Macrophage polarization in bacterial infections. *J Immunol* 181: 3733-3739.
13. Martinez FO, Sica A, Mantovani A, Locati M (2008) Macrophage activation and polarization. *Front Biosci* 13: 453-461.
14. Nieuwenhuizen L, Schutgens RE, Coeleveld K, Mastbergen SC, Roosendaal G, et al. (2014) Hemarthrosis in hemophilic mice results in alterations in M1-M2 monocyte/macrophage polarization. *Thromb Res* 133: 390-395.
15. Mantovani A, Sica A, Sozzani S, Allavena P, Vecchi A, et al. (2004) The chemokine system in diverse forms of macrophage activation and polarization. *Trends Immunol* 25: 677-686.
16. Cassetta L, Cassol E, Poli G (2011) Macrophage polarization in health and disease. *ScientificWorldJournal* 11: 2391-2402.
17. Murray PJ, Allen JE, Biswas SK, Fisher EA, Gilroy DW, et al. (2014) Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity* 41: 14-20.
18. Miles SA, Conrad SM, Alves RG, Jeronimo SM, Mosser DM (2005) A role for IgG immune complexes during infection with the intracellular pathogen *Leishmania*. *J Exp Med* 201: 747-754.
19. Mosmann TR, Cherwinski H, Bond MW, Giedlin MA, Coffman RL (1986) Two types of murine helper T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J Immunol* 136: 2348-57.
20. Stein M, Keshav S, Harris N, Gordon S (1992) Interleukin 4 potently enhances murine macrophage mannose receptor activity: a marker of alternative immunologic macrophage activation. *J Exp Med* 176: 287-92.
21. Muraille E, Leo O, Moser M (2014) TH1/TH2 paradigm extended: macrophage polarization as an unappreciated pathogen-driven escape mechanism? *Front Immunol* 5: 603.
22. Gordon S (2003) Alternative activation of macrophages. *Nat Rev Immunol* 3: 23-35.
23. Gordon S, Martinez FO (2010) Alternative activation of macrophages: mechanism and functions. *Immunity* 32: 593-604.
24. Mantovani A, Sozzani S, Locati M, Allavena P, Sica A (2002) Macrophage polarization: tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. *Trends Immunol* 23: 549-555.
25. Sica A, Mantovani A (2012) Macrophage plasticity and polarization: in vivo veritas. *J Clin Invest* 122: 787-795.
26. Mantovani A, Biswas SK, Galdiero MR, Sica A, Locati M (2013) Macrophage plasticity and polarization in tissue repair and remodelling. *J Pathol* 229: 176-185.
27. Lawrence T, Natoli G (2011) Transcriptional regulation of macrophage polarization: enabling diversity with identity. *Nat Rev Immunol* 11: 750-761.
28. Watkins SK, Egilmez NK, Suttles J, Stout RD (2007) IL-12 rapidly alters the functional profile of tumor-associated and tumor-infiltrating macrophages in vitro and in vivo. *J Immunol* 178: 1357-1362.
29. Gratchev A, Kzhyshkowska J, Köthe K, Muller-Molinet I, Kannookadan S, et al. (2006) Mphi1 and Mphi2 can be re-polarized by Th2 or Th1 cytokines, respectively, and respond to exogenous danger signals. *Immunobiology* 211: 473-486.
30. Sag D, Carling D, Stout RD, Suttles J (2008) Adenosine 5'-monophosphate-activated protein kinase promotes macrophage polarization to an anti-inflammatory functional phenotype. *J Immunol* 181: 8633-8641.
31. Basso B (2013) Modulation of immune response in experimental Chagas disease. *World J Exp Med* 3: 1-10.
32. Brener Z, Gazzinelli RT (1997) Immunological control of *Trypanosoma cruzi* infection and pathogenesis of Chagas' disease. *Int Arch Allergy Immunol* 114: 103-110.
33. Camargo MM, Andrade AC, Almeida IC, Travassos LR, Gazzinelli RT (1997) Glycoconjugates isolated from *Trypanosoma cruzi* but not from *Leishmania* species membranes trigger nitric oxide synthesis as well as microbicidal activity in IFN-gamma-primed macrophages. *J Immunol* 159: 6131-9.
34. Hideko Tatakihara VL, Cecchini R, Borges CL, Malvezi AD, Graça-de Souza VK, et al. (2008) Effects of cyclooxygenase inhibitors on parasite burden, anemia and oxidative stress in murine *Trypanosoma cruzi* infection. *FEMS Immunol Med Microbiol* 52: 47-58.
35. Penas F, Mirkin GA, Vera M, Cevay Á, González CD, et al. (2015) Treatment in vitro with PPARα and PPARγ ligands drives M1-to-M2 polarization of macrophages from *T. cruzi*-infected mice. *Biochim Biophys Acta* 1852: 893-904.
36. Jouanguy E, Dörfinger R, Dupuis S, Pallier A, Altare F, et al. (1999) IL-12 and IFN-gamma in host defense against mycobacteria and salmonella in mice and men. *Curr Opin Immunol* 11: 346-351.
37. Bastos KR, Alvarez JM, Marinho CR, Rizzo LV, Lima MR (2002) Macrophages from IL-12p40-deficient mice have a bias toward the M2 activation profile. *J Leukoc Biol* 71: 271-278.
38. Sathler-Avelar R, Vitelli-Avelar DM, Teixeira-Carvalho A, Martins-Filho OA (2009) Innate immunity and regulatory T-cells in human Chagas disease: what must be understood? *Mem Inst Oswaldo Cruz* 104 Suppl 1: 246-251.
39. Abrahamssohn IA, da Silva AP, Coffman RL (2000) Effects of interleukin-4 deprivation and treatment on resistance to *Trypanosoma cruzi*. *Infect Immun* 68: 1975-1979.
40. Gupta S, Silva TS, Osizugbo JE, Tucker L, Spratt HM, et al. (2014) Serum-mediated activation of macrophages reflects TcVac2 vaccine efficacy against Chagas disease. *Infect Immun* 82: 1382-1389.
41. Gupta S, Garg NJ (2010) Prophylactic efficacy of TcVac2 against *Trypanosoma cruzi* in mice. *PLoS Negl Trop Dis* 4: e797.
42. Subramani PA, Reddy MC, Narala VR (2013) The need for physiologically relevant peroxisome proliferator-activated receptor-

- gamma (PPAR-gamma) ligands. Endocr Metab Immune Disord Drug Targets;13: 175-83.
43. Hovsepian E, Mirkin GA, Penas F, Manzano A, Bartrons R, et al. (2011) Modulation of inflammatory response and parasitism by 15-Deoxy- $\Delta(12,14)$ prostaglandin J(2) in *Trypanosoma cruzi*-infected cardiomyocytes. Int J Parasitol 41: 553-562.
44. Diaz-Gandarilla JA, Osorio-Trujillo C, Hernandez-Ramirez VI, Talamas-Rohana P (2013) PPAR activation induces M1 macrophage polarization via cPLA(2)-COX-2 inhibition, activating ROS production against *Leishmania mexicana*. Biomed Res Int 2013: 215283.
45. Martins RM, Alves RM, Macedo S, Yoshida N (2011) Starvation and rapamycin differentially regulate host cell lysosome exocytosis and invasion by *Trypanosoma cruzi* metacyclic forms. Cell Microbiol 13: 943-954.
46. Hardie DG, Carling D (1997) The AMP-activated protein kinase--fuel gauge of the mammalian cell? Eur J Biochem 246: 259-273.
47. Inoki K, Zhu T, Guan KL (2003) TSC2 mediates cellular energy response to control cell growth and survival. Cell 115: 577-590.
48. Brunton J, Steele S, Ziehr B, Moorman N, Kawula T (2013) Feeding uninvited guests: mTOR and AMPK set the table for intracellular pathogens. PLoS Pathog 9: e1003552.

This article was originally published in a special issue, entitled: "**Macrophage Polarization**", Edited by David J Vigerust, Vanderbilt University School of Medicine, USA