

Macrophage Polarization in Persistent Diseases: Current Therapeutic Perspectives

Elena Kovarik*

Department of Clinical Immunology, Central European Institute of Medical Sciences, Prague, Czech Republic

DESCRIPTION

Macrophages are highly adaptable leukocytes that perform diverse biological functions extending far beyond their traditional role in phagocytosis. These cells participate in tissue development, maintenance of physiological homeostasis, removal of apoptotic cells, regulation of vascular integrity, coordination of cellular communication, and restoration of damaged tissues following injury. Nearly every organ contains resident macrophage populations that develop unique biological characteristics according to their local microenvironment. Their remarkable ability to modify functional behavior in response to changing biological conditions, commonly described as polarization, enables rapid adaptation to infections, metabolic alterations, mechanical injury, and tissue remodeling. Careful regulation of macrophage polarization is essential because persistent imbalance may contribute to prolonged tissue dysfunction involving the lungs, kidneys, liver, gastrointestinal tract, cardiovascular system, skin, nervous system, and musculoskeletal tissues. Increasing understanding of macrophage biology has therefore encouraged development of therapeutic strategies directed toward selective regulation of cellular function while preserving normal physiological defense mechanisms.

Macrophages originate from two principal developmental sources. Many tissue-resident macrophages arise during embryonic development and maintain their populations through local proliferation without continuous replacement from circulating blood cells. Examples include microglia within the central nervous system, Kupffer cells in the liver, alveolar macrophages in the lungs, and specialized macrophages residing within the spleen and skin. Additional macrophages develop from circulating monocytes generated in the bone marrow. Following migration into tissues, these monocyte-derived cells differentiate according to signals provided by stromal cells, epithelial cells, endothelial cells, extracellular matrix proteins, cytokines, and local metabolic conditions. The coexistence of embryonic and monocyte-derived macrophages contributes substantially to functional diversity throughout the body.

Polarization describes the capacity of macrophages to adopt distinct functional characteristics according to environmental cues. Earlier biological models categorized macrophages into two broad groups commonly referred to as classically activated and alternatively activated phenotypes. Contemporary investigations demonstrate that macrophage behavior exists along a broad functional spectrum rather than within two rigid categories. Individual cells continuously modify cytokine production, receptor expression, metabolic activity, phagocytic capacity, and tissue repair functions according to changing biological conditions. This functional flexibility allows adaptation to highly diverse physiological requirements.

Removal of apoptotic cells represents one of the most important physiological responsibilities of macrophages. Billions of cells undergo programmed cell death each day within the human body. Efficient recognition and clearance of these dying cells prevent release of intracellular contents capable of stimulating prolonged tissue injury. Specialized receptors expressed on macrophage surfaces recognize phosphatidylserine and additional molecular signals exposed by apoptotic cells, initiating phagocytosis without provoking excessive cytokine production. Disturbances affecting this clearance process have been associated with chronic systemic disorders involving multiple organs.

Metabolic adaptation has emerged as a major determinant of macrophage function. Distinct polarization states are associated with characteristic patterns of glucose utilization, fatty acid oxidation, amino acid metabolism, and mitochondrial activity. Cells emphasizing antimicrobial activity frequently display increased glycolysis, whereas macrophages participating in tissue repair rely more extensively upon oxidative phosphorylation and fatty acid metabolism. Nutrient availability, oxygen concentration, and intracellular metabolic pathways therefore influence cellular behavior alongside traditional cytokine-mediated signaling.

Mitochondria perform functions extending beyond energy production. These organelles regulate calcium homeostasis, reactive oxygen species generation, apoptosis, and synthesis of metabolic intermediates affecting gene expression. Changes in

Correspondence to: Elena Kovarik, Department of Clinical Immunology, Nordic Institute of Medical Sciences, Aarhus, Denmark, E-mail : elena.kovarik@ceims-czech.edu

Received: 26-May-2026, Manuscript No. IDIT-26-43963; **Editor assigned:** 28-May-2026, PreQC No. IDIT-26-43963(PQ); **Reviewed:** 11-Jun-2026, QC No. IDIT-26-43963; **Revised:** 18-Jun-2026, Manuscript No. IDIT-26-43963 (R); **Published:** 25-Jun-2026, DOI: 10.35248/2593-8509.26.11.259

Citation: Kovarik E (2026). Macrophage Polarization in Persistent Diseases: Current Therapeutic Perspectives. Immunol Disord Immunother. 11:259.

Copyright: © 2026 Kovarik E. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

mitochondrial dynamics, including fusion, fission, and selective removal of damaged mitochondria through mitophagy, significantly influence macrophage polarization. Continued investigation seeks to determine whether pharmacological modulation of mitochondrial function may provide therapeutic benefit in chronic tissue disorders.

Extracellular vesicles released by macrophages facilitate communication with neighboring and distant cells. These membrane-bound particles transport proteins, messenger ribonucleic acid, micro ribonucleic acid, lipids, and signaling molecules capable of modifying recipient cell behavior. Macrophage-derived extracellular vesicles influence fibroblast activation, endothelial function, epithelial regeneration, and communication with lymphocyte populations. Their molecular composition reflects the functional state of the originating macrophages, making them attractive candidates for biomarker development.

Cell-based therapeutic approaches have also attracted scientific interest. Laboratory-expanded mesenchymal stromal cells release soluble mediators and extracellular vesicles capable of modifying macrophage polarization toward tissue-repair functions. Experimental investigations continue evaluating the safety, biological activity, and long-term effectiveness of these cellular therapies across several chronic disorders.

Nanotechnology offers additional opportunities for selective drug delivery because macrophages naturally internalize particulate material through phagocytosis. Biodegradable nanoparticles carrying therapeutic compounds may accumulate preferentially within macrophages, increasing intracellular drug concentrations while reducing systemic exposure. Surface modification techniques further improve targeting efficiency toward selected macrophage populations residing within affected tissues.

CONCLUSION

Continued investigation into macrophage polarization is expanding understanding of cellular communication, tissue regeneration, metabolic regulation, vascular biology, and molecular adaptation. These scientific developments are expected to support increasingly selective therapeutic strategies capable of restoring physiological macrophage function while preserving essential protective activities. Ongoing collaboration among laboratory researchers, clinicians, pharmaceutical scientists, and biomedical engineers will remain fundamental for translating these discoveries into practical clinical applications that improve long-term outcomes for individuals affected by persistent systemic disorders.