OPEN ACCESS Freely available online

Orthopedic & Muscular System: Current Research

Opinion Article

Medical Treatment Strategies for Osteoporosis in Elderly Patients

Sophie Grant *

Department of Geriatric Medicine and Orthopedic Research, Cambridge Clinical Institute, Cambridge, United Kingdom

DESCRIPTION

Osteoporosis in elderly patients represents a major public health concern due to its strong association with fragility fractures, disability, loss of independence, and increased mortality. In my opinion, the management of osteoporosis in older adults requires a carefully balanced medical approach that goes beyond simple fracture prevention and focuses on improving overall skeletal strength, reducing fall risk, and enhancing quality of life. The aging population presents unique clinical challenges, including multiple comorbidities, polypharmacy, altered drug metabolism, and reduced physiological reserves, all of which must be considered when designing an effective treatment strategy.

One of the central principles in the medical treatment of osteoporosis in elderly patients is early identification of individuals at high risk of fracture. Many patients remain undiagnosed until they experience a low-impact fracture, which often marks a turning point in their functional decline. Therefore, proactive screening using clinical risk assessment tools, bone density evaluation, and fracture history analysis is essential. In my view, timely diagnosis provides an opportunity to initiate therapy before irreversible skeletal damage occurs, thereby significantly reducing the burden of future fractures.

Pharmacological treatment remains the cornerstone of osteoporosis management in elderly individuals. Antiresorptive medications, particularly bisphosphonates, are widely used due to their proven effectiveness in reducing vertebral and hip fractures. These medications work by inhibiting bone resorption and stabilizing bone density over time. However, in older patients, careful consideration must be given to renal function, gastrointestinal tolerance, and adherence to dosing instructions. Alternative therapies, such as monoclonal antibody treatment that inhibits bone resorption through different mechanisms, offer additional options for patients who cannot tolerate first-line medications or who require more potent fracture protection.

Anabolic agents that stimulate bone formation represent an important advancement in the treatment of severe osteoporosis, especially in elderly patients with multiple fractures or very low bone density. These medications promote new bone formation and improve bone microarchitecture, thereby increasing skeletal strength more effectively than antiresorptive agents alone in certain high-risk populations. In my opinion, the strategic use of anabolic therapy followed by maintenance treatment with antiresorptive medication represents a rational and increasingly accepted approach in complex elderly cases.

Another critical aspect of medical treatment is the optimization of calcium and vitamin D levels. Deficiencies in these nutrients are common in elderly populations due to inadequate dietary intake, reduced sunlight exposure, and impaired absorption. Supplementation plays a supportive role in enhancing the effectiveness of pharmacological therapy and maintaining bone metabolism. However, supplementation alone is insufficient and must always be combined with disease-specific drug therapy and lifestyle modification.

Elderly patients often have multiple chronic diseases that complicate osteoporosis management. Conditions such as diabetes mellitus, chronic kidney disease, and cardiovascular disorders may influence drug selection and treatment duration. Polypharmacy increases the risk of drug interactions and adverse effects, making individualized treatment planning essential. In my view, a patient-centered approach that takes into account life expectancy, functional status, and personal treatment goals is more appropriate than a uniform treatment protocol.

Equally important is the issue of medication adherence. Many elderly patients discontinue therapy due to fear of side effects, misunderstanding of treatment benefits, or cognitive decline. Healthcare providers must therefore emphasize patient education, regular follow-up, and simplified dosing regimens to improve adherence. Support from caregivers and family members can also play a significant role in ensuring consistent treatment.

Although pharmacological therapy is central, it should not be viewed in isolation. The integration of fall prevention strategies, physical activity programs, and nutritional optimization significantly enhances treatment outcomes. Strength training and balance exercises can reduce the risk of falls, while proper nutrition supports overall musculoskeletal health.

Correspondence to: Sophie Grant. Department of Geriatric Medicine and Orthopedic Research, Cambridge Clinical Institute, Cambridge, United Kingdom, E-mail: sophie.grant@cambridgeclinical.ac.uk

Received: 02-Jun-2025, Manuscript No. OMCR-25-43038; **Editor assigned:** 05-Jun-2025, PreQC No. OMCR-25-43038 (PQ); **Reviewed:** 19-Jun-2025, QC No. OMCR-25-43038; **Revised:** 23-Jun-2025, Manuscript No. OMCR-25-43038 (R); **Published:** 30-Jun-2025, DOI: 10.35248/2161-0533.25.14.430.

Citation: Grant S (2025) Medical Treatment Strategies for Osteoporosis in Elderly Patients. J Orthop Muscular Syst. 14.430.

Copyright: © 2025 Grant S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.