Journal of Cancer Research and Immuno-Oncology

OPEN ACCESS Freely available online

Opinion Article

Microbiome-Derived Metabolites Modulating Cancer Progression and Therapeutic Responsivenes

Sofia Lindholm*

Department of Cancer Systems Biology, Scandinavian Institute of Biomedical Sciences, Helsinki, Finland

DESCRIPTION

Cancer develops through continuous interactions among malignant cells, stromal populations, vascular structures, extracellular matrix components, inflammatory cells, and numerous biochemical signals that collectively determine disease behavior. During the past decade, another influential biological factor has gained increasing scientific attention: the human microbiome. Trillions of microorganisms inhabit the gastrointestinal tract, oral cavity, respiratory system, skin, and additional body surfaces, forming highly organized microbial communities that participate in nutrient metabolism, maintenance of tissue integrity, regulation of inflammatory activity, and production of bioactive molecules. Rather than affecting health solely through direct contact with host tissues, these microorganisms continuously generate metabolites that circulate throughout the body and influence cellular physiology in distant organs. Within cancer research and immuno-oncology, microbiome-derived metabolites have become an important area of investigation because they contribute to cellular communication, metabolic regulation, inflammatory balance, and variability in therapeutic responsiveness among individuals with similar clinical diagnoses.

The human microbiome contains bacteria, fungi, viruses, archaea, and additional microorganisms existing in dynamic equilibrium with host tissues. Under physiological conditions, these microbial populations assist digestion, vitamin synthesis, nutrient absorption, and maintenance of epithelial barriers that separate host tissues from environmental microorganisms. The composition of microbial communities differs considerably among individuals owing to dietary habits, age, medication exposure, geography, lifestyle, environmental conditions, and host genetics. Despite this diversity, healthy microbial ecosystems generally maintain stable functional relationships with host physiology through continuous biochemical communication.

One of the principal mechanisms by which microorganisms influence human biology involves production of metabolites generated during nutrient fermentation and cellular metabolism. These metabolites include short-chain fatty acids, secondary bile acids, amino acid derivatives, indoles, polyamines, vitamins, and numerous additional low-molecular-weight compounds capable of entering systemic circulation. Once transported through the bloodstream, these molecules interact with epithelial cells, fibroblasts, endothelial cells, macrophages, dendritic cells, lymphocytes, and malignant cells. Consequently, microbial metabolism extends its biological influence well beyond the gastrointestinal tract.

Short-chain fatty acids have received considerable scientific attention because of their diverse physiological activities. Produced primarily through bacterial fermentation of dietary fiber, these metabolites influence epithelial barrier integrity, cellular differentiation, metabolic regulation, and inflammatory signaling. Within malignant disease, alterations in short-chain fatty acid concentrations modify communication among stromal populations and inflammatory cells while influencing cellular metabolism. Their biological activities demonstrate how dietary components, microbial metabolism, and host physiology become closely interconnected during cancer progression.

Bile acid metabolism represents another important pathway connecting microbial activity with cancer biology. Primary bile acids synthesized in the liver undergo chemical modification by intestinal microorganisms, generating numerous secondary bile acid derivatives possessing distinct biological properties. These compounds influence epithelial proliferation, lipid metabolism, cellular signaling, and inflammatory regulation. Changes in microbial composition alter the balance among different bile acid species, contributing to variability in cellular responses observed across patient populations. Ongoing investigations continue examining how bile acid metabolism influences malignant disease affecting both gastrointestinal and extraintestinal organs.

Dendritic cells located within intestinal tissues and peripheral organs also respond extensively to microbial metabolites. These specialized antigen-presenting cells process abnormal proteins and communicate with lymphocyte populations responsible for adaptive responses. Microbial compounds influence dendritic

Correspondence to: Sofia Lindholm, Department of Cancer Systems Biology, Scandinavian Institute of Biomedical Sciences, Helsinki, Finland, E-mail: sofia.lindholm@sibs-finland.edu

Received: 27-Feb-2026, Manuscript No. JCRIO-26-43985; **Editor assigned:** 02-Mar-2026, PreQC No. JCRIO-26-43985(PQ); **Reviewed:** 16-Mar-2026, QC No. JCRIO-26-43985; **Revised:** 23-Mar-2026, Manuscript No. JCRIO-26-43985 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2684-1266.26.12.284

Citation: Lindholm S (2026). Microbiome-Derived Metabolites Modulating Cancer Progression and Therapeutic Responsivenes. J Cancer Res Immunooncol. 11:284.

Copyright: © 2026 Lindholm S. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

cell maturation, migration, cytokine secretion, and antigen presentation efficiency. Through these mechanisms, microbial metabolism contributes to regulation of host defense while affecting biological interactions occurring within malignant tissues.

Lymphocyte populations likewise respond to metabolites generated by intestinal microorganisms. Short-chain fatty acids, indole derivatives, bile acid metabolites, and polyamines influence cellular differentiation, metabolic adaptation, proliferation, and cytokine production. Variability in microbial composition among individuals therefore contributes to differences in lymphocyte behavior observed during cancer progression and therapeutic intervention. Continued investigation seeks to clarify relationships between specific microbial metabolites and distinct lymphocyte populations across different malignancies.

Malignant cells themselves adapt to biochemical conditions created by microbiome-derived metabolites. Several microbial compounds influence intracellular signaling pathways regulating metabolism, proliferation, oxidative stress responses, and survival. These effects vary according to metabolite concentration, tissue location, and molecular characteristics of individual cancer types. Such observations indicate that microbial metabolism participates in continuous communication with malignant cells rather than functioning solely through indirect effects on surrounding stromal populations.

The intestinal epithelial barrier represents an important regulator of systemic exposure to microbial metabolites. Healthy epithelial integrity controls movement of microbial products into circulation while preventing excessive translocation of microorganisms. Alterations in epithelial permeability modify circulating metabolite concentrations and influence systemic inflammatory activity. Consequently, gastrointestinal physiology significantly affects communication between microbial communities and distant tissues involved in malignant disease.

CONCLUSION

Future progress in cancer research and immuno-oncology will depend upon continued investigation of biochemical communication linking microbial ecosystems with host tissues. Microbiome-derived metabolites influence epithelial physiology, stromal organization, vascular biology, inflammatory regulation, metabolism, and communication among diverse cellular populations participating in malignant disease. Their widespread biological activities illustrate the close relationship between microbial ecology and systemic human physiology. As analytical technologies continue advancing, comprehensive characterization of microbial metabolic networks will improve biomarker discovery, strengthen individualized patient assessment, and support development of innovative therapeutic strategies across a broad spectrum of human malignancies.