

Mini Review: Karyotypic Survey in Triatominae Subfamily (Hemiptera, Heteroptera)

Kaio Cesar Chaboli Alevi^{1*}, Joao Aristeu da Rosa², and Maria Tercilia Vilela de Azeredo Oliveira¹

¹Laboratory of Cell Biology, Department of Biology, Institute of Biosciences, Humanities and the Exact Sciences, Sao Paulo State University-Julio de Mesquita Filho (UNESP/BILCE), Sao Jose do Rio Preto, Sao Paulo, Brazil

²Laboratory of Parasitology, Department of Biological Sciences, College of Pharmaceutical Sciences, Sao Paulo State University-Julio de Mesquita Filho, (UNESP/FCFAR), Araraquara, Sao Paulo, Brazil

Abstract

The Triatominae subfamily consists of 145 species distributed in 18 genera and grouped in six tribes. Currently, there are 86 karyotypes described in the literature, distributed in 11 genera. There are five chromosomal complements described for these bloodsucking insects, out more, 22 (20A+XY), 23 (20A+X₁X₂Y), 24 (20A+X₁X₂X₃Y), 21 (18A+X₁X₂Y), 25 (22A+X₁X₂Y). Thus, we review all triatomine species with the number of chromosomes described in the literature. Through these data highlight the importance of further analysis cytogenetic with karyotype description in Triatominae subfamily, since it can help as an important tool cytotaxonomy and mainly allows the understanding of the evolution of this important group of insect vectors of Chagas disease.

Keywords: Karyotype; Holocentric chromosome; Diploid chromosome set

Mini review

Triatomines are insects that are taxonomically included in the Hemiptera order and the Heteroptera suborder within the Reduviidae family and in the Triatominae subfamily [1]. These organisms are of great medical importance, because all of the species that belong to the Triatominae subfamily are both bloodsucking and susceptible to infection from the protozoan *Trypanosoma cruzi* Chagas, 1909 (Kinetoplastida, Trypanosomatidae) and are therefore potential vectors of Chagas disease [2].

The Triatominae subfamily consists of 145 species [3-8] distributed in 18 genera and grouped in six tribes [9,10] (Table 1).

In cytogenetics, biological models of triatomine bugs are important because they have holocentric chromosomes, which have

diffuse kinetochores distributed along the chromosomes and not a single kinetochore located in the centromeric region. These insects also perform an unusual meiosis in which the segregation of sex chromosomes is post-reductional [11,12].

The description of the diploid chromosome set in Triatominae subfamily was initiated in 1909 with the description of the karyotype of *Triatoma sanguisuga* LeConte, 1856 [13]. In 1950, the karyotypic studies were resumed and new karyotypes were described in the literature [14].

Ueshima, in 1966, described the karyotype of twenty new species and proposed cytogenetic studies as a tool in the taxonomy of these vectors (cytotaxonomy). Moreover, the author analyzes the genetic viability during meiosis of experimental hybrids resulting from crosses between species of triatomines [12].

Currently, there are 86 karyotypes described in the literature, distributed in 11 genera (Table 1). Three tribes and seven genera show no cytogenetic studies maybe due to the limited geographic location which difficulty the granting of specimens of the species. The three principal genera known and studied maybe by large epidemiological importance are *Triatoma*, *Rhodnius* and *Panstrongylus*. However, even in these three genera, we emphasize the importance of new karyotypic studies, since, according to Table 1, we are aware of approximately 65%, 57% and 78%, the number of chromosomes in their respective genera.

Ueshima, in 1996, propose that the type number of chromosomes of Triatominae subfamily is 22 (20A+XY). However, he points out that there are five chromosomal complements described for these bloodsucking insects, out more, 22 (20A+XY), 23 (20A+X₁X₂Y),

Tribe	Genus	Number of Species	Karyotype Described
Alberproseniini	<i>Alberprosenia</i>	2	0
Bolboderini	<i>Belminus</i>	8	2
	<i>Bolbodera</i>	1	0
	<i>Microtriatoma</i>	2	0
	<i>Parabelminus</i>	2	0
Cavernicolini	<i>Cavernicola</i>	2	0
Linhcosteini	<i>Linhcosteus</i>	6	0
Rhodniini	<i>Psammolestes</i>	3	2
	<i>Rhodnius</i>	18	14
Triatomini	<i>Dipetalogaster</i>	1	1
	<i>Eratyrus</i>	2	2
	<i>Hermanlentia</i>	1	0
	<i>Meccus</i>	6	6
	<i>Mepraia</i>	3	3
	<i>Nesotriatoma</i>	3	2
	<i>Panstrongylus</i>	14	8
	<i>Paratriatoma</i>	1	1
	<i>Triatoma</i>	70	45
Total		145	86

Table 1: Revision of the Triatominae subfamily (species and karyotypes).

*Corresponding author: Kaio Cesar Chaboli Alevi, Instituto de Biociencias, Letras e Ciencias Exatas, IBILCE-UNESP, Rua Cristovao Colombo, 2265, Jardim Nazareth, 15054-000-Sao Jose do Rio Preto, SP-Brazil, Tel: (17) 32212380; E-mail: kaiochaboli@hotmail.com

Received May 25, 2013; Accepted June 14, 2013; Published June 17, 2013

Citation: Alevi KCC, da Rosa JA, de Azeredo Oliveira MTV (2013) Mini Review: Karyotypic Survey in Triatominae Subfamily (Hemiptera, Heteroptera). Entomol Ornithol Herpetol 2: 106. doi:[10.4172/2161-0983.1000106](https://doi.org/10.4172/2161-0983.1000106)

Copyright: © 2013 Alevi KCC, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Karyotype (2n)	Genus: species
21=18A+X ₁ X ₂ Y	<i>Panstrongylus: megistus</i> <i>Triatoma: nitida</i>
22=20A+XY	<i>Psammolestes: coreodes, tertius</i> <i>Rhodnius: brethesi, colombiensis, domesticus, ecuadoriensis, milesi, montenegrensis, nasustus, neglectus, neivai, pallescens, pictipes, prolixus, robustus, stali</i> <i>Dipetalogaster: maximus</i> <i>Paratriatoma: hirsuta</i> <i>Triatoma: arthurneivai, baratai, brasiliensis (b. brasiliensis, b. macromelanosoma), carcavalloi, circummaculata, costalimai, delpontei, garciabesi, guasayana, guazu, infestans (i. melanosoma), juazeirensis, jurbergi, klugi, lectularia, lenti, maculata, matogrossensis, melanica, patagonica, petrochiae, platensis, pseudomaculata, rubrovaria, sherlocki, sordida, vandae, williami</i>
23=20A+X ₁ X ₂ Y	<i>Belminus: herreri, corredori</i> <i>Eratyrus: cuspidatus, mucronatus</i> <i>Meccus: bassolsae, longipennis, mazzottii, pallidipennis, phyllosoma, picturatus</i> <i>Mepraia: gajardoi, parapatrica, spinolai</i> <i>Nesotriatoma: bruneri, flavida</i> <i>Panstrongylus: chinai, geniculatus, howardi, lignarius, rufotuberculatus, tupynambai</i> <i>Triatoma: barberi, dimidiata (d. capita, d. dimidiata, d. maculipennis), gerstaegeri, hegneri, mexicana, peninsularis, protracta, rubida, ryckmani, sanguisuga, sinaloensis, tibiamaculata</i>
24=20A+X ₁ X ₂ X ₃ Y	<i>Panstrongylus: lutzi</i> <i>Triatoma: eratyrusiformis, melanocephala, vitticeps</i>
25=22A+X ₁ X ₂ Y	<i>Triatoma: rubrofasciata</i>

Table 2: Chromosome number of 86 species of Triatomines with karyotypic analysis. A: autosomes.

24 (20A+X₁X₂X₃Y), 21 (18A+X₁X₂Y), 25 (22A+X₁X₂Y). Thus, we review the literature and grouped species according to the number of chromosomes (Table 2).

As can be observed in Table 2, some genera are poorly described derivatives as compared to the common ancestor, if we consider that the common ancestor of these bloodsucking insects presented 22 chromosomes (type number). On the other hand, if we analyze the tribe Triatomini, we observed that different chromosomal complements can be found, corroborating the hypothesis that the tribe is polyphyletic Triatomini.

Recently, through karyosystematic, Alevi and collaborated proposed a subcomplex of species were reorganized. The authors described the karyotype of the species *Triatoma lenti* Sherlock and Serafim, 1967 [15] and *Triatoma melanocephala* Neiva and Pinto, 1923 [16], and observed that *Triatoma melanocephala* Neiva and Pinto, 1923, as well as *Triatoma vitticeps* Stal, 1859 and *Triatoma tibiamaculata* Pinto, 1926 showed fragmentation of the sex chromosome X, different from other species also grouped in Brasiliensis subcomplex. Thus, as the initial proposal of the grouping of species in the subcomplex was performed using basically morphological data and geographical distribution [5], the number of chromosomes showed to be an important tool cytogenetic and, therefore, was proposed the exclusion of the three species with fragmented sex chromosome X of Brasiliensis subcomplex [16].

Thus, we review all triatomine species with the number of chromosomes described in the literature. Through these data highlight the importance of further analysis cytogenetic with karyotype description in Triatominae subfamily, since it can help as an important tool cytogenetic and mainly allows the understanding of the evolution of this important group of insect vectors of Chagas disease.

References

- Lent H, Wygodzinsky PW (1979) Revision of the Triatominae (Hemiptera, Reduviidae) and their significance as vector of Chagas's disease. Bull Am Mus Nat Hist 163: 123-520.
- Noireau F, Diasque P, Jansen AM (2009) *Trypanosoma cruzi*: adaptation to its vectors and its hosts. Vet Res 40: 26.
- Ayala JM (2009) Una nueva especie de *Panstrongylus* Berg de Venezuela. (Hemiptera: Reduviidae, Triatominae). Entomotropica. 24: 105-109.
- Jurberg J, Rocha DS, Galvão C (2009) *Zeledoni Rhodnius* sp. November

order of *Rhodnius paraensis* Sherlock, Guitton & Miles, 1977 (Hemiptera, Reduviidae, Triatominae). Biota Neotrop 9: 123-128.

- Schofield CJ, Galvão C (2009) Classification, evolution, and species groups within the Triatominae. Acta Trop. 110: 88-100.
- Frias-Lasserre D (2010) A new species and karyotype variation in the bordering distribution of *Mepraia spinolai* (Porter) and *Mepraia gajardoi* Frias et al (Hemiptera: Reduviidae: Triatominae) in Chile and its parapatric model of speciation. Neot Entomol 39: 572-583.
- Rosa JA, Rocha CS, Gardim S, Pinto MC, Mendonça, VJ, et al. (2012) Description of *Rhodnius montenegrensis* n. sp (Hemiptera: Reduviidae: Triatominae) from the state of Rondonia, Brazil. Zootaxa. 3478: 62-76.
- Gonçalves TCM, Teves-Neves SC, Santos-Mallet JR, Carballo-de-la-Fuente AL, Lopes CM (2013) *Triatoma jatai* sp. nov. in the state of Tocantins, Brazil (Hemiptera: Reduviidae: Triatominae). Mem Inst Oswaldo Cruz 108: 429-437.
- Galvão C, Carvalho RU, Rocha DS, Jurberg J (2003) A checklist of the current valid species of the subfamily Triatominae Jeannel, 1919 (Hemiptera, Reduviidae) and their geographical distribution, with nomenclatural and taxonomic notes. Zootaxa. 202: 1-36.
- Tartarotti E, Azeredo-Oliveira MT, Ceron CR (2006) Phylogenetic approach to the study of triatomines (Triatominae, Heteroptera). Braz J Biol. 66: 703-708.
- Barth R (1956). Estudos anatômicos e histológicos sobre a subfamília Triatominae (Heteroptera, Reduviidae). VI parte: Estudo comparativo sobre a espermatoctogênese das espécies mais importantes. Mem Inst Oswaldo Cruz 54: 599-616.
- Ueshima N (1966) Cytotaxonomy of the triatominae (Reduviidae: Hemiptera). Chromosoma 18: 97-122.
- Payne F (1909) Some new types of chromosome distribution and their relation to sex. Biol Bull 16: 119-166.
- Schreiber G, Pellegrino J (1950) Eteropicosi di autosomi come possibile meccanismo di speciazione. Sci Genet. 3: 215-226.
- Alevi KCC, Mendonça PP, Pereira NP, Rosa JA, Azeredo-Oliveira MTV (2012) Karyotype of *Triatoma melanocephala* Neiva and Pinto (1923). Does this species fit in the the Brasiliensis subcomplex? Infect Genet Evol. 12: 1652-1653.
- Alevi KCC, Mendonça PP, Pereira NP, Succi M, Rosa JÁ, et al. (2012) Karyotype of *Triatoma lenti* (Hemiptera: Triatominae), a potential Chagas vector. Gen Mol Res. 11: 4278-4284.