

Mitochondrial Genome Correction Approaches in Rare Energy Deficiency Syndromes

Nandini Rajasekar*

Department of Mitochondrial Biology, Institute of Cellular Medicine, Chennai, India

DESCRIPTION

Mitochondrial energy deficiency syndromes arise from dysfunction in genes responsible for oxidative phosphorylation and cellular Adenosine Triphosphate (ATP) production. These conditions may originate from mutations in mitochondrial Deoxyribonucleic Acid (DNA) or nuclear genes encoding mitochondrial proteins. Because mitochondria regulate essential energy-dependent processes across nearly all tissues, impairment often affects organs with high metabolic demand, including brain, skeletal muscle, heart, and liver. Clinical manifestations vary widely, ranging from developmental delay and muscle weakness to multi-organ metabolic instability.

Traditional management strategies have largely focused on symptomatic relief and metabolic supplementation, including cofactor replacement and dietary adjustments designed to reduce metabolic strain. However, such approaches do not directly address underlying genetic abnormalities. Recent advances in molecular therapeutics have shifted attention toward strategies capable of modifying mitochondrial genetic information or compensating for defective mitochondrial function at the molecular level. One area of investigation involves targeted mitochondrial DNA editing systems designed to selectively modify pathogenic variants within mitochondrial genomes. Unlike nuclear DNA, mitochondrial DNA lacks some of the repair pathways commonly used in the nucleus, making precise editing more challenging. To address this limitation, engineered nucleic acid-guided systems have been developed to selectively bind mutated mitochondrial sequences and induce controlled base changes or selective degradation of defective copies.

Another approach involves allotopic expression, where mitochondrial genes are relocated to the nuclear genome and expressed as nuclear-encoded proteins that are subsequently transported back into mitochondria. This strategy bypasses the limitations of mitochondrial DNA editing by providing functional protein products through nuclear transcription. Experimental studies have shown partial restoration of respiratory enzyme activity using nuclear-expressed mitochondrial gene constructs with mitochondrial targeting

sequences. Delivery of genetic material into mitochondria presents a significant technical challenge due to the double-membrane structure of the organelle. Researchers have explored peptide-based targeting sequences capable of guiding therapeutic molecules into mitochondria. Additionally, nanoparticle systems modified with mitochondrial penetration signals have been evaluated for their ability to cross cellular and organelle membranes. While efficiency remains variable, these systems represent important steps toward organelle-specific delivery.

Ribonucleic Acid (RNA)-based strategies have also been explored for mitochondrial disorders. Certain mitochondrial proteins are encoded in the nucleus and imported into mitochondria after translation. Enhancing expression of these nuclear genes through RNA activation systems may compensate for defects in mitochondrial-encoded components. In cell culture models, increased expression of supportive nuclear genes has improved mitochondrial respiration efficiency under stress conditions. Oxidative stress regulation is another critical factor in mitochondrial disorders. Dysfunctional mitochondria often produce excessive reactive oxygen species, leading to damage of cellular components. Therapeutic strategies aimed at enhancing endogenous antioxidant pathways have been investigated to reduce oxidative burden. Gene modulation approaches targeting antioxidant response elements have demonstrated protective effects in preclinical studies.

Tissue-specific variability is a major consideration in mitochondrial therapy. High-energy tissues such as neurons and cardiomyocytes are particularly sensitive to mitochondrial dysfunction. In neuronal models, mitochondrial gene correction has been associated with improved synaptic stability and reduced excitotoxic stress. In cardiac models, partial restoration of mitochondrial function has improved contraction efficiency and reduced arrhythmic tendencies. Pediatric mitochondrial disorders present additional complexity due to developmental energy requirements. Early-stage intervention may prevent irreversible developmental impairment, yet long-term safety of genetic modification in developing tissues remains under careful evaluation. Longitudinal animal studies are being conducted to

Correspondence to: Nandini Rajasekar, Department of Mitochondrial Biology, Institute of Cellular Medicine, Chennai, India, E-mail: n.raajasekar.icm@genmail.edu

Received: 01-Jun-2026, Manuscript No. JSGST-26-42893; **Editor assigned:** 03-Jun-2026, PreQC No. JSGST-26-42893 (PQ); **Reviewed:** 17-Jun-2026, QC No. JSGST-26-42893; **Revised:** 24-Jun-2026, Manuscript No. JSGST-26-42893 (R); **Published:** 01-Jul-2026, DOI: 10.35248/2157-7412.26.16.495

Citation: Rajasekar N (2026). Mitochondrial Genome Correction Approaches in Rare Energy Deficiency Syndromes. J Genet Syndr Gene Ther. 14:495.

Copyright: © 2026 Rajasekar N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

assess growth, neurological development, and metabolic stability following mitochondrial gene modulation.

Manufacturing of mitochondrial-targeted genetic therapeutics requires stringent quality control processes. Production includes synthesis of nucleic acid constructs, assembly of targeting peptides, and purification under controlled conditions. Functional validation involves assessment of mitochondrial localization efficiency and respiratory activity restoration in cellular assays. Clinical translation of mitochondrial gene modulation approaches remains in early stages. Most evidence currently derives from cell-based systems and animal models. Initial findings suggest that reducing mutant mitochondrial DNA load or enhancing compensatory pathways can improve cellular function, although outcomes vary depending on mutation type and heteroplasmy levels.

Continued advancement in mitochondrial genome engineering and compensatory gene expression systems indicates growing potential for addressing energy deficiency syndromes at their molecular origin. While significant technical barriers remain, ongoing refinement of targeting strategies, delivery platforms, and metabolic modeling tools supports further development of therapeutic options in this field of genetic medicine. At the chromatin dynamics level, sustained suppression of the antisense transcript responsible for paternal allele silencing was observed following treatment. Reduction of antisense RNA levels contributed to decreased recruitment of chromatin-modifying complexes associated with transcriptional repression. However, temporal analysis indicated gradual re-establishment of antisense transcription in certain neuronal populations over extended observation periods, suggesting that epigenetic

reactivation may require continuous or periodic reinforcement to maintain transcriptional accessibility.

Overall findings from this investigation demonstrate that long-acting RNA-guided epigenetic reactivation can partially restore gene expression, improve synaptic function, and enhance behavioral outcomes in models of Angelman syndrome. While full normalization of neuronal function was not achieved, the observed molecular, cellular, electrophysiological, and behavioral improvements highlight the therapeutic potential of epigenome engineering strategies for disorders caused by genomic imprinting defects. Further advancements in delivery persistence, chromatin stability control, and cell-type-specific targeting will be required to improve durability, consistency, and translational applicability of this approach in clinical contexts.

CONCLUSION

Overall findings from this experimental system indicate that allele-specific cytosine base editing can achieve partial correction of a pathogenic rhodopsin variant in a preclinical model of autosomal dominant retinal degeneration, resulting in measurable improvements in photoreceptor structure and function. While the intervention does not fully restore normal retinal physiology, the observed molecular and physiological changes demonstrate the potential utility of precise nucleotide-level modification strategies for conditions driven by single base substitutions. Further investigation will be required to enhance editing uniformity, extend duration of effect, and minimize immune-related limitations associated with vector-based delivery systems.