

Mitochondrial Genome Diversity and Evolutionary Dynamics in Pathogenic Fungi

Dorian Keller*

Department of Molecular Mycology, Northern Coast University, Bergen, Norway

DESCRIPTION

Fungal pathogens affect plants, animals, and humans across diverse ecosystems. While much attention has traditionally focused on nuclear genomes, mitochondrial genomes have become increasingly important in studies of fungal biology. These compact genetic systems perform essential cellular functions and contribute to energy production, metabolism, environmental adaptation, and evolutionary diversification. Advances in genome sequencing have enabled detailed examination of fungal mitochondrial DNA, revealing remarkable diversity in genome structure, organization, and evolutionary patterns among pathogenic species.

Mitochondria originated from ancient bacterial ancestors that established a symbiotic association with primitive eukaryotic cells. Over evolutionary time, many genes were transferred to the nucleus, leaving a reduced but functionally significant mitochondrial genome. In fungi, mitochondrial DNA generally exists as circular molecules, although linear forms have also been identified in several species. Genome sizes vary considerably, ranging from fewer than twenty thousand base pairs to well over two hundred thousand base pairs depending on species and lineage. One of the most striking characteristics of fungal mitochondrial genomes is their structural variability. Closely related pathogenic fungi may possess substantial differences in genome length, gene arrangement, intron content, and repetitive sequences. These variations arise through multiple mechanisms including insertions, deletions, recombination events, and mobile genetic element activity. Such diversity provides valuable information regarding fungal evolution and population history.

Energy production remains the primary function of fungal mitochondria. Genes encoded within mitochondrial DNA participate in oxidative phosphorylation, a process responsible for generating adenosine triphosphate, the major energy currency of cells. Pathogenic fungi require efficient energy production to support growth, reproduction, host colonization, and survival under stressful conditions. Mutations affecting mitochondrial genes can alter respiratory efficiency and influence pathogenic behavior. Comparative genomic

investigations have demonstrated that mitochondrial genes evolve at different rates among fungal groups. Some genes remain highly conserved across distant species due to strong functional constraints. Others accumulate mutations more readily, providing useful markers for phylogenetic analysis. Researchers frequently employ mitochondrial sequences to reconstruct evolutionary relationships and estimate divergence times among fungal pathogens.

Intron content contributes significantly to mitochondrial genome diversity. Introns are non-coding segments located within genes that are removed during Ribonucleic Acid (RNA) processing. Certain pathogenic fungi contain numerous mitochondrial introns, whereas others possess relatively few. Many introns carry open reading frames encoding homing endonucleases or reverse transcriptases capable of facilitating intron mobility. These elements can spread within populations and influence genome expansion. Repetitive Deoxyribonucleic Acid (DNA) sequences also play an important role in shaping mitochondrial architecture. Tandem repeats, dispersed repeats, and other repetitive elements create opportunities for recombination. Such events may generate genome rearrangements, duplications, or deletions that alter mitochondrial organization. In some pathogenic fungi, extensive repeat accumulation contributes substantially to differences in genome size between strains.

Pathogenic fungi encounter numerous environmental challenges during host infection. Temperature fluctuations, oxidative stress, nutrient limitations, and host immune responses place substantial demands on cellular metabolism. Mitochondrial function influences the ability of fungi to withstand such pressures. Genomic analyses have identified mitochondrial variants associated with stress tolerance, suggesting that mitochondrial evolution contributes to ecological adaptation. Drug resistance represents another area where mitochondrial genetics has attracted attention. Certain antifungal compounds target cellular processes linked directly or indirectly to mitochondrial activity. Mutations affecting mitochondrial genes may influence susceptibility to these treatments. Understanding

Correspondence to: Dorian Keller, Department of Molecular Mycology, Northern Coast University, Bergen, Norway, E-mail: d.keller.mycology@ncu-research.no

Received: 27-Feb-2026, Manuscript No. FGB-26-42476; **Editor assigned:** 02-Mar-2026, PreQC No. FGB-26-42476 (PQ); **Reviewed:** 16-Mar-2026, QC No. FGB-26-42476; **Revised:** 23-Mar-2026, Manuscript No. FGB-26-42476 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2165-8056.26.16.302

Citation: Keller D (2026). Mitochondrial Genome Diversity and Evolutionary Dynamics in Pathogenic Fungi. *Fung Genom Biol.* 16:302.

Copyright: © 2026 Keller D. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the genetic basis of such responses may support future efforts to improve disease management strategies.

Recent sequencing projects have uncovered evidence of horizontal transfer involving mitochondrial elements. Genetic material occasionally moves between unrelated fungal lineages through mechanisms that remain incompletely understood. These transfers introduce novel sequences into mitochondrial genomes and contribute to evolutionary complexity. Such findings challenge traditional assumptions regarding strictly vertical inheritance of mitochondrial DNA.

Mitochondrial-nuclear interactions represent a significant aspect of fungal biology. Although mitochondria possess their own genomes, many proteins required for mitochondrial function are encoded by nuclear genes. Cellular performance depends on coordinated communication between these genetic compartments. Evolutionary changes affecting one genome may influence selection pressures acting upon the other. Studies examining these interactions have revealed complex patterns of co-evolution in pathogenic fungi.

Advancements in long-read sequencing technologies have greatly improved mitochondrial genome assembly. Earlier approaches often struggled to resolve repetitive regions and structural complexity. Modern sequencing platforms generate longer reads

that facilitate accurate reconstruction of complete mitochondrial genomes. As a result, researchers can now investigate genomic features previously inaccessible through conventional methods. Another emerging area involves the examination of mitochondrial transcriptomes. Beyond DNA sequence analysis, researchers investigate patterns of mitochondrial gene expression under various conditions. Changes in transcript abundance provide insights into metabolic responses during infection, stress exposure, and developmental transitions. Integrating genomic and transcriptomic data offers a more comprehensive understanding of mitochondrial function.

CONCLUSION

The mitochondrial genome represents far more than a simple energy-producing component of the fungal cell. Its structure, sequence variation, inheritance patterns, and evolutionary history provide valuable information regarding biological processes operating across multiple scales. From cellular metabolism to species diversification, mitochondrial genetics influences numerous aspects of fungal life. Continued exploration of mitochondrial genome diversity will enhance understanding of pathogenic fungi and contribute to future developments in fungal genomics and biology.