

## Model Roles of the Hypothalamo-Neurohypophysial System in Neuroscience Study

Dan Hou<sup>#</sup>, Feng Jin<sup>#</sup>, Jiaoyan Li<sup>#</sup>, Jie Lian<sup>#</sup>, Mingyang Liu<sup>#</sup>, Xinna Liu<sup>#</sup>, Yanan Xu<sup>#</sup>, Chaoyu Zhang<sup>#</sup>, Chen Zhao<sup>#</sup>, Shuwei Jia<sup>#</sup>, Runsheng Jiao<sup>#</sup>, Xiaoyu Liu<sup>#</sup>, Xiaoran Wang<sup>#</sup>, Ying Zhang<sup>#</sup> and Yu-Feng Wang<sup>\*</sup>

Department of Physiology, Harbin Medical University, Harbin, China

<sup>#</sup>All authors contributed equally

### Abstract

Studies on the Hypothalamo-Neurohypophysial System (HNS) have played a pivotal role in establishing the fundamental principles of neuroscience. Examples include the identification of excitation-secretion coupling, coexistence of peptides-containing large dense core vesicles with classical neurotransmitters, neuronal-glia interactions, functions-associated morphological plasticity, and domain-specific regulation of the secretory activity, etc. Moreover, mechanisms underlying the secretion by the HNS and functions of neuropeptides oxytocin and vasopressin remain the central topic of the neuroendocrine society, novel findings of which keep substantiating the model role of this system. Nevertheless, challenges are still severe in this study field, such as extrahypothalamic connections of the HNS, scaffolding and guiding roles of glial fibrillary acidic protein in astrocyte plasticity, cellular and molecular processes determining different firing patterns, the position of the HNS in neuroendocrine-immune network, and the nose-hypothalamic route that serves as a key mediator of drug effects on brain activity. Nowadays, translational studies of the HNS also become the prime time of researchers, such as vasopressin involvement in brain injury in ischemic stroke, and therapeutic potentials of oxytocin in lactation failure and its associated health issues. In this paper, we review these challenges in modern neuroscience studies on the HNS.

**Keywords:** Neuropeptides; Neuroscience; Oxytocin; Supraoptic nucleus; Vasopressin

**Abbreviations:** GFAP: Glial Fibrillary Acidic Protein; HNS: Hypothalamo-Neurohypophysial System; HPA Axis: Hypothalamic-Pituitary-Adrenal Axis; MNCs: Magnocellular Neurosecretory Cells; OXT: Oxytocin; OXTR: OXT Receptor; PVN: Paraventricular Nucleus; SON: Supraoptic Nucleus; VP: Vasopressin

### Introduction

In the hypothalamus, there are two types of peptidergic magnocellular neurosecretory cells (MNCs): oxytocin (OXT)-producing neurons and vasopressin (VP)-producing neurons. OXT and VP (also called antidiuretic hormone, ADH) are the first group of neuropeptide hormones being sequenced and synthesized biochemically by Du Vigneaud, the Nobel prize laureate in 1955. This achievement largely determines the chemical nature of the mysterious pituitrin, pitressin and pitocin factors of the posterior pituitary that had been questioned since the identification of the vasopressor, uterus contraction, milk-letdown and antidiuretic effects of pituitary extracts. In studying on the neuroendocrine cells, particularly the MNCs of the Hypothalamo-Neurohypophysial System (HNS) that was initially established by Bargmann and Scharrer, mechanisms underlying the secretion and functions of VP and OXT have been the central topic of neuroendocrine society. From functional studies of circulating nonapeptides that extensively modulate the activities of almost all organ systems, to their brain actions that are represented by the autocrine regulation and prosocial effects, and then to the hidden secretion of peripherally produced OXT and VP that subtly adjust the activities of individual cells, studies on the HNS have been playing a pivotal role in establishing the fundamental principles of the neuroscience and the working models of neuropeptides. These achievements have been excellently reviewed and will not be further presented here [1,2].

Among several hypothalamic nuclei containing the MNCs, the supraoptic nucleus (SON) has many features that are suitable for characterizing neuron activities and their underlying mechanisms.

For example, these MNCs have relatively large cell bodies for their identification and electrophysiological manipulation, large amounts of secretory products for biochemical quantification, and relatively homogeneous cell populations that are easy to separate from other brain regions [3]. Thus, studies on the SON have greatly contributed to the concepts of the excitation-secretion coupling, neuronal-glia interactions, functions-associated morphological plasticity, and domain-specific regulation of the secretory activity, etc. In addition, the SON also has the following important features: 1) A clear separation of the somatodendritic zone from the axon terminals, which allows separate studies on the release profile of neuropeptides in the brain and in the circulation; 2) A distinguished spatial orientation of astrocyte processes, which is suitable for observing glial-neuronal interactions at microdomain and molecular levels; 3) Coexpression of classical neurotransmitters with neuropeptides, which provides a model to observe the interactions between different types of neurotransmitters in the autoregulation of MNC activity, and 4) Dependence of release patterns of the blood neuropeptides on their intranuclear release and on the activity of decomposing enzyme. After the initial findings, these features are also identified in many other types of neurons, particularly neuroendocrine cells [4,5]. Therefore, the SON remains a "model system" in our pursuing neuroscience research. In this review, we outline the challenges that we face in current studies on the HNS and our perspectives of future studies.

**\*Corresponding author:** Yu-Feng Wang, Department of Physiology, School of Basic Medical Sciences Harbin Medical University, 157 Baojian Rd, Nangang, Harbin, Heilongjiang 150081, China, Tel: +86-451-86674538; Fax: +86-451-86674538; E-mail: [yufengwang@ems.hrbmu.edu.cn](mailto:yufengwang@ems.hrbmu.edu.cn)

Received May 19, 2016; Accepted June 16, 2016; Published June 20, 2016

**Citation:** Hou D, Jin F, Li J, Lian J, Liu M, et al. (2016) Model Roles of the Hypothalamo-Neurohypophysial System in Neuroscience Study. *Biochem Pharmacol (Los Angel)* 5: 211. doi:10.4172/2167-0501.1000211

**Copyright:** © 2016 Hou D, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Histological features of the HNS

In the MNCs, OXT with its carrier protein neurophysin I and VP with neurophysin II are packaged in large, membrane-bound dense-core vesicles that are transported down the long axons to the nerve endings in the posterior pituitary [6]. In response to changes in extracellular environment, the MNCs are excited to generate action potentials that propagate down the axons and elicit exocytosis of some of these vesicles through excitation-secretion coupling. Released into extracellular space, these neuropeptides diffuse into the bloodstream through fenestrated capillaries and act on corresponding receptors including V1a, V1b and V2 subtypes of VP receptors and OXT receptor (OXTR) [7,8]. Moreover, the firing activity and release profiles of the MNCs are modulated by synaptic innervations, astrocytic-neuronal interactions, blood-borne factors and auto-regulation of MNC activities [9-12]. Broadly speaking, MNC-associated presynaptic terminals and astrocytes are also the integrative parts of the HNS that is mainly composed of the MNCs and their terminals.

The MNCs in the SON have 1-3 large dendrites that mainly project ventrally to receive most of the presynaptic terminals from afferent neurons and participate in information processing. Moreover, the dendrites and axonal collaterals of supraoptic neurons can release OXT and VP to exert neuromodulatory effects. The axonal efferent from the SON can project into various extrahypothalamic regions including the olfactory bulb (OB), the cortex, the amygdala, and the mammillary body complex in addition to a densely labeled pathway running from the SON to the posterior pituitary through the internal median eminence [13]. Although some cells around the SON could absorb radio-labeled leucine in the aforementioned study and partially account for the neural projection from the SON to these extrahypothalamic regions, it is clear that MNCs from the SON can modulate both hypothalamic and extrahypothalamic neuron activity. This proposal is further supported by the following evidence. A small proportion of the MNCs in the SON can send axon collaterals to the suprachiasmatic nucleus, a brain center regulating circadian rhythm, the antero-ventral third ventricular region near the organum vasculosum of the lamina terminalis where osmosensory neurons are identified [14], and to the medial preoptic area where gonadotropin-releasing hormone neurons are present [15]. Furthermore, dendrites of the MNCs can reach the ventricular system and thus exert neurohormonal effects on extensive CNS areas through the CSF [16,17]. In addition, there are close interactions between the SON and the paraventricular nucleus (PVN), which could be mediated by both neural innervations and paracrine modulations [18-20]. These connections form a structural basis of the simultaneous activation of a large pool of HNS neurons in the bilateral hypothalamus, typically observed in OXT neurons during suckling [21]. Importantly, interactions of some brain functions with OXT neurons in the SON are different from those in the PVN, such as that the activity of the gonadotropin-releasing hormone neurons in the medial preoptic area has close interactions with OXT neurons in the SON but not in the PVN [15]. These facts indicate that the neuromodulatory effects of MNCs in the SON on brain activity and behaviors are largely underestimated over the years, largely because parvocellular division of the PVN has been considering as the major source of brain OXT and VP after its early findings [22]. Moreover, the SON likely exerts different effects from the PVN does on some brain functions, such as the regulation of estrous cycle [23] and of olfaction [24]. Nevertheless, further studies are required to depict the exact trajectories of these neural projections from the MNCs. Figure 1 is a schematic illustration of the histology of the SON and the targets of its neural projections.

## Morphological and functional plasticity


An important feature of the HNS is its morphological plasticity. As revealed by electron microscopic observations in rats, dehydration increases the size of the MNCs and significantly reduces glial coverage of these cells in the SON [25]; lactation increases the size of OXT neurons, the numbers of shared synapses, and the apposition between adjacent OXT neurons [26]. This plastic change was originally thought to be specific to OXT neurons [27] that are mainly located in the dorsal part of the SON [28]. However, a more detailed morphologic analysis revealed that VP-ergic dendrites become more extensive in the SON, which is likely to fill in the open space left by the reduction of dendritic branches and total dendritic length of OXT neurons during lactation [29]. This morphological plasticity can partially explain the synchronized activation of OXT neurons during suckling as well as the changes in the excitability of MNCs in response to osmotic challenges that have been previously reviewed [10,30] and thus will not be further discussed here.

It is worth noting that the chronic morphological plasticity can be compensated by some other mechanisms since disabling this plasticity in the SON did not significantly influence OXT neuronal activity as shown in studying the burst discharges of OXT neurons during suckling stimulation [31]. In contrast, acutely occurred morphological plasticity in the SON is likely irreplaceable for the milk-ejection reflex and for the osmotic reaction and thus constitutes the structural basis of acute functional plasticity of the HNS [32-34].

Acute morphological plasticity of the HNS under different functional states is closely related to acute astrocytic plasticity [10] that is largely dependent upon plastic changes in GFAP molecules [35]. In the studies on supraoptic astrocytic morphological plasticity in rat models of lactation and dehydration [36-38], GFAP plasticity has been used as an indicator of gliosis or the retraction of astrocyte processes. The GFAP plasticity is closely related to astrocyte functions, such as buffering changes in extracellular  $K^+$  and glutamate levels [39], regulating extracellular volume transduction [40], adjusting physical barriers between adjacent neurons and many others [33,34,39].

The contribution of GFAP to astrocytic plasticity has been well shown in the SON during suckling [37] and in response to hypoosmotic challenges [41,42]. However, the scaffolding and guiding roles of GFAP in acute astrocytic plasticity are not well known. For example, in glutamate metabolism, GFAP has molecular associations with glutamine synthetase and serine racemase and thus, could relocate these molecules during its polymerization and depolymerization [37], thereby modulating glutamate-glutamine cycle in a microdomain-specific manner as previously discussed [32,43]. Similarly, GFAP has also molecular association with vesicular GABA transporters and thus can modulate GABA metabolism and actions [41]. These evidences suggest that GFAP can be a scaffold to provide a platform for the interactions between different molecules while serving as a transporter to guide their spatial localization of functions.

The contribution of the scaffolding and guiding roles of GFAP to astrocytic plasticity is further supported by the following studies. Aquaporin 4 (AQP4), another binding partner of GFAP, shows changes in its expression levels as the retraction or expansion of astrocyte processes around OXT neurons at different stages of suckling stimulation [37]. The membrane installation of AQP4 following the occurrence of the milk-ejection burst could allow more water to get into the cells and facilitate the expansion of astrocyte processes. This could be accompanied by the activation of  $Na^+$ ,  $K^+$ ,  $2 Cl^-$  cotransporter-1




[44] and AQP4-coupled inwardly rectifying  $K^+$  channel 4.1 [39], as a result of burst-evoked increase in extracellular  $K^+$  levels [45,46]. The activation of these ion transporters decreases extracellular  $K^+$  levels and could also lead to the post-burst inhibition [47] in OXT neurons. Conversely, the reduction of AQP4 during glial retraction exerts an opposite effect: reducing absorption of extracellular  $K^+$  and glutamate, leading to gradual increases in the inter-burst firing rate in OXT neurons during suckling [47]. Since GFAP plasticity occurs extensively in a variety of physiological and pathological conditions, such as the site-dependent GFAP plasticity in ischemic stroke [39,48], further exploration of the scaffolding and guiding roles of GFAP in astrocytic plasticity will profoundly impact our understandings of the glial-neuronal interactions and in turn manifest new targets for treatment of astrocyte-associated diseases. Figure 2 is a diagrammatic drawing of the scaffolding and guiding roles of GFAP in acute astrocytic plasticity and their influences on neuronal activity.

### Cellular and molecular processes leading to different firing patterns

In response to different physiological demands, VP neurons and OXT neurons exhibit different firing patterns and secretory profiles of the neuropeptides. For example, hyperosmotic and hypovolemic stimuli can increase the release of VP that in turn causes water reabsorption in the kidney and vasoconstriction to reduce blood osmolality and elevate blood pressure. In this process, the firing rate of VP neurons increases in a phasic pattern that occurs periodically based on a plateau potential [49]. Hyperosmotic stimulation also increases the firing activity of OXT neurons and OXT release that appear in a tonic pattern but not

in phasic or burst pattern [50]. During lactation [51] and parturition [52], the firing activity of OXT neurons appears in an intermittently bursting pattern on the background of a continuous or irregular firing activity. This milk-ejection burst is highly synchronized among OXT neurons in the SON and the PVN [21,53] and is followed by a bolus release of OXT [54]. In general, the phasic and burst firing patterns are more efficient for hormone release than the tonic firing pattern does with the same number of spikes [49]. Compared to the phasic firing activity of VP neurons, the burst firing in OXT neurons is much shorter and more intense [55]. This high frequency burst discharges of OXT neurons can account for the bolus release of OXT during suckling stimulation while the randomly occurred phasic firing in VP neurons can explain the intensively increased VP release in response to hyperosmotic challenges. This association between firing patterns and neurosecretion of the MNCs provides a suitable model for studying the mechanisms underlying pattern generators of neuronal activities and their association with different functions.

In investigating mechanisms underlying the differences in firing patterns between OXT and VP neurons, many questions have been explored, such as the afferent pathways transmitting different stimuli and the chemical nature of synaptic connections [56], electrophysiological features of VP and OXT neurons *in vivo* [51] and *in vitro* [57], genes associated with OXT- or VP-neurons [58], signaling molecules [59] and ion channels [57,60,61] that trigger special firing patterns. Among them, burst firing activity of OXT neurons specifically depends on suckling stimulation [62], intranuclear release of OXT [63], and a neuromodulation of the “burst synchronization center” in the mammillary body complex [64]. Relative to the regulation



of OXT neuronal activity, the regulation of VP neuronal activity is “non-specific.” The type of stimulation and extracellular factors relating to the phasic firing are relatively loose although the activity of VP neurons is closely regulated by the anteroventral region of the third ventricle [65]. At the cellular level, OXT but not VP neurons possess a rebound depolarization that underlies a brief burst of action potentials while exhibiting a broader spike width and a larger Ca<sup>2+</sup>-dependent afterhyperpolarization [66]. However, the identified *in vivo* electrophysiological features largely reflect the extracellular environment. For example, hyperosmotic stimulation triggers phasic firing while OXT is the permissive factor of burst discharge, which does not show the link between the signaling features of these MNCs and firing patterns. *In vitro* studies on chemically-identified MNCs can touch the cellular and molecular processes accurately; however, previously identified electrophysiological differences between the two types of MNCs remain controversial. That is, in previous study, the presence of a sustained outward rectifying potassium current and/or an inward rectifying hyperpolarization-activated current are presumably present in OXT neurons but absent in VP neurons [67]. However, the phenotypes classified according to the electrophysiological protocol in brain slices do not match their molecular counterparts because VP and intermediate neurons also exhibit both outward and inward rectifying currents [68]. As a whole, differences in the cellular and molecular processes leading to the different firing patterns are largely unclear.

What we know better are the association between phasic burst in VP neurons and a series of cellular and molecular processes as excellently reviewed [57,69]. In contrast, only a few of publications

using *in vitro* approaches adopted strict criteria that reflect the features of burst firing of OXT neurons. By simulating the neurochemical environment existing during lactation in the SON, several burst models were created in the last two decades. One was produced in a low calcium medium in brain slices from lactating rats [55] or male rats [70] by bath-application of phenylephrine, the  $\alpha$ 1 adrenoceptor agonist. Another one was established in brain slices from lactating rats [71] by extracellularly applying OXT, the permissive factor of burst occurrence *in vivo* [63,72]. In the third one, a spontaneous burst of OXT neurons was identified in organotypic cultures of the hypothalamic explants from neonate rats [73]. By using these models, the differences in the firing patterns between the two types of MNCs are outlined. That is, bursts in OXT neurons are shorter in duration, higher in peak firing rates, wider in the half width of action potentials and not superimposed on a plateau potential. Moreover, burst occurrence depends on the activation of OXTRs and glutamatergic synaptic transmission [73] but not on GABAergic synaptic transmission in brain slices [71]. Interestingly, OXT suppresses tonic glutamatergic and GABAergic synaptic transmission [11] but evokes short clustered glutamate and GABA release that is highly correlated with the burst features in OXT neurons [71]. One contradictory phenomenon is the lack of reports of burst evoked directly by exogenous glutamate so far while exogenous GABA in the SON could evoke burst discharge in lactating rats [74]. Thus, it is likely that the burst occurrence in OXT neurons needs an excitatory environment, based on which the mobilization of OXTR-specific signaling cascade can trigger the burst discharges as reviewed previously [75]. In addition, clustered GABAergic inputs on OXT

neurons are also likely involved in the burst generation [76], which is assumed to trigger a hyperpolarization-evoked rebound depolarization [66]. However, causal relationship of these clustered EPSCs and IPSCs with the burst onset remains to be elucidated.


At molecular levels, relative to the classical  $\alpha$ -subunit signaling in OXT actions,  $G\alpha_{q/11}$ -associated  $G\beta\gamma$  subunit signaling plays a dominant role in the burst firing of OXT neurons at both pre- and postsynaptic sites [71]. In the downstream of the  $G\beta\gamma$  subunits, phosphorylated extracellular signal-regulated protein kinase 1/2 and actin reorganization at the membrane subcortical areas [77] are essential to the burst generation. However, the dependence of phosphorylated extracellular signal-regulated protein kinase 1/2 [78] and actin reorganization [79] are also essential for the phasic firing in VP neurons. The common involvement of these signaling molecules in different types of the MNCs does not mean that the signaling processes are the same for both the phasic and burst firing patterns. It is possible that the two types of MNCs express these molecules in different spatial and temporal orders in response to their adequate stimuli. That is, the actin cytoskeletal reorganization occurs before the burst in OXT neurons following a cascade of cytosolic signaling molecules during suckling [75] while osmotic challenges in VP neurons elicit actin reorganization following a series of membrane events and triggering the phasic firing [79]. Another line of evidence explaining the differences in the firing patterns is from the studying on cell-specific expressions of calcium binding proteins. Immunohistological studies revealed that OXT neurons have more calbindin-D28k and calretinin than VP neurons [80], which allows  $Ca^{2+}$  oscillation in OXT neurons to be buffered quicker than that in VP neurons. This feature determines that OXT neurons discharge action potentials in either continuous or burst pattern while VP neurons adopt the phasic firing pattern [81]. Nevertheless, it remains to clarify how these different signaling processes are linked to changes in the extracellular milieu and ion channel activity that underlies the different firing patterns.

As more ion channels and their functions are identified in the MNCs, such as ATP-activated transient receptor potential vanilloid channels [82], the approaches of their contributions to the burst firing are also questioned. Further study should clarify if the burst onset is associated with a group of ion channels or a single channel, based on appropriate burst-firing models. Overall, burst model-based experiments are required to identify the expression, distribution, and functions of individual channels as well as their associations with burst-evoking signaling molecules, which should fully consider factors at levels of system, local neural circuit, MNCs, glial-neuronal interaction, receptors and intracellular signaling as well as ion channels of an organism (Figure 3).

### The HNS-immune network

The first and a major hypothalamic peptidergic system involving in neuroimmunologic regulation is considered to be the hypothalamic-pituitary-adrenal (HPA) axis [83]. The HPA axis refers to a complex set of direct influences and feedback interactions between the hypothalamus, the pituitary gland, and the adrenal glands. Under the regulation and coordination of hypothalamic corticotrophin-releasing hormone, the HPA axis can prime the immune system and potentiate acute defensive responses while suppressing immunologic activities by antagonizing adrenergic pro-inflammatory actions [84].

In parallel with the role of the HPA axis in immune regulation, the HNS is also implicated in neuroendocrine regulation of immunologic



activities. On the one hand, the HNS is the target of immunologic activities. For instance, thymectomy causes general atrophy of the SON, PVN and the median eminence [85]; immune cytokines [86] and pathogens [87] can change the activity of neuroendocrine neurons and neuropeptide secretion. On the other hand, the HNS can profoundly modulate the development, differentiation and functions of the immune system. For example, removal of the intermediate and posterior lobes of the pituitary reduces hormone- and cell-mediated immune responses to pathological challenges in rats as recently reviewed [88]. In rat tissue cultures, VP inhibits while OXT promotes the growth of thymic glands [89]. OXT reduces stress-induced psychological illness by strengthening social bonding while VP and corticotrophin-releasing hormone enhance autonomic reactivity and increase susceptibility to adverse mental and physical impacts through its facilitating arousal [90]. In addition, OXT can extensively modulate the activity of immunologic system, such as increasing the production of hematopoietic progenitor cells, reducing inflammatory cytokines, inhibiting the invasion of neutrophils [88]. Consistently, blocking OXTRs inhibits T-cell differentiation in the thymus [91] and activates the excretion of the inflammatory cytokines, interleukin-6 and Chemokine (C-C motif) ligand 5 [92,93] while neutralization of VP specifically and reversibly blocks interferon- $\gamma$  production in mouse splenic lymphocytes [94]. Moreover, intracerebroventricular administration of VP suppresses the proliferative response of splenic T-cells and natural killer cell cytotoxicity. Consistent with the actions of these two neuropeptides, OXTRs are found in monocytes and macrophages [95], thymic T-cells [96], and mesenchymal stromal cells of adult bone marrow [97] while VP receptors are found on splenic membrane, splenic lymphocytes [96] and CD4+ and CD8+ thymic cells [98]. Clearly, there are bidirectional communications between the HNS and the immune system, in which the HNS could function as a higher neuroendocrine regulatory center of the immune system and modulate immunologic responses differentially and coordinately through OXT and VP, respectively. This proposal expands the view that the OXT-secreting system is a major part of the neuroendocrine center regulating immunologic activity [88].


To claim that the HNS is a higher neuroendocrine regulatory center of the immune system, it is essential to clarify the relationship between

the HNS and the HPA axis. On the one hand, glucocorticoids secreted in response to stress activation of the HPA axis can exert rapid negative feedback influences onto the HNS and thus, reduce the secretion of OXT and VP [99,100]. On the other hand, the activity of HPA axis is heavily modulated by VP and OXT. For example, VP can promote the secretion of adrenocorticotropin hormone in the pituitary, particularly in males [101] while OXT inhibits the activity of the HPA axis [102]. Thus, the HPA axis and the HNS form a complex feedback loop while functioning collaboratively in responses to immunologic challenges. Consistently, the central inhibition of VP on the proliferative response of splenic T-cells could be due to VP activation of HPA axis in the brain that directly inhibits splenic responses to pathologic challenges [103]. Nevertheless, it remains to explore the subtle interactions between these immune-associated neuropeptide systems.

Together with the proposal by Pittman [104], we propose that in the neuroendocrine-immune network, the HNS can integrate immunologic, neural signal, and hormonal messages and regulate immunologic activities directly as well as indirectly through the HPA axis (Figure 4). By this way, the HNS can tailor the immunologic responses to meet body's defense requirement accurately. Further study should differentiate the contributions of the VP-secreting system from that of the OXT-secreting system to the hypothalamus-immune network.

### Nose-HNS route mediating nasal drug effects on brain activities

As the accumulation of knowledge of the NHS, particularly their involvement in diseases such as acute hyponatremia [43] and ischemic stroke [105], exploring non-invasive approaches modulating HNS activity has become a strong challenge to neuroscience researchers.



**Figure 4:** The hypothalamo-neurohypophysial system (HNS) and-immune network. The HNS including both the OXT- and VP-secreting systems has close interactions with the immune organs. In response to the challenge of pathogens and immune cytokines (1), OXT-, VP- and the hypothalamic-pituitary-adrenal (HPA) axis are all activated (2). OXT inhibits and VP activates HPA axis (3), which inhibits both OXT- and VP-neuronal activity (4). In general, OXT promotes the development of immune organs but inhibits inflammatory reactions (5); (79); VP inhibits the activity of the thymus and facilitates the functions of the spleen while promoting inflammatory responses (6). A functioning thymus is also a critical factor of the integrity of the HNS (7). In addition, OXT (8) and VP (9) can reduce and increase the sensitivity to pathogens through promoting social activity and autonomic outflow, respectively.


Recently, an OB-HNS route has emerged as a potential approach mediating intranasal drug effects on brain activities. In general, intranasally applied drugs can get into the brain bypassing the BBB, through diffusion via the olfactory cells and neural sheathes, through circulation or by activating trigeminal neurons [106]. Nevertheless, drugs diffusing from the olfactory epithelium into the brain get into and accumulate in the OB first [107]. Moreover, OB cells also express OXTRs [108] and VP receptors [109] while circulating neuropeptides are less likely acting on the HNS [110]. Thus, the OB could be the first brain target of nasally-applied neuropeptides and can be the only direct target if these neuropeptide doses are lower enough [111-112]. Relative to the extensively identified brain effects of nasally-applied OXT, the effects of nasal VP are relatively weak and limited [113] and thus, exploring OXT-associated OB-SON pathway likely provides a useful model to study the mechanism underlying the actions of intranasal neuropeptides.

In fact, an OB-SON neural circuit has been well established based on close interactions of OBs with the SON. Stimulation of the OB increases activity of supraoptic neurons [114,115] via the lateral olfactory tract [116]. Conversely, increased OXT neuronal activity causes OXT release in the OB via neural terminals from the SON or its associated interneurons [13]. By activation of OXTRs in the OB, OXT can change the electrical activity of mitral cells [24], the major output neurons in the OB. Thus, there are bidirectional interactions between the OB and the SON, which allows nasal OXT to modulate the activity of OB neurons and subsequently change the activity of MNCs in the SON [117]. Since OXTR has very high sensitivity to low levels of OXT [118], if nasally-applied OXT is controlled at appropriate levels, it is possible to elicit a "specific" effect on SON neuronal activity. This is particularly meaningful to avoid unwanted effects when OXTRs are known to express in extensive brain areas and peripheral tissues [8].

Nasal therapy is a simple yet effective method for medical modulation of brain activity. For instance, OXT administration can greatly attenuate the cortisol response to laboratory tasks that strongly activate the HPA-axis, which is more robust among clinical populations concomitant with social difficulties [119], thereby exerting an anxiolytic effect. These actions could be mediated by the SON. As stated above, axonal collaterals and dendrites of the SON neurons can innervate extensive brain areas directly or through its perinuclear zone neurons or the cerebroventricular system [16,120]. Thus, brain effects of intranasal neuropeptides could be mediated by a nose-OB-SON-brain approach. To achieve a specific effect, future studies are required to identify the distribution of different neuropeptide receptors, the unique or major pathway from the OB to a brain target area, and the relative power of the main OB versus vomeronasal organ [121]. Figure 5 shows a putative nose-brain approach mediated by the OB-HNS route.

### Clinically relevant studies

The HNS is extensively involved in a variety of physiological functions while its pathological implications are also explored [122,123]. For example, VP is involved in stress, the hypophysial-adrenocortical axis, neuroimmune responses, renal function and corticotroph pituitary tumors [124]. Similarly, OXT is extensively involved in many body functions such as sexual activity, penile erection, ejaculation, pregnancy, uterus contraction, milk ejection, maternal behavior, osteoporosis, diabetes, cancer, social bonding, and stress among many others [125,126]. Thus, OXT and its receptor have been considered as potential targets for drug therapy of diseases in the brain and peripheral organs [127]. Here, we discuss the clinical relevance of OXT and VP in association with their classical functions.



Abnormality in the VP-secreting system can result in multiple disorders in body functions, typically seen in diabetes insipidus (DI) and the syndrome of inappropriate antidiuretic hormone secretion (SIADH) [128]. DI is characterized by hypotonic polyuria greater than 3 liters/24 hours in adults and persisting even during water deprivation due to a defect in VP synthesis (central DI), VP resistance (nephrogenic DI), or early decomposition of VP by placental enzymes (gestational DI). For the central DI, desmopressin, a synthetic analog of vasopressin, is the treatment option [129]. Conversely, overly produced VP can cause SIADH. SIADH is a condition that the body releases excessive VP from the posterior pituitary or another source, which induces too much water to remain inside the body and results in hypervolemia along with dilutional hyponatremia. In SIADH, physiological mechanisms regulating VP secretion are disrupted, either because of the activation of neuroendocrine tumors or non-osmotic regulatory processes, and thus, the synthesis and release of VP are not inhibited by a reduction in plasma osmolality [43]. Thus, tolvaptan, a VP antagonist can be used along with fluid restriction and diuretics to resolve the fluid retention in the SIADH [130].

Recently, the involvement of VP hypersecretion in ischemic stroke has also been highlighted. In ischemic stroke, brain edema formation is one of the most important pathologic processes [131], in which VP exerts a pivotal role [132,133]. VP release and expression of its receptors in the brain increase significantly following ischemia, trauma or subarachnoid hemorrhage in patients and in animal models of these diseases [134]. VP can increase ion transport across the cell membrane and the blood-brain barrier [135] while increasing AQP4 expression in astrocytes and thus promoting cytotoxic edema [136]. In parallel with the direct effect on hydromineral balance, excessive VP also causes brain arteriole constriction and worsens tissue hypoxia [137], thereby worsening brain edema through oxygen and glucose deprivation. These and other associated factors [138-140] can lead to disruption of the blood-brain barrier and increase hydromineral transfer into the brain, leading to further increases in intracranial pressure and neuron damages. By increasing water retention, cytotoxic edema, disruption of the blood-brain barrier and worsening ischemic injury, VP hypersecretion becomes a major cause of brain injury and disability following stroke. However, mechanisms underlying VP hypersecretion remain unclear [105] although the SON and PVN are known to be

involved in the infarct zone in rats [141]. Clarification the mechanisms underlying VP hypersecretion during stroke and suppression of its hypersecretion or the effect of VP receptor signaling processes remain a serious challenge in stroke treatment.

In parallel with the progress in studying the pathological roles of VP, OXT-associated diseases are also explored. Parturition and lactation are the two classical functions of OXT. OXT knockout does not influence parturition; however lactation or breastfeeding is clearly disrupted as shown in animal experiments [142]. Thus, roles of OXT in lactation failure and its associated diseases are discussed here. Breastfeeding has many beneficial effects on the health of mothers and the babies. However, insufficient breastfeeding and lactation failure are common socioeconomic and biomedical issues among the mothers. Insufficient breastfeeding is associated with the high incidence of postpartum depression, premenopausal breast cancer, diabetes and obesity in the mothers and autism, sudden death, and deficiency in maternal behaviors in their offspring [143]. Thus, clarification of the mechanisms underlying lactation failure and exploration of its potential therapeutic approaches are important for the health of both the mothers and the babies.

In general, cesarean section, maternal illness during pregnancy, and preterm birth are considered as the major factors that affect early breastfeeding status [144]. These factors can disrupt breastfeeding through the following approaches. 1) Pain and stress. Factors leading to lactation failure are commonly associated with pain and stress, physically or mentally. For example, cesarean section increases the incidence of postnatal depression, psychological birth trauma, birth-related post-traumatic stress disorder and postoperative pain [145], which can cause breastfeeding failure through inhibition of OXT neuronal activity by the activated HPA axis. Consistently, maternal separation causes postpartum depression, leading to maternal loss of interests toward her offspring and lactation failure [146]. 2) Immaturity of the OXT-secreting system for lactation. Around the term, the SON experiences a significant morphological and functional remodeling [25,26] that facilitates milk-ejection reflex [147]. However, maternal separation days after the parturition can reverse the facilitatory morphologic changes in the hypothalamus [148] and results in reduction of the secretion of prolactin [149] and OXT [150] along with increased corticosteroid levels [151]. Thus, mothers with preterm babies or separated from the baby could delay the proceeding of this maturation, particularly in the mother with preterm baby. 3) Failure in maintenance of the functional state of the structural basis for the milk-ejection reflex. Suckling stimulates the release of OXT and milk removal [152] while increasing the secretion of prolactin as well [153]. At cellular and molecular levels, without sufficient suckling, OXT synthesis in OXT neurons decreases and OXTRs uncouple with their downstream signals as observed in a maternal separation model of lactating rats [146]. Failure of suckling also causes accumulation of milk in breast ductal system that increases intramammary pressure, blocks milk secretion, elicits oxidative stress and results in the involution of breast tissues [148]. Thus, lacking suckling stimulation not only induces the involution of the OXT-secreting system and the mammary glands, but also disrupts the structural basis for the milk-ejection reflex.

Importantly, breastfeeding failure seeds the risk of breast cancer. A cohort study of parous premenopausal women indicated that breastfeeding is inversely associated with the incidence of breast cancer among women with a family history of breast cancer [154]. However, the biological basis of this anticancer effect of breastfeeding has not gained support of experimental evidence in spite of the

presence of many hypotheses. Recently, an animal experiment using phosphorylated extracellular signal-regulated protein kinase 1/2 and cyclooxygenase 2 as indicators of hydrogen peroxide-evoked proliferative activity revealed that lactation failure resulting from lack of pulsatile OXT release during lactation is causally related to the precancerous lesions of the mammary glands [155]. This finding, although requiring observation of long-term effects on the incidence of mammary gland tumorigenesis, suggests that intermittently pulsatile action of OXT during lactation underlies the anti-pre-menopausal breast cancer effects of breastfeeding.

## Concluding Remarks

Studies on the MNCs have greatly enriched our knowledge of the peptidergic neurons in modern neuroscience. These include not only the progress in the studies on the classical roles of the HNS, such as neuronal-glia interactions and function-associated morphological plasticity, but also exploration of the scaffolding and guiding roles of glial fibrillary acidic protein in astrocytic plasticity, cellular and molecular mechanisms underlying different firing patterns, the pivotal position of the HNS-immune network in neuroendocrine regulation of immune activity, and the OB-HNS route that serves as a key mediator of nasal drug effects on brain activity. The studies also highlight the translational potentials of modulating HNS activity, such as suppression of VP signaling process to control the pathogenesis of ischemic stroke and utilizing the pulsatile feature of OXT actions in prevention of premenopausal breast cancer. Nevertheless, a number of major questions remain to be addressed, which include but not limited to: 1) The details of neural connections between the HNS and other brain structures and between different nuclei in the HNS; 2) The compensatory mechanisms to reserve the “lost” functions in gene deficient animals, particularly in OXT and OXTR gene knockout mice [156]; and 3) Exploration of the optimal approaches and therapies to treat the HNS-associated diseases. To achieve these goals, it is necessary to apply integrative research approach in the studies on the HNS to build an intact picture from individual molecular events to whole body’s functions/behaviors. It is also important to present “more comparable” data by adopting uniformed and optimized experimental conditions in individual laboratories. By clarifying these questions and resolving emerging challenges, we can better understand the functions of the HNS and their translational potentials, and in turn provide important insights into common disease states.

## Acknowledgements

We thank Dr. Stephani C. Wang for critical reading. The project is sponsored by the National Natural Science Foundation of China (grant No. 31471113, YFW), and the higher education talents funds of Heilongjiang province (grant No. 002000154). All authors declare no conflict of interest.

## References

1. Watts AG (2011) Structure and function in the conceptual development of mammalian neuroendocrinology between 1920 and 1965. *Brain Res Rev* 66: 174-204.
2. Ottenhausen M, Bodhinayake I, Banu MA, Stieg PE, Schwartz TH (2016) Vincent du Vigneaud: following the sulfur trail to the discovery of the hormones of the posterior pituitary gland at Cornell Medical College. *J Neurosurg* 124: 1538-1542.
3. Burbach JP, Luckman SM, Murphy D, Gainer H (2001) Gene regulation in the magnocellular hypothalamo-neurohypophysial system. *Physiol Rev* 81: 1197-1267.
4. Murphy D, Konopacka A, Hindmarch C, Paton JF, Sweedler JV, et al. (2012) The hypothalamo-neurohypophysial system: from genome to physiology. *J Neuroendocrinol* 24: 539-553.
5. Leng G, Pineda R, Sabatier N, Ludwig M (2015) 60 Years of Neuroendocrinology: The posterior pituitary, from Geoffrey Harris to our present understanding. *J Endocrinol* 226: T173-T185.
6. Brownstein MJ, Russell JT, Gainer H (1980) Synthesis, transport, and release of posterior pituitary hormones. *Science* 207: 373-378.
7. Sofroniew MV, Glasmann W (1981) Golgi-like immunoperoxidase staining of hypothalamic magnocellular neurons that contain vasopressin, oxytocin or neurophysin in the rat. *Neuroscience* 6: 619-643.
8. Gimpl G, Fahrenholz F (2001) The oxytocin receptor system: structure, function, and regulation. *Physiol Rev* 81: 629-683.
9. Shibuya I, Kabashima N, Ibrahim N, Setiadiji SV, Ueta Y, et al. (2000) Pre- and postsynaptic modulation of the electrical activity of rat supraoptic neurones. *Exp Physiol* 85: 145S-151S.
10. Hatton GI (1990) Emerging concepts of structure-function dynamics in adult brain: the hypothalamo-neurohypophysial system. *Prog Neurobiol* 34: 437-504.
11. Pittman QJ, Bains J (2003) Backtalk in neurons. *Trends Endocrinol Metab* 14: 2-3.
12. Crowley WR (2015) Neuroendocrine regulation of lactation and milk production. *Compr Physiol* 5: 255-291.
13. Alonso G, Szafarczyk A, Assenmacher I (1986) Radioautographic evidence that axons from the area of supraoptic nuclei in the rat project to extrahypothalamic brain regions. *Neurosci Lett* 66: 251-256.
14. Inyushkin AN, Orlans HO, Dyball RE (2009) Secretory cells of the supraoptic nucleus have central as well as neurohypophysial projections. *J Anat* 215: 425-434.
15. Caligioni CS, Oliver C, Jamur MC, Franci CR (2007) Presence of oxytocin receptors in the gonadotrophin-releasing hormone (GnRH) neurones in female rats: a possible direct action of oxytocin on GnRH neurones. *J Neuroendocrinol* 19: 439-448.
16. Ju G, Liu S, Tao J (1986) Projections from the hypothalamus and its adjacent areas to the posterior pituitary in the rat. *Neuroscience* 19: 803-828.
17. Neumann I, Ludwig M, Engelmann M, Pittman QJ, Landgraf R (1993) Simultaneous microdialysis in blood and brain: oxytocin and vasopressin release in response to central and peripheral osmotic stimulation and suckling in the rat. *Neuroendocrinology* 58: 637-645.
18. Moos F, Richard P (1989) Paraventricular and supraoptic bursting oxytocin cells in rat are locally regulated by oxytocin and functionally related. *J Physiol* 408: 1-18.
19. Eliava M, Melchior M, Knobloch-Bollmann HS, Wahis J, da Silva Gouveia M, et al. (2016) A New Population of Parvocellular Oxytocin Neurons Controlling Magnocellular Neuron Activity and Inflammatory Pain Processing. *Neuron* 89: 1291-1304.
20. Ludwig M, Stern J (2015) Multiple signalling modalities mediated by dendritic exocytosis of oxytocin and vasopressin. *Philos Trans R Soc Lond B Biol Sci* 370.
21. Wang YF, Negoro H, Honda K (1995) Effects of hemitransection of the midbrain on milk-ejection burst of oxytocin neurones in lactating rat. *J Endocrinol* 144: 463-470.
22. Buijs RM (1978) Intra- and extrahypothalamic vasopressin and oxytocin pathways in the rat. Pathways to the limbic system, medulla oblongata and spinal cord. *Cell Tissue Res* 192: 423-435.
23. Liu XY, Hou D, Wang J, Lv C, Jia S, et al. (2016) Expression of glial fibrillary acidic protein in astrocytes of rat supraoptic nucleus throughout estrous cycle. *Neuro Endocrinol Lett* 37: 41-45.
24. Yu GZ, Kaba H, Okutani F, Takahashi S, Higuchi T, et al. (1996) The action of oxytocin originating in the hypothalamic paraventricular nucleus on mitral and granule cells in the rat main olfactory bulb. *Neuroscience* 72: 1073-1082.
25. Tweedle CD, Hatton GI (1977) Ultrastructural changes in rat hypothalamic neurosecretory cells and their associated glia during minimal dehydration and rehydration. *Cell Tissue Res* 181: 59-72.
26. Theodosios DT, Chapman DB, Montagnese C, Poulain DA, Morris JF (1986) Structural plasticity in the hypothalamic supraoptic nucleus at lactation affects oxytocin-, but not vasopressin-secreting neurones. *Neuroscience* 17: 661-678.
27. Chapman DB, Theodosios DT, Montagnese C, Poulain DA, Morris JF (1986) Osmotic stimulation causes structural plasticity of neurone-glia relationships of the oxytocin but not vasopressin secreting neurones in the hypothalamic supraoptic nucleus. *Neuroscience* 17: 679-686.

28. Shi L, Fan Y, Xu Z (2012) Development of oxytocin- and vasopressin-network in the supraoptic and paraventricular nuclei of fetal sheep. *Physiol Res* 61: 277-286.
29. Armstrong WE, Stern JE (1998) Phenotypic and state-dependent expression of the electrical and morphological properties of oxytocin and vasopressin neurones. *Prog Brain Res* 119: 101-113.
30. Rossoni E, Feng J, Tirozzi B, Brown D, Leng G, et al. (2008) Emergent synchronous bursting of oxytocin neuronal network. *PLoS Comput Biol* 4: e1000123.
31. Catheline G, Touquet B, Lombard MC, Poulain DA, Theodosis DT (2006) A study of the role of neuro-glial remodeling in the oxytocin system at lactation. *Neuroscience* 137: 309-316.
32. Wang YF, Hamilton K (2009) Chronic vs. acute interactions between supraoptic oxytocin neurons and astrocytes during lactation: role of glial fibrillary acidic protein plasticity. *ScientificWorldJournal* 9: 1308-1320.
33. Tasker JG, Oliet SH, Bains JS, Brown CH, Stern JE (2012) Glial regulation of neuronal function: from synapse to systems physiology. *J Neuroendocrinol* 24: 566-576.
34. Wang YF, Zhu H (2014) Mechanisms underlying astrocyte regulation of hypothalamic neuroendocrine neuron activity. *Sheng Li Ke Xue Jin Zhan* 45: 177-184.
35. Pekny M, Wilhelmsson U, Bogestål YR, Pekna M (2007) The role of astrocytes and complement system in neural plasticity. *Int Rev Neurobiol* 82: 95-111.
36. Salm AK, Smithson KG, Hatton GI (1985) Lactation-associated redistribution of the glial fibrillary acidic protein within the supraoptic nucleus. An immunocytochemical study. *Cell Tissue Res* 242: 9-15.
37. Wang YF, Hatton GI (2009) Astrocytic plasticity and patterned oxytocin neuronal activity: dynamic interactions. *J Neurosci* 29: 1743-1754.
38. Hawrylak N, Boone D, Salm AK (1999) The surface density of glial fibrillary acidic protein immunopositive astrocytic processes in the rat supraoptic nucleus is reversibly altered by dehydration and rehydration. *Neurosci Lett* 277: 57-60.
39. Wang YF, Parpura V (2016) Central Role of Maladapted Astrocytic Plasticity in Ischemic Brain Edema Formation. *Frontiers in Cellular Neuroscience* 10.
40. Vargova L, Sykova E (2014) Astrocytes and extracellular matrix in extrasynaptic volume transmission. *Philos Trans R Soc Lond B Biol Sci* 369: 20130608.
41. Wang YF, Sun MY, Hou Q, Hamilton KA (2013) GABAergic inhibition through synergistic astrocytic neuronal interaction transiently decreases vasopressin neuronal activity during hypoosmotic challenge. *Eur J Neurosci* 37: 1260-1209.
42. Wang YF, Sun MY, Hou Q, Parpura V (2013) Hyposmolality differentially and spatiotemporally modulates levels of glutamine synthetase and serine racemase in rat supraoptic nucleus. *Glia* 61: 529-538.
43. Wang YF, Liu LX, Yang HP (2011) Neurophysiological involvement in hypervolemic hyponatremia-evoked by hypersecretion of vasopressin. *Translational Biomedicine* 2: 3.
44. Hertz L, Xu J, Chen Y, Gibbs ME, Du T, et al. (2014) Antagonists of the Vasopressin V1 Receptor and of the beta(1)-Adrenoceptor Inhibit Cytotoxic Brain Edema in Stroke by Effects on Astrocytes - but the Mechanisms Differ. *Curr Neuropharmacol* 12: 308-323.
45. Coles JA, Poulain DA (1991) Extracellular K<sup>+</sup> in the supraoptic nucleus of the rat during reflex bursting activity by oxytocin neurones. *J Physiol* 439: 383-409.
46. Leng G, Shibuki K, Way SA (1988) Effects of raised extracellular potassium on the excitability of, and hormone release from, the isolated rat neurohypophysis. *J Physiol* 399: 591-605.
47. Wang YF, Negoro H, Honda K (1996) Milk ejection bursts of supraoptic oxytocin neurones during bilateral and unilateral suckling in the rat. *J Neuroendocrinol* 8: 427-431.
48. Zamanian JL, Xu L, Foo LC, Nouri N, Zhou L, et al. (2012) Genomic analysis of reactive astrogliosis. *J Neurosci* 32: 6391-6410.
49. Dutton A, Dyball RE (1979) Phasic firing enhances vasopressin release from the rat neurohypophysis. *J Physiol* 290: 433-440.
50. Brimble MJ, Dyball RE (1977) Characterization of the responses of oxytocin- and vasopressin-secreting neurones in the supraoptic nucleus to osmotic stimulation. *J Physiol* 271: 253-271.
51. Lincoln DW, Wakerley JB (1975) Factors governing the periodic activation of supraoptic and paraventricular neurosecretory cells during suckling in the rat. *J Physiol* 250: 443-461.
52. Summerlee AJ (1983) Hypothalamic neurone activity: hormone release and behaviour in freely moving rats. *Q J Exp Physiol* 68: 505-515.
53. Belin V, Moos F (1986) Paired recordings from supraoptic and paraventricular oxytocin cells in suckled rats: recruitment and synchronization. *J Physiol* 377: 369-390.
54. Higuchi T, Tadokoro Y, Honda K, Negoro H (1986) Detailed analysis of blood oxytocin levels during suckling and parturition in the rat. *J Endocrinol* 110: 251-256.
55. Wang YF, Hatton GI (2004) Milk ejection burst-like electrical activity evoked in supraoptic oxytocin neurons in slices from lactating rats. *J Neurophysiol* 91: 2312-2321.
56. Brown CH, Bains JS, Ludwig M, Stern JE (2013) Physiological regulation of magnocellular neurosecretory cell activity: integration of intrinsic, local and afferent mechanisms. *J Neuroendocrinol* 25: 678-710.
57. Armstrong WE, Wang L, Li C, Teruyama R (2010) Performance, properties and plasticity of identified oxytocin and vasopressin neurones in vitro. *J Neuroendocrinol* 22: 330-342.
58. Augustine RA, Bouwer GT, Seymour AJ, Grattan DR, Brown CH (2016) Reproductive Regulation of Gene Expression in the Hypothalamic Supraoptic and Paraventricular Nuclei. *J Neuroendocrinol* 28.
59. Sladek CD, Song Z (2012) Diverse roles of G-protein coupled receptors in the regulation of neurohypophysial hormone secretion. *J Neuroendocrinol* 24: 554-565.
60. Iovino M, Guastamacchia E, Giagulli VA, Licchelli B, Iovino E, et al. (2014) Molecular mechanisms involved in the control of neurohypophysial hormones secretion. *Curr Pharm Des* 20: 6702-6713.
61. Prager-Khoutorsky M, Bourque CW (2015) Mechanical basis of osmosensory transduction in magnocellular neurosecretory neurones of the rat supraoptic nucleus. *J Neuroendocrinol* 27: 507-515.
62. Wakerley JB, Lincoln DW (1973) The milk-ejection reflex of the rat: a 20- to 40-fold acceleration in the firing of paraventricular neurones during oxytocin release. *J Endocrinol* 57: 477-493.
63. Moos F, Poulain DA, Rodriguez F, Guerné Y, Vincent JD, et al. (1989) Release of oxytocin within the supraoptic nucleus during the milk ejection reflex in rats. *Exp Brain Res* 76: 593-602.
64. Wang YF, Negoro H, Higuchi T (2013) Lesions of hypothalamic mammillary body desynchronize milk-ejection bursts of rat bilateral supraoptic oxytocin neurones. *J Neuroendocrinol* 25: 67-75.
65. Oliveira GR, Franci CR, Rodovalho GV, Franci JA, Morris M, et al. (2004) Alterations in the central vasopressin and oxytocin axis after lesion of a brain osmotic sensory region. *Brain Res Bull* 63: 515-520.
66. Stern JE, Armstrong WE (1996) Changes in the electrical properties of supraoptic nucleus oxytocin and vasopressin neurons during lactation. *J Neurosci* 16: 4861-4871.
67. Armstrong WE, Stern JE (1998) Electrophysiological distinctions between oxytocin and vasopressin neurons in the supraoptic nucleus. *Adv Exp Med Biol* 449: 67-77.
68. da Silva MP, Merino RM, Mecawi AS, Moraes DJ, Varanda WA (2015) In vitro differentiation between oxytocin- and vasopressin-secreting magnocellular neurons requires more than one experimental criterion. *Mol Cell Endocrinol* 400: 102-111.
69. Brown CH, Ruan M, Scott V, Tobin VA, Ludwig M (2008) Multi-factorial somatodendritic regulation of phasic spike discharge in vasopressin neurons. *Prog Brain Res* 170: 219-228.
70. Wang YF, Hatton GI (2005) Burst firing of oxytocin neurons in male rat hypothalamic slices. *Brain Res* 1032: 36-43.
71. Wang YF, Hatton GI (2007) Dominant role of betagamma subunits of G-proteins in oxytocin-evoked burst firing. *J Neurosci* 27: 1902-1912.
72. Cosgrave AS, Richardson CM, Wakerley JB (2000) Permissive effect of centrally administered oxytocin on the excitatory response of oxytocin neurones to ventral tegmental stimulation in the suckled lactating rat. *J Neuroendocrinol* 12: 843-852.

73. Israel JM, Le Masson G, Theodosios DT, Poulain DA (2003) Glutamatergic input governs periodicity and synchronization of bursting activity in oxytocin neurons in hypothalamic organotypic cultures. *Eur J Neurosci* 17: 2619-2629.
74. Moos FC (1995) GABA-induced facilitation of the periodic bursting activity of oxytocin neurones in suckled rats. *J Physiol* 488: 103-114.
75. Hatton GI, Wang YF (2008) Neural mechanisms underlying the milk ejection burst and reflex. *Prog Brain Res* 170: 155-166.
76. Popescu IR, Morton LA, Franco A, Di S, Ueta Y, et al. (2010) Synchronized bursts of miniature inhibitory postsynaptic currents. *J Physiol* 588: 939-951.
77. Wang YF, Hatton GI (2007) Interaction of extracellular signal-regulated protein kinase 1/2 with actin cytoskeleton in supraoptic oxytocin neurons and astrocytes: role in burst firing. *J Neurosci* 27: 13822-13834.
78. Dine J, Ducourneau VR, Fénelon VS, Fossat P, Amadio A, et al. (2014) Extracellular signal-regulated kinase phosphorylation in forebrain neurones contributes to osmoregulatory mechanisms. *J Physiol* 592: 1637-1654.
79. Prager-Khoutorsky M, Bourque CW (2010) Osmosensation in vasopressin neurons: changing actin density to optimize function. *Trends Neurosci* 33: 76-83.
80. Miyata S, Khan AM, Hatton GI (1998) Colocalization of calretinin and calbindin-D28k with oxytocin and vasopressin in rat supraoptic nucleus neurons: a quantitative study. *Brain Res* 785: 178-182.
81. Li Z, Decavel C, Hatton GI (1995) Calbindin-D28k: role in determining intrinsically generated firing patterns in rat supraoptic neurones. *J Physiol* 488: 601-608.
82. Woudenberg-Vrenken TE, Bindels RJ, Hoenderop JG (2009) The role of transient receptor potential channels in kidney disease. *Nat Rev Nephrol* 5: 441-449.
83. Webster EL, Torpy DJ, Elenkov IJ, Chrousos GP (1998) Corticotropin-releasing hormone and inflammation. *Ann N Y Acad Sci* 840: 21-32.
84. Uchoa ET, Aguilera G, Herman JP, Fiedler JL, Deak T, et al. (2014) Novel aspects of glucocorticoid actions. *J Neuroendocrinol* 26: 557-572.
85. Ficek W (1982) The physiological relationship between the hypothalamus and thymus of Wistar rats. I. Some morphological and microstructural changes in the hypothalamus in newborn rats following thymectomy. *Gegenbaurs Morphol Jahrb* 128: 732-740.
86. Yang H, Wang L, Ju G (1997) Evidence for hypothalamic paraventricular nucleus as an integrative center of neuroimmunomodulation. *Neuroimmunomodulation* 4: 120-127.
87. McQuaid RJ, McInnis OA, Abizaid A, Anisman H (2014) Making room for oxytocin in understanding depression. *Neurosci Biobehav Rev* 45: 305-322.
88. Wang P, Yang HP, Tian S, Wang L, Wang SC, et al. (2015) Oxytocin-secreting system: A major part of the neuroendocrine center regulating immunologic activity. *J Neuroimmunol* 289: 152-161.
89. Ficek W (1983) Physiological dependency between the hypothalamus and the thymus of Wistar rats. IV. Organotypic culture of the thymus in the presence of hypophysial hormones, vasopressin, and oxytocin. *Gegenbaurs Morphol Jahrb* 129: 445-458.
90. Gobrogge K, Wang Z (2015) Neuropeptidergic regulation of pair-bonding and stress buffering: Lessons from voles. *Horm Behav* 76: 91-105.
91. Hansenne I (2005) Thymic transcription of neurohypophysial and insulin-related genes: impact upon T-cell differentiation and self-tolerance. *J Neuroendocrinol* 17: 321-327.
92. Deing V, Roggenkamp D, Kuhn J, Gruschka A, Stab F, et al. (2013) Oxytocin modulates proliferation and stress responses of human skin cells: implications for atopic dermatitis. *Exp Dermatol* 22: 399-405.
93. Kim SH, MacIntyre DA, Firmino Da Silva M, Blanks AM, Lee YS, et al. (2015) Oxytocin activates NF- $\kappa$ B-mediated inflammatory pathways in human gestational tissues. *Mol Cell Endocrinol* 403: 64-77.
94. Johnson HM, Torres BA (1988) A novel arginine vasopressin-binding peptide that blocks arginine vasopressin modulation of immune function. *J Immunol* 141: 2420-2423.
95. Szeto A, Nation DA, Mendez AJ, Dominguez-Bendala J, Brooks LG, et al. (2008) Oxytocin attenuates NADPH-dependent superoxide activity and IL-6 secretion in macrophages and vascular cells. *Am J Physiol Endocrinol Metab* 295: E1495-E501.
96. Elands J, Resink A, De Kloet ER (1990) Neurohypophysial hormone receptors in the rat thymus, spleen, and lymphocytes. *Endocrinology* 126: 2703-2710.
97. Tsai MS, Hwang SM, Chen KD, Lee YS, Hsu LW, et al. (2007) Functional network analysis of the transcriptomes of mesenchymal stem cells derived from amniotic fluid, amniotic membrane, cord blood, and bone marrow. *Stem Cells* 25: 2511-2523.
98. Hansenne I, Rasier G, Charlet-Renard CH, DeFresne MP, Greimers R, et al. (2004) Neurohypophysial receptor gene expression by thymic T cell subsets and thymic T cell lymphoma cell lines. *Clin Dev Immunol* 11: 45-51.
99. Di S, Tasker JG (2008) Rapid synapse-specific regulation of hypothalamic magnocellular neurons by glucocorticoids. *Prog Brain Res* 170: 379-388.
100. Pereira AS, Giusti-Paiva A, Vilela FC (2015) Central corticosterone disrupts behavioral and neuroendocrine responses during lactation. *Neurosci Lett* 60: 688-693.
101. Spanakis EK, Wand GS, Ji N, Golden SH (2016) Association of HPA axis hormones with copeptin after psychological stress differs by sex. *Psychoneuroendocrinology* 63: 254-261.
102. Smith AS, Tabbaa M, Lei K, Eastham P, Butler MJ, et al. (2015) Local oxytocin tempers anxiety by activating GABA receptors in the hypothalamic paraventricular nucleus. *Psychoneuroendocrinology* 63: 50-58.
103. Decker DA, Grant C, Oh L, Becker PM, Young D, et al. (2014) Immunomodulatory effects of H.P. Acthar Gel on B cell development in the NZB/W F1 mouse model of systemic lupus erythematosus. *Lupus* 23: 802-812.
104. Pittman QJ (2011) A neuro-endocrine-immune symphony. *J Neuroendocrinol* 23: 1296-1297.
105. Jia SW, Liu XY, Wang SC, Wang YF (2016) Vasopressin Hypersecretion-Associated Brain Edema Formation in Ischemic Stroke: Underlying Mechanisms. *J Stroke Cerebrovasc Dis* 25: 1289-1300.
106. Liu Y, Liu X (2015) Intranasal delivery: circumventing the iron curtain to treat neurological disorders. *Expert Opin Drug Deliv* 12: 1717-1725.
107. Dhuria SV, Hanson LR, Frey WH (2010) Intranasal delivery to the central nervous system: mechanisms and experimental considerations. *J Pharm Sci* 99: 1654-1673.
108. Meddle SL, Bishop VR, Gkoumassi E, van Leeuwen FW, Douglas AJ (2007) Dynamic changes in oxytocin receptor expression and activation at parturition in the rat brain. *Endocrinology* 148: 5095-5104.
109. Stevenson EL, Caldwell HK (2012) The vasopressin 1b receptor and the neural regulation of social behavior. *Horm Behav* 61: 277-282.
110. Leng G, Russell JA (2016) The Peptide Oxytocin Antagonist F-792, When Given Systemically, Does Not Act Centrally in Lactating Rats. *J Neuroendocrinol* 28.
111. Leng G, Ludwig M (2016) Intranasal Oxytocin: Myths and Delusions. *Biol Psychiatry* 79: 243-250.
112. Freeman SM, Samineni S, Allen PC, Stockinger D, Bales KL, et al. (2016) Plasma and CSF oxytocin levels after intranasal and intravenous oxytocin in awake macaques. *Psychoneuroendocrinology* 66: 185-194.
113. Ludwig M, Tobin VA, Callahan MF, Papadaki E, Becker A, et al. (2013) Intranasal application of vasopressin fails to elicit changes in brain immediate early gene expression, neural activity and behavioural performance of rats. *J Neuroendocrinol* 25: 655-667.
114. Hatton GI, Yang QZ (1990) Activation of excitatory amino acid inputs to supraoptic neurons. I. Induced increases in dye-coupling in lactating, but not virgin or male rats. *Brain Res* 513: 264-269.
115. Modney BK, Yang QZ, Hatton GI (1990) Activation of excitatory amino acid inputs to supraoptic neurons. II. Increased dye-coupling in maternally behaving virgin rats. *Brain Res* 513: 270-273.
116. Smithson KG, Weiss ML, Hatton GI (1989) Supraoptic nucleus afferents from the main olfactory bulb--I. Anatomical evidence from anterograde and retrograde tracers in rat. *Neuroscience* 31: 277-287.
117. Price JL, Slotnick BM, Revial MF (1991) Olfactory projections to the hypothalamus. *J Comp Neurol* 306: 447-461.
118. Wang YF, Ponzio TA, Hatton GI (2006) Autocrine effects of progressively rising oxytocin concentrations on supraoptic oxytocin neuronal activity in slices from lactating rats. *Am J Physiol Regul Integr Comp Physiol* 290: R1191-R1198.

119. Cardoso C, Kingdon D, Ellenbogen MA (2014) A meta-analytic review of the impact of intranasal oxytocin administration on cortisol concentrations during laboratory tasks: moderation by method and mental health. *Psychoneuroendocrinology* 49: 161-170.
120. Veening JG, de Jong T, Barendregt HP (2010) Oxytocin-messages via the cerebrospinal fluid: behavioral effects; a review. *Physiol Behav* 101: 193-210.
121. Matsunami H, Amrein H (2003) Taste and pheromone perception in mammals and flies. *Genome Biol* 4: 220.
122. Calabro RS, Italiano D, Ferrara D, Mondello S, Conti-Nibali V, et al. (2012) The hypothalamic-neurohypophysial system: current and future treatment of vasopressin and oxytocin related disorders. *Recent Pat Endocr Metab Immune Drug Discov* 6: 235-250.
123. Meyer-Lindenberg A, Domes G, Kirsch P, Heinrichs M (2011) Oxytocin and vasopressin in the human brain: social neuropeptides for translational medicine. *Nat Rev Neurosci* 12: 524-538.
124. Rotondo F, Butz H, Syro LV, Yousef GM, Di Ieva A, et al. (2016) Arginine vasopressin (AVP): a review of its historical perspectives, current research and multifunctional role in the hypothalamo-hypophysial system. *Pituitary*.
125. Yang HP, Wang L, Han L, Wang SC (2013) Nonsocial functions of hypothalamic oxytocin. *ISRN Neurosci* 2013: 179272.
126. Ramos L, Hicks C, Kevin R, Caminer A, Narlawar R, et al. (2013) Acute prosocial effects of oxytocin and vasopressin when given alone or in combination with 3,4-methylenedioxymethamphetamine in rats: involvement of the V1A receptor. *Neuropsychopharmacology* 38: 2249-2259.
127. Viero C, Shibuya I, Kitamura N, Verkhatsky A, Fujihara H, et al. (2010) REVIEW: Oxytocin: Crossing the bridge between basic science and pharmacotherapy. *CNS Neurosci Ther* 16: e138-e156.
128. Verbalis JG (2014) Disorders of water metabolism: diabetes insipidus and the syndrome of inappropriate antidiuretic hormone secretion. *Handbook of Clinical Neurology* 124: 37-52.
129. Lubecka EA, Sikorska E, Sobolewski D, Prah A, Slaninová J, et al. (2016) Potent antidiuretic agonists, deamino-vasopressin and desmopressin, and their inverse analogs: NMR structure and interactions with micellar and liposomal models of cell membrane. *Biopolymers* 106: 245-259.
130. Ichimura S, Fahlbusch R, Ludemann W (2015) Treatment of Hyponatremia with Tolvaptan in a Patient after Neurosurgical Treatment of a Pituitary Tumor: Case Report and Review of Literature. *J Neurol Sur Rep* 76: e279-e281.
131. Zador Z, Stiver S, Wang V, Manley GT (2009) Role of aquaporin-4 in cerebral edema and stroke. *Handb Exp Pharmacol*, pp: 159-170.
132. Rutkowski JM, Wallace BK, Wise PM, O'Donnell ME (2011) Effects of estradiol on ischemic factor-induced astrocyte swelling and AQP4 protein abundance. *Am J Physiol Cell Physiol* 301: C204-C212.
133. O'Donnell ME, Chen YJ, Lam TI, Taylor KC, Walton JH, et al. (2013) Intravenous HOE-642 reduces brain edema and Na uptake in the rat permanent middle cerebral artery occlusion model of stroke: evidence for participation of the blood-brain barrier Na/H exchanger. *J Cereb Blood Flow Metab* 33: 225-234.
134. Kozniwska E, Romaniuk K (2008) Vasopressin in vascular regulation and water homeostasis in the brain. *J Physiol Pharmacol* 59: 109-116.
135. Yuen N, Lam TI, Wallace BK, Klug NR, Anderson SE, et al. (2014) Ischemic factor-induced increases in cerebral microvascular endothelial cell Na/H exchange activity and abundance: evidence for involvement of ERK1/2 MAP kinase. *Am J Physiol Cell Physiol* 306: C931-C942.
136. Kleindienst A, Dunbar JG, Glisson R, Marmarou A (2013) The role of vasopressin V1A receptors in cytotoxic brain edema formation following brain injury. *Acta Neurochir (Wien)* 155: 151-164.
137. Alonso G, Gallibert E, Lafont C, Guillon G (2008) Intrahypothalamic angiogenesis induced by osmotic stimuli correlates with local hypoxia: a potential role of confined vasoconstriction induced by dendritic secretion of vasopressin. *Endocrinology* 149: 4279-4288.
138. Armstead WM (2001) Vasopressin-induced protein kinase C-dependent superoxide generation contributes to atp-sensitive potassium channel but not calcium-sensitive potassium channel function impairment after brain injury. *Stroke* 32: 1408-1414.
139. Yokoyama T, Saito T, Ohbuchi T, Hashimoto H, Suzuki H, et al. (2010) TRPV1 gene deficiency attenuates miniature EPSC potentiation induced by mannitol and angiotensin II in supraoptic magnocellular neurons. *J Neurosci* 30: 876-884.
140. Krowicki ZK, Kapusta DR (2011) Microinjection of glycine into the hypothalamic paraventricular nucleus produces diuresis, natriuresis, and inhibition of central sympathetic outflow. *J Pharmacol Exp Ther* 337: 247-255.
141. Curras-Collazo MC, Patel UB, Hussein MO (2002) Reduced susceptibility of magnocellular neuroendocrine nuclei of the rat hypothalamus to transient focal ischemia produced by middle cerebral artery occlusion. *Exp Neurol* 178: 268-279.
142. Nishimori K, Young LJ, Guo Q, Wang Z, Insel TR, et al. (1996) Oxytocin is required for nursing but is not essential for parturition or reproductive behavior. *Proc Natl Acad Sci U S A* 93: 11699-11704.
143. Ip S, Chung M, Raman G, Chew P, Magula N, et al. (2007) Breastfeeding and maternal and infant health outcomes in developed countries. *Evid Rep Technol Assess (Full Rep)*, pp: 1-186.
144. Arun E, Yalçın SS, Madendağ Y, Ustünyurt-Eras Z, Kutluk S, et al. (2010) Factors associated with breastfeeding initiation time in a Baby-Friendly Hospital. *Turk J Pediatr* 52: 10-16.
145. Enabudoso EJ, Ezeanochie MC, Olagbuji BN (2011) Perception and attitude of women with previous caesarean section towards repeat caesarean delivery. *J Matern Fetal Neonatal Med* 24: 1212-1214.
146. Wang YF, Hatton GI (2009) Oxytocin, lactation and postpartum depression. *Frontiers in Neuroscience* 3252-3253.
147. Jiang QB, Wakerley JB (1995) Analysis of bursting responses of oxytocin neurones in the rat in late pregnancy, lactation and after weaning. *J Physiol* 486: 237-248.
148. Montagnese CM, Poulain DA, Vincent JD, Theodosis DT (1987) Structural plasticity in the rat supraoptic nucleus during gestation, post-partum lactation and suckling-induced pseudogestation and lactation. *J Endocrinol* 115: 97-105.
149. Abou-Saleh MT, Ghubash R, Karim L, Krymski M, Bhai I (1998) Hormonal aspects of postpartum depression. *Psychoneuroendocrinology* 23: 465-475.
150. Tops M, van Peer JM, Korf J (2007) Individual differences in emotional expressivity predict oxytocin responses to cortisol administration: relevance to breast cancer? *Biol Psychol* 75: 119-123.
151. Cox EQ, Stuebe A, Pearson B, Grewen K, Rubinow D, et al. (2015) Oxytocin and HPA stress axis reactivity in postpartum women. *Psychoneuroendocrinology* 55: 164-172.
152. Wagner KU, Young WS, Liu X, Ginns EI, Li M, et al. (1997) Oxytocin and milk removal are required for post-partum mammary-gland development. *Genes Funct* 1: 233-244.
153. Schagen FH, Knigge U, Kjaer A, Larsen PJ, Warberg J (1996) Involvement of histamine in suckling-induced release of oxytocin, prolactin and adrenocorticotropin in lactating rats. *Neuroendocrinology* 63: 550-558.
154. Stuebe AM, Willett WC, Xue F, Michels KB (2009) Lactation and incidence of premenopausal breast cancer: a longitudinal study. *Arch Intern Med* 169: 1364-1371.
155. Liu X, Jia S, Zhang Y, Wang YF (2016) Pulsatile but not Tonic Secretion of Oxytocin Plays the Role of Anti-Precancerous Lesions of the Mammary Glands in Rat Dams Separated from the Pups during Lactation. *M J Neuro* 1: 002.
156. Yoshida M, Takayanagi Y, Inoue K, Kimura T, Young LJ, et al. (2009) Evidence that oxytocin exerts anxiolytic effects via oxytocin receptor expressed in serotonergic neurons in mice. *J Neurosci* 29: 2259-2271.