

Opinion Article

New Emerging Human Parasites in 21th Century

Cheepsattayakorn A 1,2*

¹10th Zonal Tuberculosis and Chest Disease Center, Chiang Mai, Thailand

²10th Office of Disease Prevention and Control, Department of Disease Control, Ministry of Public Health, Chiang Mai, Thailand

*Corresponding author: Cheepsattayakorn A, 143 Sridornchai Road, Changklan Muang Chiang Mai, 50100 Thailand, Tel: 66-53-140767; E-mail: attaponche@yahoo.com

Rec date: October 2, 2014, Acc date: October 6, 2014, Pub date: October 16, 2014

Citation: Cheepsattayakorn A (2014) New Emerging Human Parasites in 21th Century. Poult Fish Wildl Sci 2: 118. doi: 10.4172/2375-446X.1000118

Copyright: © 2014 Cheepsattayakorn A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Introduction

A number of new human parasites have urged the interest of scientific community in addition to many zoonotic potential parasites. Potential human parasites transmitted from wildlife domestic hosts are included *Dirofilaria ursi*, *Dirofilaria subdermata*, *Dipetalonema* species, *Loa Loa*, *Thelazia callipaeda*, *Babesia duncani*, *Babesia venatorum*, *Babesia*-liked organisms, *Babesia divergens*-liked organism, *Leishmania braziliensis*, *Leishmania major*, *Plasmodium knowlesi*, *Plasmodium simium*, *Onchocerca gutturosa*, *Onchocerca cervicalis*, *Onchocerca jakutensis*, *Onchocerca dewittei japonica*, *Onchocerca lupi*, *Diphyllobothrium dendriticum*, etc. Macaques (*Macaca fascicularis*) are believed to be the primary hosts of *Plasmodium knowlesi* was identified by PCR in 58% of human malaria cases in the Kapit division of Sarawak, Malaysia, between 2000 and November 2002. It is transmitted by the forest dwelling *Anopheles latens*, a mosquito species that feeds on both the monkey natural host as well as on humans. A single human malaria case of *Plasmodium knowlesi*, confirmed by PCR, has been reported from Thailand. Outbreaks of Chagas disease transmitted by accidental ingestion of *Trypanosoma cruzi*-contaminated guava, sugar cane or acai' juices have been reported in South America.