

New Therapeutic Approaches for Rare Inborn Enzyme Deficiencies Using Targeted Gene Delivery Systems

Arjun Mehta*

Department of Medical Genetics, Institute of Human Molecular Studies, Hyderabad, India

DESCRIPTION

Inherited enzyme deficiencies affecting metabolic processing pathways represent a diverse group of rare conditions that often present early in life with multisystem involvement. These conditions arise due to alterations in genes responsible for encoding specific proteins required for cellular metabolism. The resulting biochemical imbalance can affect growth, neurological development, hepatic function, and energy utilization. Conventional care has largely relied on dietary control and enzyme replacement strategies, yet these approaches often provide partial correction and require lifelong administration. The exploration of gene-based interventions has opened new possibilities for long-term management by directly addressing the molecular source of dysfunction.

Recent investigations have focused on vector-mediated gene delivery systems designed to introduce functional copies of defective genes into target tissues. Viral vectors derived from adeno-associated platforms have demonstrated relatively stable expression in non-dividing cells such as hepatocytes and neurons. These systems are selected for their reduced immunogenic profile and ability to maintain episomal persistence, which limits integration-related genomic disturbances. Research groups have optimized tissue specificity by modifying capsid proteins, allowing preferential uptake by organs most affected in metabolic disorders.

One example includes disorders associated with urea cycle dysfunction, where accumulation of ammonia leads to neurological impairment. Experimental models have demonstrated that hepatic-directed gene transfer can partially restore enzymatic activity required for nitrogen processing. In preclinical studies using murine systems, treated groups exhibited improved survival rates and normalized biochemical markers compared to untreated controls. While complete restoration is not consistently achieved, even partial enzyme activity has shown measurable improvement in systemic stability.

Another area of investigation involves lysosomal storage conditions characterized by progressive accumulation of undegraded substrates

within cellular compartments. Gene delivery strategies targeting hematopoietic stem cells have been explored to provide continuous enzyme supply through differentiated progeny cells. This approach allows systemic distribution of corrective proteins, potentially affecting multiple organ systems simultaneously. Long-term follow-up in animal models has indicated sustained enzyme presence and reduced substrate accumulation in neural and visceral tissues. Regulatory considerations remain central to clinical translation. The introduction of foreign genetic material into human cells raises concerns regarding immune recognition and long-term genomic behavior. To address these issues, researchers have implemented transient immunosuppression protocols during vector administration. Additionally, self-complementary vector constructs have been developed to increase transcription efficiency while reducing required dosage levels. These modifications aim to balance therapeutic effect with biological safety.

Another significant area of interest involves messenger Ribonucleic Acid (RNA)-based therapeutic delivery. Unlike Deoxyribonucleic Acid (DNA)-based vectors, mRNA does not require nuclear entry and is translated directly in the cytoplasm. This reduces the possibility of integration into host chromosomes. However, the transient nature of mRNA expression necessitates repeated administration. Chemical modification of nucleotide structures has improved stability and reduced innate immune activation, allowing more consistent protein production in experimental systems. Clinical trial data from early-phase studies have provided encouraging biochemical outcomes in selected metabolic conditions. Patients receiving gene transfer therapy have demonstrated reductions in toxic metabolite accumulation and partial correction of enzymatic function. Despite these improvements, variability in response remains a challenge. Factors influencing therapeutic variability include pre-existing antibodies, vector distribution efficiency, and differences in disease severity at the time of intervention.

Pediatric applications require additional caution due to developmental sensitivity. Early intervention may prevent irreversible organ damage, yet the long-term consequences of

Correspondence to: Arjun Mehta, Department of Medical Genetics, Institute of Human Molecular Studies, Hyderabad, India, E-mail: arjun.mehta.ihms@genmail.edu

Received: 01-Jun-2026, Manuscript No. JSGST-26-42898; **Editor assigned:** 03-Jun-2026, PreQC No. JSGST-26-42898 (PQ); **Reviewed:** 17-Jun-2026, QC No. JSGST-26-42898; **Revised:** 24-Jun-2026, Manuscript No. JSGST-26-42898 (R); **Published:** 01-Jul-2026, DOI: 10.35248/2157-7412.26.16.500

Citation: Mehta A (2026). New Therapeutic Approaches for Rare Inborn Enzyme Deficiencies Using Targeted Gene Delivery Systems. J Genet Syndr Gene Ther. 14:500.

Copyright: © 2026 Mehta A. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

altering gene expression in developing tissues remain under observation. Longitudinal studies are being conducted to evaluate growth patterns, neurocognitive development, and endocrine function following treatment exposure. Ethical frameworks emphasize informed consent, especially in cases where parental authorization is required for neonatal participation. Manufacturing processes for gene-based medicinal products involve strict quality control procedures to ensure purity, potency, and consistency. Production systems utilize cultured cell lines engineered to produce viral particles under controlled conditions. Downstream purification techniques remove residual proteins and nucleic acid contaminants. Each batch undergoes analytical validation before clinical use, including genomic sequencing verification and infectivity assays.

One ongoing challenge involves immune memory formation against vector components, which can reduce efficacy upon repeat administration. Strategies to mitigate this include capsid engineering and use of serotype switching in sequential treatments. Research is also exploring exosome-based delivery systems as an alternative to viral carriers. These naturally occurring vesicles offer biocompatibility advantages, although scaling production remains a technical limitation. Economic considerations also influence accessibility of gene-based

interventions. High production costs and specialized infrastructure requirements limit availability in resource-constrained regions. Collaborative initiatives between academic institutions and biotechnology manufacturers are being explored to reduce cost barriers and expand clinical reach. Policy discussions emphasize equitable distribution without compromising safety standards.

CONCLUSION

Future investigative directions include multiplex gene delivery systems capable of addressing conditions involving multiple gene defects simultaneously. Research is also focusing on regulatory switches that allow controlled activation or suppression of introduced genes based on physiological cues. Such systems aim to provide adaptive therapeutic responses aligned with patient needs over time. Continued refinement of delivery systems, coupled with improved understanding of metabolic pathways, is expected to expand therapeutic options for individuals affected by inherited enzyme deficiencies. While limitations persist, the integration of molecular engineering, clinical research, and computational modeling supports steady advancement in this area of genetic medicine.