

Nutritional Composition of Quail Meatballs and Quail Pickled Eggs

Bayomy HM¹, Rozan MA^{1*} and Mohammed GM²

¹Department of Food Science and Technology, Damanhour University, Egypt

²Department of Nutrition and food Science, Tabuk University, KSA

*Corresponding author: Rozan MA, Department of Food Science and Technology, Damanhour University, Egypt, Tel: +20 45 3368069; E-mail: Mahmoud.abdelgalil@agr.dmu.edu.eg

Received date: March 17, 2017; Accepted date: March 21, 2017; Published date: March 28, 2017

Copyright: © 2017 Bayomy HM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

The aims of this study were to evaluate the acceptance of meatballs and pickled quail eggs, determine the nutritional composition of the quail pickled eggs and quail meatballs. Data indicate that quail meatballs and pickled quail eggs are an acceptable market product. The quail meatballs have 70.56% moisture content and 29.44% dry matter which include 7.78%, 12.98%, 7.10% and 2.48% in wet basis for total lipids, proteins, ash and carbohydrate, respectively. Total cholesterol content in quail meatballs was 0.087 g/100g. Regarding pickled quail eggs, the values of moisture, dry matter, total lipids, proteins, ash and carbohydrate are 53.32%, 46.68%, 20.09%, 20.38%, 4.70% and 1.51% in wet basis, respectively. Average of cholesterol content in pickled quail eggs is 2.06 g/100 g yolk. Total essential amino acids were 6.17 g/100 g protein in quail meatballs, while total non-essential amino acids were 5.87g/100g protein in quail meatballs. In pickled quail eggs the values of total essential amino acids and non-essential amino acids were 6.84 g/100g and 8.81 g/100g protein, respectively. Total saturated fatty acids (SFAs) in quail meatballs and quail pickled eggs are 26.37% and 37.52%, respectively. Total mono-unsaturated fatty acids and poly-unsaturated fatty acids contents in quail meatballs are 36.61% and 36.21, respectively, while these values for pickled quail eggs are 54.12% and 8.36%, respectively.

Keywords: Quail; Meatballs; Pickled egg; Nutritional composition

Introduction

Poultry meat and their products are consumed in large quantities and the consumption is growing all over the world [1]. Quail products have gained popularity in the last few years by consumers. Quail is belonging to the species *Coturnix coturnix japonica*. It was first domesticated in Japan as a source of both egg and meat after that quail distribution in the wild spreads over large areas of Asia, Europe and Africa, [2]. Quail meat is recommended for the low-fat diet because it contains low amount of fat and cholesterol especially thanks to its thin skin and low fat accumulation between its tissues [3]. Japanese quail meat may be taken into consideration in diet for prevention of heart diseases because of high C18:1 content. The ratio between PUFAs and SFAs was 0.43 in female quail meat and 0.40 in male quail meat. Both ratios are in accordance with the WHO recommendations. Quail eggs have a very interesting chemical composition, being rich in fat soluble vitamins and B-complex; in proteins, amino acids, macro and microelements, but are low in cholesterol [4], triglycerids and saturated fatty acids [5]. Quail eggs used to help treat tuberculosis, asthma, and even diabetes by Chinese. Also quail eggs can help prevent sufferer of liver, kidney or gallbladder stones, and remove these types of stones. Quail eggs have higher nutritional value than those offered by other eggs with they are rich sources of antioxidants, minerals, and vitamins, and give us a lot of nutrition than do other foods [6]. Because of its valuable chemical compositions, quail egg is considered to be a true universal panacea, placed 3rd in the Chinese natural medicine after snake venom and Ginseng [7,8]. The eggs of quail are used in the same manner as those of chickens. Because of their small size and attractive appearance quail eggs are often used for special purposes. Pickled quail eggs are a specialty. Pickles occupy an important place

among processed foods. With the increasing popularity of quail (*Coturnix coturnix japonica*), there is a good scope for the utilization of small and tinted quail eggs in the form of pickle. Technological details pertaining to the manufacture of pickled quail eggs have been worked out with a view to offer such nutritious, ready-to-eat product to the consumers.

Quail meat is considered a delicacy. It can be fried or roasted, and many recipes have been developed. Meatballs are a popular meat product in Egypt. This includes products such as chicken meatballs, fish meatballs, beef meatballs; quail meat is also potential raw materials for meatballs production. According to [9] quail meat can be used for the manufacture of meatballs successfully by using different types of flour as an alternative to the use of other meats such as beef and chicken.

Egyptian consumer preferably eats chicken eggs. But quail eggs sold in the form much more restricted, commonly used in the ornamentation of cold dishes or snacks. In recent years, it has become spread the quail eggs consumed by the belief that these do not contain cholesterol or its content is irrelevant and greater value nutritious, belief that has been fueled to some extent by the producers themselves at the lack of information about their composition.

In developed countries continues to drive consumption chicken egg through the development of new egg products are used in various dishes, however quail egg by their different physical characteristics to chicken cannot be used for preparing the same egg products but can be exploited in other presentations. The objective of this work was to evaluate the acceptance of meatballs and pickled quail eggs. Determine the nutritional composition of the quail pickled eggs and quail meatballs.

Materials and Methods

Materials

Quail (*Coturnix coturnix japonica*) carcasses and eggs were collected from the National Center for Agricultural Research in Dokki, Egypt. Other materials were collected from local markets in Damanhour city, Egypt.

Methods

Preparation of Samples

Preparation of pickled quail eggs

According to Indian standard [10] fresh quail eggs stored at ambient temperature for at least 24 hours, for about 10 min at simmering temperature. After boiling, the eggs shall be cooled immediately in running tap water and peeled off manually. There shall be no pieces of egg shell or shell membrane adhering over the peeled eggs. Peeled eggs showing torn or rugged-looking albumen surface shall not be used for pickling. Pickling solution shall consist of vinegar and water (50: 50 v/v), 8% (w/v) common salt, and 2% (w/v) each spice mixture, onion (fresh and chopped), garlic and ginger. The solution shall be boiled for 10 min and then filtered using clean muslin cloth. Pickling solution shall be heated to $70 \pm 2^\circ\text{C}$ and poured onto the peeled eggs. Cover the peeled eggs with pickling solution. The pickled eggs shall be aged for 48 hours at ambient temperature in pre-sterilized glass container.

Preparation of meatballs

The quail carcasses were deboned mechanically. The mean carcass weights were 203.04 ± 16.67 g. meatballs blend was prepared using the formula given by [11]. as follows: Meat 90% was mixed into house mincer, flour 3%, fresh onion 3% were peeled and minced before the addition, 2% salt and 2% spices. These ingredients were mixed together, divided into balls to formulate quail meatballs. Samples were freeze stored at $-18 \pm 2^\circ\text{C}$. Meatballs cooked before analysis at 95°C for 5 min.

Sensory properties

The sensory properties of consumers were applied for the pickled quail eggs and meatballs. Sixty panelists evaluated the sensory attributes of the quail products. Panelists were given no time limit for the evaluation. Pickled quail eggs and meatballs were cut into small pieces and placed on a plastic plate. The attributes evaluated were color, flavor, texture and overall acceptability. For each sample, panelists scored their liking of these characteristics using a five-point hedonic scale (1 = dislike very much, 2 = dislike slightly, 3 = neither like nor dislike, 4 = like slightly, 5 = like very much) [12].

Analytical Methods

Chemical analysis

Moisture, protein, fat and ash contents were determined in accordance with standard AOAC methods of [13]. Protein determination involved a Kjeldahl assay (Nx6.25). Fat was determined extracting samples in a Folch apparatus using chloroform: Ethanol (2:1 v/v) as a solvent. Moisture was quantified by oven-drying 10 g samples at 100°C overnight. Ash was determined after incineration in a furnace

at 500°C and carbohydrate content was calculated by computing the difference. Minerals (Ca; Na; P & Fe) contents were performed by the classical method of chemical analysis [13].

Total cholesterol content

It was determined by the method of Lieberman-Buchart [14].

Amino acid profile

The technique recommended by [15] for fractionation of amino acids was applied by Gas Liquid Chromatography (GLC).

Fatty acid methyl esters analyses

For fat content determination lipids were extracted from 10 g breast by the method described by [16,17]. Five mL chloroform/methanol (2:1 v/v) solution was added to sample and shook thoroughly by vortex for 3 min. One mL of 0.9% NaCl was added to mix and shake again. The chloroform phase containing lipids were collected. Lipid extracts were converted to fatty acid methyl esters as described by [17]. It was prepared after alkaline hydrolysis following procedures. Briefly, lipid extract was mixed with 0.5N 2mL methanolic NaOH and held at 100°C for 10 min. This solution was cooled to room temperature and mixed with 2mL of (14% Boron tri fluoride, 86% Methanol). Then after, solution obtained by adding BF_3 was incubated at 100°C for 2 min and cooled down to again room temperature. 1mL n-heptane was added to final solution and thoroughly mixed by vortex for 3min and reheated to 100°C for 1min. Final solution was cooled down to room temperature and centrifuged at 300 rpm for 5 min. Upper layer (heptane phase) was transferred to glass tube for the GC analysis. The final concentration of the Fatty acid methyl esters analyses was approximately 7mg/mL in heptane.

Statistical analysis

The obtained data were statistically analyzed by routine statistical methods by means \pm standard deviation of Microsoft Excel software.

Results and Discussion

Sensory properties

Sensory properties was conducted to measure the organoleptic properties of organic, which include color, taste, smell, texture and general acceptance of both meatballs quail and pickled quail eggs. General precautions to taste tests have been taken. Evaluation of sensory properties of quail meatballs and pickled eggs are shown in Table (1). In general, the assessment of panelists indicates moderate likeability, may be due to the fact that the panelists were not very familiar with meatballs made from quail meat. Regarding to pickled product, the data indicate that pickled quail eggs are an acceptable market product.

Data of chemical composition for quail meatballs and pickled quail eggs are presented in Table 2. Results demonstrated that quail meatballs have 70.56% moisture content and 29.44% dry matter which include 7.78%, 12.98%, 7.10% and 2.48% in wet basis for total lipids, proteins, ash and carbohydrate, respectively. These results are agreed with results produced by [9]. They found that quail meatballs have value of 65.94% moisture, 13.53(% wet basis) protein, 10.59 (% wet basis) fat, 2.30(% wet basis) ash and 7.63(% wet basis) carbohydrates. According to [18] the proximate analysis values of quail meatballs are

not so different with those of meatballs made from chicken which prepared by [18]. They found that chicken meatballs have 64.33-71.81%, 9.94-15.96 (% wet basis), 4.26-14.00(% wet basis), 1.92-2.82(% wet basis) and 5.54-20.85(% wet basis) for moisture, protein, fat, ash and carbohydrate content, respectively.

Properties	Meatball Quail	Pickled quail egg
Colour	4.7 ± 0.23	4.2 ± 0.51
Taste	4.5 ± 0.35	4.5 ± 0.35
Odour	5.0 ± 0.37	4.0 ± 0.65
Flavour	4.3 ± 0.27	4.3 ± 0.45
Texture	4.3 ± 0.17	4.1 ± 0.65
Overall acceptability	4.5 ± 0.24	4.2 ± 0.45
The value ± standard deviation (SD)		

Table 1: Sensory properties of quail meatballs and pickled quail egg.

Chemical composition

Chemical composition (g/100 g DW)	Quail meatballs	Pickled quail eggs
Moisture	70.56 ± 2.04	53.32 ± 1.74
Dry matter	29.44 ± 1.63	46.68 ± 0.86
Total lipids	7.78 ± 0.68	20.09 ± 0.85
Cholesterol	0.087 ± 0.005	2.06 ± 0.56 g/100 g yolk
Proteins	12.98 ± 1.03	20.38 ± 2.03
Carbohydrate	6.10 ± 0.74	1.51 ± 0.04
Ash	2.48 ± 0.22	4.70 ± 0.33
Na (mg/kg)	1425.23 ± 21.67	1058.96 ± 7.97
P (mg/kg)	3838.79 ± 18.31	955.19 ± 6.55
Ca (mg/kg)	1448.60 ± 8.08	613.21 ± 1.27
Fe (mg/kg)	36.028 ± 0.17	22.476 ± 0.68
All values are means of triplicate determinations ± standard deviation (SD). DW: Dry Weight; TC: Total Carbohydrates		

Table 2: Chemical composition of quail meatballs and pickled quail egg (g/100 g in wet basis).

According to the results of [19], quail meatballs in Malaysian markets have higher protein content and lower carbohydrate content than various kinds of commercial chicken meatballs, who reported protein and carbohydrate contents ranging from 12.83-13.71% and 5.23-8.25%, respectively. Total cholesterol content in quail meatballs was 0.087 g/100g. This value is agreed with that reported by [20] who found that cholesterol content in quail carcass was 0.097 and 0.094 g/100g for male and female, respectively. It is clear that quail meatballs have high minerals content this may be due to the mixing of soft bone with flesh of quail when manufacturing.

With regard to pickled quail eggs, Table 2 indicated that the values of moisture, dry matter, total lipids, proteins, ash and carbohydrate are 53.32%, 46.68%, 20.09%, 20.38%, 4.70% and 1.51% in wet basis, respectively. Average of cholesterol content in pickled quail eggs is 2.06 g/100 g yolk. [21] found that the proximal composition of quail eggs with a protein content of 13.6 ± 2.1%, total lipids of 12.59 ± 2.2 and 1.13 ± 0.33% cholesterol in wet basis. [22] reported that the quail eggs have 1.06, 4.01, 9.89, 12.7, and 72.25 g/100g for ash, carbohydrate, fat, protein and moisture, respectively. Tunsaringkarn et al. indicated that the quail egg yolk has a 1.21 ± 0.07 g of cholesterol per 100 g of yolk [23]. According to Bragagnalo et al. [24] the highest cholesterol content was observed in ostrich and duck eggs, whereas the lowest in [23] compared the cholesterol content of Brazilian chicken and quail eggs. They found that the cholesterol level between chicken and quail eggs in terms of yolk were not significant difference. Opposite results were presented by Kazmierska et al. [25], who reported higher cholesterol content in quail egg yolk (13.6 mg/g) compared to Tunsaringkarn et al. [23].

Amino acids content

Amino acid composition is one of the most important factors to determine the quality of food protein. In general, high protein food is also high in contents of amino acids including eight which are essential in human nutrition: lysine, valine, leucine, isoleucine, threonine, tryptophan, methionine and phenylalanine [26]. Analysis of amino acids (g/100g sample) in quail meatballs and pickled quail eggs are recorded in Table 3.

Essential amino acids were 6.17 g/100 g sample this value equals 47.5% from protein in quail meatballs. Among the essential amino acids lysine has the greatest value (1.63%) of amino acids. lysine is assists in calcium absorption, required for growth and bone development in children, and assists in maintaining the correct nitrogen balance in the body, as well as maintaining lean body mass. Lysine is also needed to produce antibodies, enzymes, collagen formation, hormones, as well as repair of tissue [27]. While total non-essential amino acids was 5.87g/100g sample in quail meatballs. In pickled quail eggs, total essential amino acids and total non-essential amino acids were 6.84 g/100g sample and 8.81 g/100g sample, respectively. The greatest value of amino acids refers to leucine as essential amino acid. Leucine is a branched chain amino acid along with valine and isoleucine. It is beneficial and functional to protein structure for 60-70% in human body, and blood sugar level regulation which maintains a balance of insulin and glucose [28]. It proposed as a promising pharmaconutrient in the prevention and treatment of sarcopenia and/or type 2 diabetes [29]. The “ideal” protein should contain the following concentrations of essential amino acids: lysine – 5.5%, leucine – 7%, isoleucine – 4%, valine – 5%, methionine + cysteine – 3.5%, threonine – 4%, phenylalanine + tyrosine – 6% as reported in FAO/WHO specialists in 1973 [30].

Fatty acids composition

Quail is the good source of both saturated and unsaturated fatty acids. In our studies, we found that quail meatballs and pickled quail eggs used in study contain saturated and unsaturated fatty acids. As shown in Table (4), Palmitic acid (C14:0) is present in higher number in saturated fatty acids with values of up to 20.48% and 28.11% for quail meatballs and pickled quail eggs, respectively. It is similar with results reported by [31]. They found that Palmitic acid (C16:0) was the major fatty acid, followed by stearic acid (C18:0) in female and male

quail meats. Myristic acid was in very low concentration in comparison to palmitic and stearic acid. Sum of saturated fatty acids (Σ SFA) in quail meatballs and quail pickled eggs are 26.37% and 37.52%, respectively. Saturated fatty acids play important role in different biological functions like source of metabolic energy and in cell membrane structure formation. Whereas Oleic (C18:1) is present in highest value in all fatty acids. Sum of mono unsaturated fatty acids (Σ MUFA) and Sum of poly unsaturated fatty acids (Σ PUFA) in quail meatballs are 36.61% and 36.21, respectively. While Σ MUFA and Σ PUFA values for pickled quail eggs are 54.12% and 8.36%, respectively. These data are similar with that reported by [32]. He found that Σ MUFA and Σ PUFA values for egg in Pharaoh Japanese quail are 51.3 % and 9.9%, respectively. Generally, bird's meat is relatively rich in the essential Ω -3 α -linolenic fatty acid (C18:3). Our results for α -linolenic acid content in quail meatballs and pickled quail eggs are 23.6% and 5.36%, respectively and higher than to those of [33] who reported that content of α -linolenic acid was 1.74% in chicken meat. The α -linolenic acid content in the meat of the different duck populations varies in a wider range – 0.8-1.62% [34].

Amino acids	Quail meatballs	Pickled quail eggs
Essential amino acids		
Lysine	1.63 ± 0.41	1.36 ± 0.16
	1.63 ± 0.41	1.36 ± 0.16
Methionine	0.48 ± 0.08	0.78 ± 0.03
Leucine	0.89 ± 0.17	1.47 ± 0.09
Isoleucine	0.97 ± 0.03	0.77 ± 0.02
Phenylalanine	0.94 ± 0.08	0.81 ± 0.08
Threonine	0.47 ± 0.24	0.71 ± 0.03
Valine	0.79 ± 0.16	0.94 ± 0.12
Σ essential amino acids	6.17	6.84
Non-essential amino acids		
Cysteine	0.12 ± 0.01	0.83 ± 0.06
Tyrosine	0.43 ± 0.06	0.51 ± 0.04
Histidine	0.62 ± 0.11	0.47 ± 0.02
Arginine	0.51 ± 0.09	0.47 ± 0.04
Glycine	0.64 ± 0.15	0.51 ± 0.07
Glutamic acids	1.02 ± 0.28	2.02 ± 0.04
Serine	0.38 ± 0.06	1.41 ± 0.07
Alanine	0.59 ± 0.13	0.73 ± 0.05
Proline	0.51 ± 0.03	0.60 ± 0.07
Asparatic acids	1.05 ± 0.06	1.26 ± 0.03
Σ Non-essential amino acids	5.87	8.81

All values are means of triplicate determinations ± standard deviation (SD)

Table 3: Amino acids content in quail meatballs and pickled quail eggs %.

Sample Fatty acid	Quail meatballs	Pickled quail eggs
Myristic (C14:0)	0.95 ± 0.06	0.79 ± 0.07
Myristoleic (C14:1)	1.15 ± 0.06	6.29 ± 0.51
Palmitic (C16:0)	20.42 ± 0.34	28.11 ± 0.62
Palmitoleic (C16:1)	0.46 ± 0.03	2.60 ± 0.04
Stearic (C18:0)	5.00 ± 0.08	8.62 ± 0.11
Oleic (C18:1)	35.80 ± 0.74	45.23 ± 0.65
Linoleic (C18:2)	23.62 ± 0.05	5.36 ± 0.04
γ -Linolenic (C18:3)	3.16 ± 0.02	2.27 ± 0.13
Eicosapentaenoic acid (C20:5)	9.43 ± 0.28	0.73 ± 0.07
Σ SFA	26.37	37.52
Σ MUFA	36.61	54.12
Σ PUFA	36.21	8.36

All values are means of triplicate determinations ± standard deviation (SD)

Table 4: Fatty acid content (%) of quail meatballs and pickled quail eggs (%) from total lipids.

Wolaszyn et al. [35] reported that Quail eggs have lower content of polyunsaturated fatty acid (25.1%) than chicken eggs (31.3%) and higher concentration of monounsaturated fatty acid in yolk (45%) compared to chicken eggs (39.1%). The total concentration of saturated fatty acid in yolk was similar in both species.

Conclusion

Quail meatballs and pickled quail eggs are an acceptable market product, have a very interesting chemical composition. Quail meatballs have high concentration of protein content, lower carbohydrate content and high minerals content. Sum of essential amino acids in quail meatballs was 6.17 g/100 g sample this value equals 47.5% from protein, while in pickled quail eggs total essential amino acids and was 6.84 g/100g sample which represents 33.56% from protein content in pickled egg. Sum of saturated fatty acids (Σ SFA) in quail meatballs and quail pickled eggs are 26.37% and 37.52%, respectively. So we should transfer knowledge to people for good nutrient benefits of quail eggs as good nutritional foods and may be the alternative resolving problem of people in some or all nutritional nutrients necessary for human health in developing countries.

Acknowledgements

The authors gratefully acknowledge National Center for Agricultural Research in Dokki, Egypt for providing necessary facilities for the successful completion of this research work.

References

- Mielnik MB, Aaby K, Rolfsen K, Ellekjær MR, Nilsson A (2002) Quality of comminuted sausages formulated from mechanically deboned poultry meat. *Meat Sci* 61: 73-84.
- Mizutani M (2003) The Japanese Quail. Laboratory Animal Research Station, Nippon Institute for Biological Science, Kobuchizawa, Yamanashi, Japan.
- Alarслан OF (2006) Modern quail manufacturing and basic nutrition principles.
- Teuşan A, Prelipcean A, Teuşan V (2011) Research regarding the structure, chemical composition and calorificity of quail eggs (*Coturnix coturnix japonica*), deposited at the beginning phase of lying.
- Vacaru-Opriş I (2002) Poultry treaty. Publisher Ceres: Bucharest.
- Lalwani P (2011) Quail Egg Nutrition.
- Polen T, Erman V (2007) Tips about raising quail, Rev. Collection Farm Publishing World press: Timisoara.
- Teuşan (Prelipcean) Anca (2010) Contributions to the knowledge qualities hatching quail eggs (*Coturnix coturnix japonica*) and embryonic development in this species.
- Ikhlas B, Huda N, Noryati I (2011) Chemical composition and physicochemical properties of meatballs prepared from mechanically deboned quail meat using various types of flour. *Int J Poult Sci* 10: 30-37.
- Indian Standard (1988) Poultry Products-Pickled Quail Eggs.
- Youssef MH (1972) Meat Cooling, Storage, Preservation and Industry. Agriculture Ministry, General Management for Veterinary Laboratories and Researches.
- Hoang DN (2008) Sensory evaluation of food. Ho Chi Minh City University of technology Press. Ho Chi Minh City (in Vietnamese).
- AOAC (2002) Official Methods of Analysis of the Association of Official Analytical Chemists (14th Rew edn.). Washington, DC, USA.
- Carmo BM (2005) Lipid Extraction and Cholesterol Quantification: A Simple Protocol. *Journal Chemical Education* 82: 103-104.
- Mabbott AG (1990) Qualitative amino acid analysis of small peptide by GC/MS. *J Chem Educ* 67: 441-445.
- Folch JM, Lees M, Stanley GHS (1957) A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 226: 497-509.
- AOAC (1995) Official Methods of Analysis of the Association of Official Analytical Chemists (16th edn.). Washington, DC, USA.
- Huda N, Yap HS, Yong LH (2009) Proximate composition, color, textural profile of Malaysian chicken balls. *Pak J Nutr* 8: 1555-1558.
- Huda N, Shen YH, Huey YL, Ahmad R, Mardiah A (2010) Evaluation of physico-chemical properties of Malaysian commercial beef meatballs. *Am J Food Tech* 5: 13-21.
- Genchev A, Mihaylova G, Ribarski S, Pavlov A, Kabakchiev M (2008) Meat quality and composition in Japanese quails. *Trakia J Sci* 6: 72-82.
- Fernando GS, Aideé HU (2011) Sensory evaluation of pickled quail eggs and nutritional composition. *Veterinary Journal Magazine*.
- Tunsaringkarn T, Tungjaroenchai W, Siri Wong W (2012) Nutrient benefits of quail (*Coturnix Coturnix Japonica*) Eggs. *Ann Food Sci Technol* 13: 122-131.
- Bragagnalo N, Rodriguez-Amaya DB (2003) Comparison of the cholesterol content of Brazilian chicken and quail eggs. *J Food Comp Anal* 16: 147-153.
- Kaźmierska M, Jarosz B, Korzeniowska M, Trziszka T, Dobrzański Z (2005) Comparative analysis of fatty acid profile and cholesterol content of egg yolks of different bird species. *Pol J Food Nutr Sci* 55: 69-73.
- Sinanoglou VJ, Strati IF, Miniadis-Meimaroglou S (2011) Lipid, fatty acid and carotenoid content of edible egg yolks from avian species: A comparative study. *Food Chem* 124: 971-977.
- Kim BH, Lee HS, Jang YA, Lee JY, Cho YJ, et al. (2009) Development of amino acid composition database for Korean foods. *J Food Comp Anal* 22: 44-45.
- Vitalhealthzone (2007) Lysine amino acid.
- Khan S (2012) Leucine Structure.
- Van Loon LJ (2012) Leucine as a pharmaconutrient in health and disease. *15: 71-77*.
- Ribarova F, Shishkov S, Baklova I (1987) Amino acid content of the Bulgarian foodstuffs.
- Gecgel U, Yilmaz I, Gurcan EK, Karasu S, Dulger GC (2015) Comparison of fatty acid composition between female and male japanese quail meats. *J Chem* 2015: 1-8
- Genchev A (2012) Quality and composition of Japanese quail eggs (*Coturnix japonica*). *10: 91-101*.
- Rondia P, Delmotte C, Maene D, Blecker C, Toussaint JF, et al. (2003) Effect of the inclusion time of extruded linseed supplementation before slaughter on n-3 fatty acids enrichment of chicken meat. *Proceeding of the XVIth European Symposium on the Quality of Poultry Meat, Saint-Brieuc, France*.
- Wolaszyn J, Ksiazkiewicz J, Orkusz A, Skrabka-Blotnicka T, Biernat J, et al. (2003) Fatty acid profile of lipids from duck muscles of three polish conservative flocks. *Proceeding of the XVIth European Symposium on the Quality of Poultry Meat, 23-26 September 2003, Saint-Brieuc, France*.
- Polat ES, Cital OB, Garip M (2013) Fatty acid composition of yolk of nine poultry species kept in their natural environment. *Anim Sci Pap Rep* 31: 363-368.