

# Obstetric and Perinatal Outcomes in Teenage Pregnancy: a Literature Review

Grazzia Rey\*

Department of Medical Education, School of Medicine, Montevideo, Uruguay

## ABSTRACT

Pregnancy during adolescence continues to be a frequent situation, mostly in developing countries. Maternal age is a key element to be taken into account when managing obstetric patients, as it is related with obstetric and perinatal outcomes.

We conducted a literature review to address the obstetric results of pregnancy in teenage mothers. We reviewed articles for the last 5 years in different databases including 'Pregnancy', 'Teenager', 'Adolescent', 'Complications' and 'Outcomes' as key words.

We obtained 2260 articles after the initial search, but only 28 met the inclusion criteria and were therefore reviewed. Most studies were excluded for not clearly specifying one of more groups of teenage patients, lacking comparison of these groups with no-teenage patients, or not addressing the obstetric outcomes of those pregnancies.

The outcomes were divided into complications of pregnancy, childbirth, puerperium and neonatal complications. We found that there is an association between young maternal age and preterm birth, pre-eclampsia/eclampsia, fetal growth restriction and stillbirth, among others. For the newborn we found outcomes such as low APGAR scores and admission to intensive care were more frequent in teenage mothers.

We conclude that maternal age is an important feature, as there are several obstetric and perinatal compications that are more frequent in teenage mothers than in other age groups.

Key words: Pregnancy; Teenager; Adolescent; Complications; Outcomes

### INTRODUCTION

The Maternal age is one of the most important pieces of information to be taken into account in obstetrics and on it depends different risks and prognosis that add up to other clinical features. Under that logic, adolescence is a particularly relevant period involving special features when compared to other moments within the reproductive life. The World Health Organization defines adolescence as a stage of life in which the individual acquires reproductive capacity, transiting the psychological patterns from childhood to adulthood; considering that it runs between 10 and 19 years. It also classifies it in early adolescence (from 10 to 13 years old), middle adolescence (between 14 to 16 years old); and late adolescence (17 to 19 years), each one representing different risks during pregnancy.

Even though there is a decreasing number in teen pregnancies worldwide, it is still particularly relevant in developing countries, accounting for a significant percentage of all pregnancies. According to World Health Organization approximately 11% of all births in the world still occur among girls aged 15 to 19. In turn, some 16 million girls aged 15 to 19 and approximately 1 million girls under 15 give birth each year, mostly in low and middle-income countries. Complications during pregnancy and childbirth are the second leading cause of death among girls aged 15 to 19 worldwide and babies of teenage mothers face a significantly higher risk of complications than those born to women aged 20 to 24 years.

Our objective was to examine whether teenage pregnancy is associated with increased risk of maternal and neonatal complications compared to adults through a systematic literature review to help clinicians in their actions before and during pregnancy in adolescents.

#### MATERIALS AND METHODS

A search was carried out in the electronic databases: PubMed, Medline, Scielo and Lilacs to identify relevant articles. The search was conducted with the following terms: 'Pregnancy, 'Teenager', 'Adolescent', 'Complications' and 'Outcomes'. The search was restricted to articles in english or spanish and published between

Correspondence to: Grazzia Rey, Department of Medical Education, School of Medicine, Montevideo, Uruguay Email: grey@fmed.edu.uy Received: February 07, 2021, Accepted: February 17, 2021, Published: February 24, 2021

Citation: Rey G (2021) Obstetric and Perinatal Outcomes in Teenage Pregnancy: a Literature Review. Gynecol Obstet. 3: 554.

**Copyright:** © 2021 Grazzia R, This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

#### Grazzia R

2014 and 2019. The last search was conducted on december 23, 2019.

The search of all articles found with these terms was carried out and their selection by title and abstract began to determine if they were consistent with the objectives of the review. Search was carried out by two independent reviewers. Articles that were potentially eligible for inclusion were read in their entirety. When the question remained whether the article was eligible or not, it was also read in its entirety. Disagreements and questions on eligibility for inclusion were discussed among all authors. An exhaustive reading of the selected articles was carried out and a synoptic table was completed. The results were presented descriptively.

#### **Inclusion** Criteria

The choice of the studies was made based on the description of: complications during pregnancy (abortion, threatened abortion, preeclampsia eclampsia syndrome, threat of preterm delivery, gestational diabetes, urinary tract infections, anaemia, premature rupture of membranes, chorioamnioitis, intrauterine growth restriction, stillbirth and maternal admission to intensive care unit); complications during childbirth (suspected fetal hypoxia, presentation abnormalities, cephalopelvic disproportion, caesarean section, instrumental vaginal delivery, perineal tear and episiotomy); complications during the puerperium (postpartum infection and postpartum hemorrhage); and neonatal complications (preterm birth, low birth weight, large for gestational age, Apgar below 7 points at the first minute of life and at 5 minutes, admission of the neonate to intensive care unit, neonatal death, congenital malformations and neonatal sepsis).

Articles from the last 5 years (between December 2014 and December 2019) that clearly defined a group of adolescents under 20 years of age who had a comparison group with adult women of 20 years or more were included.

### **Exclusion** Criteria

Author letters, editorials, case reports, epidemiological studies and reviews were excluded.

#### Data Extraction

Data from each study were extracted taking into account: country, year, type of study, method of collection, sample size, comparison groups, inclusion and exclusion criteria, and results (complications during pregnancy, childbirth, puerperium and neonatals mentioned previously).

### RESULTS

In the initial search, a total of 2.260 were obtained. After the analysis according to title and abstract, 2.195 of the articles were excluded because they did not contemplate the objectives. When reading the full text, 33 more articles were excluded because they did not present comparison groups with an adult population, 2 were excluded because they were revisions, 1 was excluded for being duplicate and 1 was excluded for being in Chinese. A total of 28 articles that met the inclusion criteria were obtained.

As for the year of publication, 2 were published in 2014, 7 in 2015, 6 in 2016, 8 in 2017, 5 in 2018 and 0 in 2019. In relation to the country producing the study: 12 were produced by Latin American countries (3 in Colombia, 2 in México, 3 in Perú, 1 in Brazil, 1 in Honduras, 1 in Cuba and 1 in Ecuador, 9 in Asia, 2 in Thailand,

1 in South Korea and 1 in India, 3 in Europe, 1 in Spain, 1 in Greece, 2 in North America (both of them in U.S.A) and 2 in África (both of them in Cameroon.

Of the 28 studies, 21 corresponded to cohort studies and 7 correspond to case control studies. Regarding the method of data collection 23 they did it from medical records, 4 from survey and 1 that did not specify the method of data collection.

The studies had age groups for adolescents and for the control group variables. Of the 28 studies: 1 study compared adolescents aged 14 to 19 with women aged 20 to 34, 4 study compared women under 19 years of age and older than 19 years, 1 study compared adolescents aged 10 to 19 years with ages 20 or older, 1 study compared adolescents aged 10 to 19 divided into early adolescents (10 to 14 years) and late adolescents (15 to 19 years) with women aged 20 to 35, 1 study compared adolescents aged 10 to 19 divided into early adolescents (10-14 years) and late adolescents (15-19 years) with women older than or equal to 20 years, 3 study compared adolescents aged 10 to 19 years with adults aged 20-34 years, 1 study compared women under 20 with women aged 20 to 34, 1 study compared women under or equal to 19 years divided into early adolescents (under 15 years) and late adolescents (15 to 19 years) with women aged 25 to 30 years, 1 study compared adolescents aged 10 to 19 divided into early adolescents (10-16 years) and late adolescents (17-19 years) with women older than or equal to 20 years, 1 study compared women under 18 or equal to women aged 19 to 36, 1 study compared adolescents aged 13 to 19 divided into early (aged 13 to 16) and late (aged 17 to 19) with women aged 20 to 21, 1 compared women aged 15 to 19 with women aged 20 to 49, 1 compares children under 19 with births nationwide, 1 study compared children under 20 with women aged 20 to 34, 1 study compared adolescents aged 15 to 19 with women aged 20 to 34, 1 study compared adolescents aged 13 to 19 years with over 20 years, 1 study compared patients under or equal to 19 years divided into children under 16 to 16 years and 16 to 19 years with women aged 20 to 34 years, 1 study compared patients younger than or equal to 19 years divided into children under 15 to 16 years and 16 to 19 years with women aged 20 to 25, 1 study compared adolescents aged 12 to 19 with women aged 20 to 34, 1 study compared patients younger than or equal to 19 years divided into children under 16 to 16 years and 16 to 19 years with women aged 20 to 30, 1 study compared children under 20 with women aged 20 to 30 (24), 1 study compared adolescents aged 12 to 17 with women aged 18 to 29 and 1 study compared adolescents aged 10 to 19 with women aged 20 to 29.

### **Puerperal Complications**

Regarding postpartum hemorrhage 13 studies show results where 5 show increased risk: 3 statistically significant and 2 not. While 8 studies do not show an increased risk for it: 2 are statistically significant, 5 do not and one does not calculate statistical significance.

On the other hand, 5 studies show data on the risk of puerperal infection where 4 show increased risk: 3 are statistically significant and 1 does not. Only 1 study does not show an increased risk for it, not calculating statistical significance.

### **Neonatal Complications**

Regarding preterm births and teenage pregnancy, 22 studies showed results. Of these, 19 showed an increase in risk for it: 15 are statistically significant and 4 are not. While only 3 studies did

#### Grazzia R

not show an increased risk, none were statistically significant.

## DISCUSSION

The results of our review indicate that there is an association between young maternal age and low birth weight, preterm birth, pre-eclampsia/eclampsia, anaemia, fetal growth restriction, stillbirth and premature rupture of membranes.

As for complications during childbirth there is an increase in the number of perineal tears and episiotomies. However, there is no increased risk of fetal hypoxia, cephalopelvic disproportion, caesarean section, instrumental vaginal delivery, or abnormal fetal presentation.

Regarding puerperal complications, there is an increased risk of puerperal infection but not of postpartum hemorrhage.

As for the newborn, there is a higher risk of low birth weight, Apgar score less than 7 points at 1 and 5 minutes, and admission to intensive care unit.

Regarding this study, it is important to highlight that no published reviews of the complications of teenage pregnancies in Latin America were found in the past 5 years.

#### CONCLUSIONS

The findings of this review indicate that there is an association between young maternal age and low birth weight, preterm birth, pre-eclampsia / eclampsia, anaemia, fetal growth restriction, stillbirth and premature rupture of membranes, increased number of perineal tears and episiotomies, there is a greater risk for the newborn of low birth weight, apgar score less than 7 points at the first and 5 minutes of life and greater risk of admission to intensive care. We can conclude then that adolescent pregnancy in developing continents presents greater risks than pregnancies in adult women.

We may assume this risk is greater when taking into account other aspects such as the socio-economic and cultural impact for adolescents or the weight of parenting, data that do not arise from our research but should be the object of further studies. Finally, we think that future research addressing adolescent pregnancy could help better understand this phenomenon and its implications.

#### REFERENCES

- 1. Dutta DC. Text book of Gynecology. 7th ed. Benign lesions of thecervix. Jaypee Brothers. 2016:217-218.
- Goldacre MJ, Loudon N, Watt B, Grant G, Loudon JD,McPherson K, et al. Epidemiology and clinical significance of cervical erosion in women attending a family planning clinic. BrMed J. 1978:1748-750.
- Critchlow CW, Wölner-Hanssen P, Eschenbach DA, Kiviat NB,Koutsky LA, Steven CE, et al. Holmes Determinants of cervicalectopia and of cervicitis: age, oral contraception, specific cervicalinfection, smoking, and douching. Am J Obstet Gynecol.1995;173:534-43.
- Morrison CS, Bright P, Wong EL, Kwok C, Yacobson I, GaydosCA, et al. Hormonal contraceptive use, cervical ectopy, and theacquisition of cervical infections. Sex Transm Dis.2004;31:561-567.
- 5. Anter SS. Clinical effect of tranexamic acid on curing cervicalerosion. IJIR. 2017;3:5-9.
- Machado Junior LC, Dalmaso AS, Carvalho HB. Evidence forbenefits from treating cervical ectopy: literature review. Sao PauloMedical Journal. 2008;126:132-139.
- 7. Reed MR, Woolley LT. Uses of tranexamic acid. BJA Education.2015;15:32-7.
- 8. Dries DJ. Tranexamic acid: is it about time? Lancet.2018;391:97-98.
- 9. Bigliardi PL, Alsagoff SA, El-Kafrawi HY, Pyon JK, Wa CT, VillaMA. Povidone iodine in wound healing: A review of currentconcepts and practices. Int J Surg. 2017;44:260-268.
- 10. CRASH-2 Trial Collaborators. Effects of tranexamic acid ondeath, vascular occlusive events, and blood transfusion in traumapatients with significant haemorrhage (CRASH-2): a randomised,placebo-controlled trial. Lancet. 2010;376:23-32.