OPEN ACCESS Freely available online

Journal of Clinical Microbiology and Antimicrobials

Commentary

Parasitic Diseases in Children: Diagnosis, Treatment and Prevention

Fariha Guez*

Department of Microbiology, University Hospital Aachen, Aachen, Germany

ABOUT TO STUDY

Parasitic diseases remain a significant health challenge worldwide, especially in children. The World Health Organization (WHO) estimates that over 1 billion people are affected by parasitic infections, with a substantial proportion being children in low- and middle-income countries. These infections can lead to severe morbidity, developmental delays, and even mortality. The ability to diagnose parasitic infections early, provide appropriate treatment, and implement preventive strategies is critical for reducing the burden of these diseases. This article explores the key aspects of parasitic diseases in children, including diagnosis, treatment, and prevention.

Common parasitic diseases in children

Intestinal protozoa infections: Protozoa are single-celled organisms that can infect the intestines, causing diarrhea, malnutrition, and dehydration. Common protozoal infections include:

Giardiasis, caused by *Giardia lamblia*, is prevalent in children in areas with poor sanitation. It causes chronic diarrhea, abdominal pain, and malabsorption.

Amebiasis, caused by *Entamoeba histolytica*, can lead to dysentery, liver abscesses, and, if untreated, severe complications.

Cryptosporidiosis, caused by *Cryptosporidium*, is known for causing watery diarrhea, which can be particularly dangerous for malnourished children.

Helminth infections: Helminths, or worms, are another major cause of morbidity in children. These include:

Ascariasis, caused by *Ascaris lumbricoides* (roundworms), leads to intestinal obstruction, malnutrition, and stunted growth in children.

Trichuriasis, caused by *Trichuris trichiura* (whipworms), can result in rectal prolapse, malnutrition, and anemia.

Hookworm disease, caused by *Ancylostoma duodenale* or *Necator americanus*, can cause anemia and malnutrition.

Malaria: Malaria, caused by *Plasmodium* species, is a major parasitic infection affecting children, especially in sub-Saharan Africa. It is transmitted through the bite of an infected mosquito and leads to high fever, chills, anemia, and, if untreated, organ failure and death. Children under the age of 5 are particularly at risk.

Toxocariasis: Toxocariasis, caused by the roundworms *Toxocara canis* and *Toxocara cati* (from pet dogs and cats), is another parasitic infection that can cause visceral larva migrans or ocular toxocariasis in children. The larvae migrate through the tissues, potentially leading to blindness and organ damage.

Diagnosis of parasitic diseases in children

Accurate diagnosis of parasitic diseases is critical for effective treatment. Diagnosis typically involves a combination of clinical evaluation, laboratory tests, and imaging, depending on the suspected parasite.

Microscopic examination of stool: The most common method for diagnosing intestinal protozoa and helminths is the microscopic examination of stool samples. Fecal flotation and concentration techniques help identify the eggs or cysts of helminths and protozoa, such as *Giardia*, *Entamoeba histolytica*, and *Ascaris*.

Antigen detection: Enzyme-Linked Immunosorbent Assays (ELISA) and other immunological tests can detect specific antigens released by parasites, such as *Giardia* or *Cryptosporidium*. These tests are particularly useful for detecting infections that are difficult to identify by microscopy.

Polymerase Chain Reaction (PCR): PCR is a powerful molecular diagnostic tool that amplifies the DNA or RNA of the parasite, allowing for highly sensitive and specific detection. It can be used for detecting protozoa and helminths, especially in cases where stool microscopy is inconclusive.

Blood tests: Blood tests are essential for diagnosing malaria and Toxocariasis. In malaria, thick and thin blood smears are used to identify *Plasmodium* species, while blood eosinophil counts can help diagnose helminth infections like ascariasis and hookworm.

Correspondence to: Fatima Guez, Department of Microbiology, University Hospital Aachen, Aachen, Germany; E-mail: guezfariha@bfarm.de

Received: 30-Dec-2024, Manuscript No. JCMA-24-36620; **Editor assigned:** 01-Jan-2025, PreQC No. JCMA-24-36620 (PQ); **Reviewed:** 15-Jan-2025, QC No. JCMA-24-36620; **Revised:** 10-Jul-2025, Manuscript No. JCMA-24-36620 (R); **Published:** 17-Jul-2025, DOI: 10.35248/ JCMA.25.09.227

Citation: Guez F (2025) Parasitic Diseases in Children: Diagnosis, Treatment and Prevention. J Clin Microbiol Antimicrob. 9:227

Copyright: © 2025 Guez F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Imaging: In cases where the infection has spread to other organs (e.g., liver abscess in Amebiasis), imaging techniques such as ultrasound or CT scans can help identify parasitic cysts or organ damage.

Treatment of parasitic diseases in children

The treatment of parasitic infections in children varies depending on the type of parasite involved. Most parasitic diseases are treatable with antiparasitic medications, but timely treatment is essential to avoid complications.

Intestinal protozoa: Giardiasis is typically treated with metronidazole or tinidazole. These medications are effective in clearing the infection and alleviating symptoms.

Amebiasis is treated with metronidazole or tinidazole for the initial infection and paromomycin or iodoquinol for asymptomatic cyst carriers.

Cryptosporidiosis generally resolves on its own in immunocompetent children, but in severe cases, nitazoxanide may be used.

Helminths: Ascariasis can be treated with albendazole, mebendazole, or ivermectin. These medications are effective in eliminating roundworms and improving nutrition in affected children.

Hookworm infections are treated with albendazole or mebendazole.

Trichuriasis is usually treated with mebendazole or albendazole.

Malaria: Malaria in children is treated with Artemisinin-based Combination Therapies (ACTs), which are the first-line treatment for *Plasmodium falciparum* malaria. Severe malaria may require intravenous artesunate or quinine.

Toxocariasis: Treatment for Toxocariasis often involves albendazole or mebendazole. If the infection leads to severe organ damage, corticosteroids may be required to reduce inflammation.

Prevention of parasitic diseases in children

Preventing parasitic infections in children requires a multifaceted approach that includes immunization, vector control, sanitation improvement, and education.

Improved sanitation: The most effective way to prevent many parasitic infections, particularly those caused by intestinal helminths and protozoa, is improving access to clean water and sanitation. Proper handwashing and hygiene practices are crucial for reducing transmission.

Vector control: In malaria-endemic regions, vector control strategies, such as the use of insecticide-treated bed nets and indoor residual spraying, can significantly reduce transmission. Early diagnosis and prompt treatment are key to malaria control in children.

Vaccination: Vaccines for parasitic diseases are limited, but there is progress in the development of vaccines for diseases like malaria. The RTS,S/AS01 malaria vaccine, approved for use in some African countries, offers partial protection against *Plasmodium falciparum*.

De-worming programs: Regular de-worming programs targeting school-age children can help reduce the prevalence of intestinal helminths. These programs provide medication to large numbers of children in endemic areas, helping reduce the overall parasite burden.

Health education: Educating families about personal hygiene, the importance of clean water, and avoiding contact with contaminated environments is essential in preventing parasitic diseases.

CONCLUSION

Parasitic diseases continue to pose significant health risks to children, particularly in resource-limited settings. Early diagnosis, effective treatment with antiparasitic medications, and preventive strategies such as improved sanitation, vector control, and health education are crucial in managing these infections. Continued efforts to enhance global health infrastructure and access to diagnostic and therapeutic tools will be key to reducing the burden of parasitic diseases in children. As new treatments and vaccines emerge, the fight against parasitic infections becomes more promising, offering hope for a future with fewer children suffering from these preventable diseases.