

Pharmacokinetics of Menaquinone-7 (Vitamin K2) in Healthy Volunteers

Knapen MHJ, Vermeer C*, Braam LAJLM and Theuwissen E

VitaK, Maastricht University, Oxfordlaan 70, 6229 EV Maastricht, The Netherlands

Abstract

Objective: Because of the wide-spread use of menaquinone-7 (MK-7, a form of vitamin K2) supplements, we have investigated its pharmacokinetics following a single intake from various formulations.

Methods: Different formulations were compared in four human intervention studies with healthy volunteers. Participants received either capsules (filling material: linseed oil; MK-7 carrier material: sunflower oil) or tablets (filling material: dicalcium phosphate dehydrate (DCPD) and cellulose; MK-7 carriers: casein, Arabic gum, or sunflower oil). MK-7-absorption profiles (24 h area-under-the-curves, 24 h AUCs) were used to compare intestinal absorption. Special attention was given to intra- and inter-subject variability.

Results: The absorption (time-to-peak) of MK-7 from tablets was slower than from capsules, 6 h as compared to 2 h-4 h, respectively. Probably, the oily matrix of the capsules released the lipophilic MK-7 more rapidly than the powder matrix of the tablets. We further found a dose response relationship for MK-7 at 24 h after the single-dose intakes (at group level). As compared to baseline, circulating MK-7 levels were still elevated at 24 h after intake, confirming the relatively long half-life of MK-7. The different MK-7 carrier materials showed similar 24 h-absorption profiles, indicating that the carrier was not influencing MK-7 absorption. Next to high intra-subject variability, high inter-subject variability was seen in the different studies. This was found for both, peak height and for total absorption (measured as 24 h-AUC). While some subjects showed a linear dose-response relation, other subjects had similar 24 h-AUCs after intake of MK-7 at different dosages

Conclusions: We have demonstrated that on a group level the bioavailability (24 h absorption) of MK-7 was similar for capsules and tablets, and also for the different MK-7 carriers. MK-7 absorption from the various formulations showed high intra- and inter-individual differences.

Keywords: Absorption; Bioavailability; Capsules; Menaquinone-7; Tablets; Vitamin K2

Introduction

Vitamin K functions as a cofactor for the γ -glutamate carboxylation of proteins involved in blood coagulation (factors II, VII, IX, X), calcium and bone metabolism (osteocalcin), cell growth regulation (Gas6), as well as soft tissue calcification (matrix Gla-protein) [1]. The carboxylated glutamate residues are called Gla-residues, and the proteins in which these unusual amino acids occur are generally referred to as the Gla-protein family. The Gla-residues in these proteins form strong calcium-binding groups that are required for their biological function. Hence, vitamin K plays a key role in important metabolic processes.

The two forms of vitamin K present in the human diet are phylloquinone (also known as vitamin K1) and the group of menaquinones (collectively known as vitamin K2). The nomenclature of the menaquinones is MK-n, where n stands for the number of unsaturated isoprenoid residues in the aliphatic side chain. Dietary relevant short-chain menaquinones are MK-4 through MK-6, of which MK-4 is by far the most abundant one; long-chain menaquinones are MK-7 through MK-10, of which MK-7 is the most prominent in the Japanese diet and MK-8 and MK-9 in the Western diet. Following intake, circulating vitamin K levels rapidly increase to reach their peak value between 2 h and 6 h postprandially, followed by a rapid decline for phylloquinone and MK-4, whereas the long-chain menaquinones were reported to have more complex pharmacokinetics and disappear much slower from the circulation [2-4]. Differences in pharmacokinetics of the various K-vitamins are probably related to differences in their molecular structure and to a number of unknown factors, including the food matrix, nature and amount of macronutrients, activity of digestive enzymes, transport efficiency across the intestinal cell, and bile production, and other individual conditions [5].

Several population-based studies have reported specific health effects of the long-chain menaquinones, notably MK-7, MK-8 and MK-9 [6-10]. Because of the wide-spread use of MK-7-containing supplements, we were interested to investigate in more detail pharmacokinetics of MK-7 following a single intake from various formulations. We compared different formulations: capsules and tablets, and different MK-7 carrier materials in four independent human intervention studies with healthy volunteers. Circulating MK-7 profiles covering 24 h after a single intake were used to compare intestinal absorption, and special attention was given to intra- and inter-subject variability.

Methods

Study 1

In this double-blind, randomized, cross-over study, absorption of MK-7 (MenaQ7, Nattopharma, Norway; daily dose of 75 μ g) from tablets of two different raw material suppliers was studied. These tablets contained the inert filling material dicalcium phosphate dehydrate (DCPD) and microcrystalline cellulose and MK-7 in three different carrier materials i.e. sunflower oil (supplier 1), soybean oil

*Corresponding author: Cees Vermeer, VitaK, Maastricht University, Oxfordlaan 70, 6229 EV Maastricht, The Netherlands, Tel: 0031-433885865; Fax: 0031-433885889; E-mail: c.vermeer@vitak.com

Received January 29, 2014; Accepted March 26, 2014; Published March 29, 2014

Citation: Knapen MHJ, Vermeer C, Braam LAJLM, Theuwissen E (2014) Pharmacokinetics of Menaquinone-7 (Vitamin K2) in Healthy Volunteers. J Clin Trials 4: 160. doi:10.4172/2167-0870.1000160

Copyright: © 2014 Knapen MHJ, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

	Study 1 (n=24)	Study 2 (n=69)			Study 3 (n=16)		Study 4 (n=12)	
Trial reg. nr.	NCT01638143	NCT00858767			NCT00742768		No	
Design	Double-blinded, randomized, cross-over	Double-blinded, randomized, parallel			Double-blinded, randomized, cross-over		Open, randomized, cross-over	
Study products	Tablets	Tablets	Tablets	Tablets	Capsules	Gelpell™ caps	Capsules	Food
Number/amount	1	2	2	3	4	4	4	40g
Daily dose	75 µg	90 µg	90 µg	90 µg	180 µg	180 µg	360 µg	360 µg
MK-7 carrier material	Sunflower oil, soybean oil or maltodextrin	Casein, Arabic gum or sunflower oil			Sunflower oil		Sunflower oil or natto food	
Filling material of supplements	Microcrystalline cellulose, DCPD	DCPD, microcrystalline cellulose			Linseed oil	Confidential	Linseed oil	
Time of intake	With breakfast	With breakfast			With breakfast		With breakfast	
Standardized breakfast / lunch	Yes	Yes			Yes		Only breakfast	
Sampling times	0 h, 2 h, 4 h, 6 h, 8 h, 24 h	0 h, 2 h, 4 h, 6 h, 8 h, 24 h			0 h, 2 h, 4 h, 6 h, 8 h, 24 h		0 h, 2 h, 4 h, 6 h, 8 h, 24 h	
Vitamin K-rich food restriction	Yes	Yes			Yes		Yes	

Table 1: Study design of four randomized clinical trials on MK-7 absorption.

(supplier 2), or maltodextrin (supplier 2) (Table 1). These three types of tablets were compared in a cross over design. Healthy men and women, aged between 20 and 40 years, were recruited in November 2010 in the southern region of the province of Limburg through placing an advertisement. Exclusion criteria were: BMI <18.5 kg/m² and >30 kg/m², metabolic or gastrointestinal diseases, chronic degenerative or inflammatory diseases, anemia, pregnancy, diabetes mellitus, soy allergy, abuse of drugs/alcohol, use of corticosteroids, oral anticoagulants, salicylates, antibiotics, and use of vitamin K supplements. Twenty-four participants (12 male / 12 female) who met the inclusion criteria were randomized to receive the first MK-7 tablet (a single dose of 75 µg MK-7). After a 2-week wash-out period, participants switched to the second type of tablet, and after another 2-week wash-out period they received the third type of tablet. On the first intervention day, blood was sampled after an overnight fast (t=0 h). Immediately after sampling, participants received a standardized breakfast (two rolls or toast with 20 g of butter, jam, 150 mL of demi-skimmed milk and tea) together with the MK-7 tablet. Subsequently, blood was sampled at 2 h, 4 h, 6 h, 8 h and 24h after the single intake of MK-7. At t=4 h a standardized lunch was offered consisting of 2 rolls and toast with butter, meat and/or jam, 250 g of demi-skimmed yogurt and coffee or tea. Two weeks before the start of the study and during the study, subjects were asked to refrain from consuming foods rich in phylloquinone (spinach, green kale, broccoli, Brussels sprouts) and rich in menaquinones (curd, cheese, natto).

Study 2

In this double-blind, randomized, parallel study, we compared absorption of MK-7 (MenaQ7, Nattopharma, Norway; daily dose of 90 µg) from tablets (filling material: DCPD and microcrystalline cellulose) containing three different MK-7 carriers, namely casein, Arabic gum, and sunflower oil (Table 1). Healthy men (n=28) and women (n=41), aged between 20 and 40 years, were recruited in October 2008 in the southern region of Limburg through placing small advertisements. Exclusion criteria were BMI >30 kg/m², metabolic or gastrointestinal diseases, chronic degenerative or inflammatory diseases, anemia, soy allergy, abuse of drugs/alcohol, use of corticosteroids, oral anticoagulants, use of vitamin K supplements, and high dietary intake of vitamin K. Eligible participants were randomly divided to consume daily either two casein tablets, two Arabic gum tablets, or three sunflower oil tablets during 8 weeks. After an overnight fast and a first blood sample (t=0 h), the participants took the MK-7 tablets together

with a standardized breakfast, followed by a standardized lunch at t=4 h (as detailed for Study 1). The subsequent samples were taken at 2 h, 4 h, 6 h, 8 h and 24 h. One week before the start of the study and during the study, participants were asked to refrain from consuming foods rich in vitamin K. This study has been shortly mentioned in the recently published paper "Vitamin K status in healthy individuals" [11]. However, different data were presented, namely the MK-7-induced changes in circulating desphospho-uncarboxylated matrix Gla-protein (marker for vitamin K status) in relation to baseline.

Study 3

Newly developed MK-7 (MenaQ7, Nattopharma, Norway) capsules with sunflower oil (Gelpell™, matrix pellets from gelatin, Gelpell AG, Swiss Caps AG, Switzerland) were compared to conventional gelatin soft gel capsules with sunflower oil and MK-7 (MenaQ7, Nattopharma, Norway) in a double-blind, randomized, cross-over study (Table 1). The difference between both capsules was that the MK-7 in the Gelpell™ capsules was encapsulated in microspheres, a technique known to give more complete absorption. Healthy men (n=8) and women (n=8), aged between 20 and 30 years, were recruited June 2008 in the southern region of Limburg through placing small advertisements. Exclusion criteria were: BMI >30 kg/m², metabolic or gastrointestinal diseases, chronic degenerative or inflammatory diseases, anemia, soy allergy, use of corticosteroids, oral anticoagulants, and use of vitamin K supplements. Eligible participants were randomized to receive a single dose of 180 µg MK-7 (four soft gel or four Gelpell™ capsules; 45 µg MK-7 per capsule). Every two weeks, participants switched to another treatment. The capsules were consumed together with the standardized breakfast followed by lunch as defined for Study 1. Blood sampling of the participants took place before (0 h, after an overnight fast) and after the intake of the capsules at 2 h, 4 h, 6 h, 8 h and 24 h. The week before intake of the capsules and during the study participants were asked to refrain from consuming foods rich in vitamin K. During the wash-out period, subjects were allowed to consume their normal diets, but not to consume more than normal amounts of green vegetables (200 g/day of spinach, broccoli, etc.) and (curd) cheese (< 50 g/day), and no natto.

Study 4

This study had an open, randomized, crossover design (Table 1). Healthy men (n=6) and women (n=6), aged between 20 and 30 years, were recruited among students of Maastricht University in January 2008. Exclusion criteria were: BMI >30 kg/m², metabolic

or gastrointestinal diseases, chronic degenerative or inflammatory diseases, use of corticosteroids, oral anticoagulants, and vitamin K supplements. Eligible volunteers were randomized in two groups to receive a single dose of either 40 g homogenized natto (360 µg of MK-7) or four MK-7 (MenaQ7, Nattopharma, Norway) capsules (90 µg MK-7 per capsule with sunflower oil) at breakfast (as defined for Study 1). Subsequently, blood samples were taken at 0 h, 2 h, 4 h, 6 h, 8 h and 24 h after intake of the single dose natto or the MK-7 capsules. After a 2-week wash-out period, participants switched to the second treatment. One week before the study and during the study period, subjects were not allowed to consume vitamin K-rich foods; only small amounts of (curd) cheese (<100 g/d) were acceptable during this period.

Ethics Statement

All studies were conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the Medical Ethics Committee of the Maastricht University Medical Center (Maastricht, The Netherlands). Written informed consent was obtained from all subjects before entering the study. Trial registration codes were clinicaltrials.gov: NCT01638143 (study 1), NCT00858767 (study 2), and NCT00742768 (study 3). Study 4 was not registered.

Circulating MK-7 levels

Plasma MK-7 levels (ng/mL) were determined according to standard procedures. In short, samples were extracted with hexane and after pre-purification on silica columns, they were analyzed by high performance liquid chromatography (HPLC) using a C-18 reversed phase column with online zinc reduction and fluorescence detection. Vitamin K1-25 (a synthetic form of vitamin K1 containing 5 isoprenoid residues) was used as an internal standard. Individual AUC values were calculated from the individual MK-7 absorption profiles using the trapezoidal rule. MK-7 measurements were performed at the research laboratory of VitaK BV (ISO9001-certified lab, Maastricht, The Netherlands).

Statistics

To ensure homogeneity of the four studies, we used the following sampling points for statistics: 0 h, 2 h, 4 h, 6 h, 8 h, and 24 h. Within one study, between-group differences in AUC values were analyzed by the non-parametric Mann-Whitney, Kruskal-Wallis, or Wilcoxon-Signed Rank tests. Differences were considered statistically significant

at $p < 0.05$ (SPSS 19). Inter-study and intra-subjects comparisons regarding MK-7 absorption were done by comparison of the time-to-peak, peak heights, and AUC-values. Finally, we identified three subjects that participated in studies 1, 2, and 3, and we could therefore perform the intra-subject comparisons.

Results

Inter-study comparisons regarding MK-7 absorption

Based on the change from baseline of the biomarkers (UACR, plasma BNP) and their percent change, the profile will be compared between the treatment groups and reviewed by repeated-measure analysis of variance with measurement time points as repetition. Primary endpoints will be analyzed by the closed testing procedure in the order of UACR, plasma BNP, and their percent change in consideration of multiplicity.

Study 1

The time-to-peak was 6 h after intake of one tablet of MK-7 (Table 2 and Figure 1A). A first peak was observed after 2 h of intake, with a second peak at 6 h corresponding to a peak height of about 1 ng/mL for all three formulations. The 24 h AUC values were calculated to range between 13.8 ± 4.3 and 16.1 ± 6.7 ng/mL x h. At 24 h after intake, circulating MK-7 levels were still increased as compared to baseline. Similar patterns were found for all three formulations, and no significant differences were observed between the average absorption curves (time-to-peak and AUC) for each formulation (Figure 1A).

Study 2

The time-to-peak was 6 h after intake of the MK-7 tablets (Table 2 and Figure 1B), to peak heights of 1.63 ± 0.74 ng/mL, 1.94 ± 1.14 ng/mL, and 1.64 ± 1.07 ng/mL for the casein, Arabic gum, and linseed oil carriers, respectively. The corresponding 24 h AUCs were calculated to be 23 ± 9 ng/mL x h, 22 ± 9 ng/mL x h, and 21 ± 8 ng/mL x h, respectively. No between-group differences were found with regards to the calculated AUCs. At 24 h after intake, circulating MK-7 levels were still higher than those at baseline for all three groups.


Study 3

The time-to-peak for MK-7 was 2 h and 4 h after intake of the standard soft gel and GelpellTM capsules, respectively (Table 2 and Figure 1C). The corresponding peak heights were 2.22 ± 0.93 ng/mL

	MK-7 at t0 (ng/mL)	MK-7 at 24 h (ng/mL)	Time to peak* (h)	Peak-Height† (ng/mL)	AUC (ng/mL x h)	p#
Study 1						
Sunflower oil	0.27 ± 0.15	0.49 ± 0.15	6	0.99 ± 0.38	16.08 ± 6.73	P=0.105
Soybean oil	0.27 ± 0.18	0.48 ± 0.18	6	0.83 ± 0.37	14.33 ± 4.72	
Maltodextrin	0.30 ± 0.15	0.44 ± 0.14	6	0.86 ± 0.37	13.78 ± 4.26	
Study 2						
Casein	0.27 ± 0.23	0.79 ± 0.39	6	1.63 ± 0.74	22.59 ± 8.86	0.904
Arabic Gum	0.31 ± 0.32	0.70 ± 0.32	6	1.94 ± 1.14	21.69 ± 8.56	
Linseed oil	0.23 ± 0.40	0.61 ± 0.26	6	1.64 ± 1.07	21.13 ± 8.20	
Study 3						
Sunflower oil	0.14 ± 0.29	1.19 ± 0.58	2	2.22 ± 0.93	37.44 ± 14.73	0.049
Sunflower oil (Gelpell)	0.18 ± 0.41	1.43 ± 0.50	4	3.38 ± 2.37	43.34 ± 14.18	
Study 4						
Sunflower oil	0.13 ± 0.13	1.51 ± 0.73	4	6.77 ± 3.30	65.90 ± 28.60	0.773
Natto	0.13 ± 0.12	1.94 ± 0.81	6	5.91 ± 2.75	63.03 ± 25.18	

Data are given as mean values ± SD. *Time to the highest peak; †Highest peak; #Between-group difference in AUC as tested by the non-parametric Kruskal-Wallis or Wilcoxon-Signed Rank tests within one study Abbreviations: AUC, area-under-the-curve; MK-7, menaquinone-7; n.s., no significant differences between any arm.

Table 2: Absorption measures of MK-7 from single-dose capsules and tablets.



and 3.38 ± 2.37 ng/mL. The 24 h AUCs were 37 ± 15 ng/mL x h and 43 ± 14 ng/mL x h for the standard soft gel and Gelpell™ capsules, respectively. A significant between-group difference was found with regards to the calculated AUCs, showing a better absorption for the patented Gelpell product. At 24 h after intake, circulating MK-7 levels were still higher than those at baseline for both groups.

Study 4

The time-to-peak was 4h after intake of the MK-7 capsules (Table 2 and Figure 1D), corresponding to a peak height of 6.77 ± 3.30 ng/mL. The 24 h AUC was calculated to be 66 ± 29 ng/mL x h. Again, circulating MK-7 levels were still elevated at 24 h after intake as compared to baseline. It was also clear that MK-7 uptake from natto was slightly slower than from the capsules, with a time-to-peak of 6 h (Figure 1D). Since the AUC values for uptake from natto (63 ± 25 ng/mL x h) was comparable to that of the capsules, the efficacy of MK-7 was regarded to be similar for both groups.

Inter-study comparisons

As can be seen in Table 2, the tablets gave a slower absorption of MK-7 than the capsules, 6 h as compared to 2 h-4 h, respectively. A dose-response curve was prepared from the average AUC values obtained for all four different dosages (Figure 2). Although the number of data points is only small, curve fitting gave a linear relationship between intake and 24 h-AUC. The fact that the line crosses the Y-axis at 7.4 ng/mL x h is explained by the baseline level of circulating MK-7 found in most participants.

Intra- and inter-subject comparisons of MK-7 absorption

There were three individuals who participated in three trials (studies 1-3), and as is shown in Table 3, there were substantial differences in the MK-7 absorption from the 75, 90 and 180 µg formulations. In subject 1 the respective AUC values were 9.9, 24.6 and 39.3 ng/mL x h, and for subject 2 these values were closely similar: 8.9, 22.0 and 31.7 ng/mL x h. For subject 3, however, the dose-response effect was negligible with respective AUC values of 22.4, 29.6 and 28.6 ng/mL x h, respectively.

Besides the observed intra-subject variability, high inter-subject variability was seen in studies 1, 2, and 3. Some subjects had little absorption, while other subjects had good absorption of MK-7 from the tablets. While some subjects showed a dose-response relation, other subjects had similar absorption after intake of MK-7 at different dosages. Differences were also seen in time-to-peak, peak height, and 24 h AUC

Discussion

Because of the widespread use of MK-7 containing supplements, it is important to study its pharmacokinetics from different formulations. We therefore measured MK-7 absorption following a single intake from oil-based capsules and powder-based tablets in four intervention trials. Tablets gave a slower absorption of MK-7 (time-to-peak) than capsules, 6 h as compared to 2 h-4 h, respectively. Probably, the oily matrix of the capsules released the lipophilic MK-7 more rapidly than the powder matrix of the tablets. Within the dose range between 75 and 360 µg/day, we found a linear 24 h dose-response relationship for MK-7 after the single-dose intakes; the curve shown in Figure 2 is based




Figure 2: Dose-response relationship between MK-7 intake and 24h-AUC circulating MK-7 values.

Subject	Study product	Daily dose (µg)	MK-7 at t0 (ng/mL)	MK-7 at 24 h (ng/mL)	Time to peak (h)*	Peak height† (ng/mL)	AUC (ng/mL x h)
Subject 1							
Sunflower oil	Tablets	75	0.34	0.42	6	0.49	9.92
Casein	Tablets	90	0	1.07	4	2.92	24.64
Sunflower oil	Capsules	180	0	1.53	2	2.13	39.27
Subject 2							
Sunflower oil	Tablets	75	0	0.4	6	0.59	8.87
Casein	Tablets	90	0	0.58	2	1.8	22.02
Sunflower oil	Capsules	180	0	1.11	2	2.3	31.73
Subject 3							
Sunflower oil	Tablets	75	0.37	0.45	6	2.04	22.43
Casein	Tablets	90	0.44	1.06	6	2.51	29.59
Sunflower oil	Capsules	180	0.13	0.77	4	1.21	28.55

Data are given as mean values ± SD. *Time to the highest peak; †Highest peak. Abbreviations: AUC, area-under-the-curve; MK-7, menaquinone-7

Table 3: Absorption measures of MK-7 from single-dose capsules and tablets in 3 healthy volunteers.

on the average AUC values from all four studies, and is only valid for the narrow dose range investigated. Circulating MK-7 levels were still elevated at 24 h after intake as compared to baseline, confirming the relatively long half-life of MK-7 [4].

In study 2, the three different MK-7 carrier materials used in the tablets showed similar absorption profiles, indicating that the carrier material was not influencing 24 h absorption. In study 3, two different encapsulation technologies were compared with regards to MK-7 absorption, i.e. standard soft gel and Gelpell™ capsules. A significant between-group difference was found with regard to the calculated 24h-AUCs, demonstrating that the Gelpell™ capsules give better absorption of MK-7 than the standard soft gel ones. This confirms the increased bioavailability as reported by the manufacturer (<http://www.gelpell.com/EN/download.html>). Next to high intra-subject variability, high inter-subject variability was seen in studies 1, 2, and 3. Some subjects had no or poor absorption, while other subjects had good absorption of MK-7 from the tablets. Whereas some subjects showed a linear dose-response relation, other subjects had similar 24 h-AUCs after intake of MK-7 at different dosages. These results further indicate that one subject can have variable absorption on different days, as seen in different studies. Absorption efficiency is influenced by a number of external factors, including the food matrix and effectors of absorption (lipids, dietary fiber, drugs); also individual conditions, like age, sex, and genetic factors are assumed to govern absorption. Although the

practical external influences on absorption in the four studies were kept similar as good as possible (i.e. overnight fast, intake with breakfast, standardized meals, restriction on vitamin K-rich foods), these may have affected absorption as well. A limitation to our study is that we evaluated retrospectively four individual trials; ideally, the different formulations need to be compared in one intervention trial.

To our knowledge, only three studies are available on single-dose kinetics of menaquinones after consumption of food supplements [12] or fortified foods [3,4]. Sato et al. [12] showed that MK-7 was well absorbed in Japanese healthy women after single-dose (420 µg MK-7) intake with maximum serum levels at 6 h after intake. In our studies 3 and 4 (using oil-based capsules), the maximum MK-7 peak was seen earlier, i.e. at 2 h-4 h after intake of the capsules. Powder was used in the Japanese gelatin capsules, which may explain the difference in time-to-peak. The maximum peak value in the Japanese participants equaled the maximum peak value after intake of 360 µg MK-7 in study 4 (6.77 ± 3.30 ng/mL). Previously, our group showed that phylloquinone, MK-4, and MK-9 consumed as vitamin K-enriched (2 µmol of each vitamin K form) corn oil gave lower plasma peaks for MK-4 and MK-9 than for phylloquinone [3]. Maximal circulating levels were reached at 2.5, 4, and 5 h postprandially for MK-4, phylloquinone, and MK-9, respectively. Their half-lives differed substantially: whereas phylloquinone and MK-4 disappeared overnight, the estimated half-

life for the long-chain MK-9 was about 60 h, which is consistent with the long half-life time of MK-7 reported in the present paper.

In conclusion, we have demonstrated similar MK-7 bioavailability for capsules and tablets, and for the different MK-7 carrier materials used in the tablets. MK-7 absorption from various formulations showed high intra- and inter-individual differences; the mechanism behind it is not yet understood. More research is needed to clarify these differences in bioavailability of MK-7.

Acknowledgement

CV designed the research; ET analyzed the data, all authors contributed to the writing of the paper.

Conflict of Interest

All authors declare no competing financial interests.

References

1. Shearer MJ, Newman P (2008) Metabolism and cell biology of vitamin K. *ThrombHaemost* 100: 530-547.
2. Schurgers LJ, Vermeer C (2000) Determination of phylloquinone and menaquinones in food. Effect of food matrix on circulating vitamin K concentrations. *Haemostasis* 30: 298-307.
3. Schurgers LJ, Vermeer C (2002) Differential lipoprotein transport pathways of K-vitamins in healthy subjects. *BiochimBiophysActa* 1570: 27-32.
4. Schurgers LJ, Teunissen KJ, Hamulyák K, Knapen MH, Vik H, et al. (2007) Vitamin K-containing dietary supplements: comparison of synthetic vitamin K1 and natto-derived menaquinone-7. *Blood* 109: 3279-3283.
5. Jones KS, Bluck LJ, Wang LY, Stephen AM, Prynne CJ, et al. (2009) The effect of different meals on the absorption of stable isotope-labelled phylloquinone. *Br J Nutr* 102: 1195-1202.
6. Geleijnse JM, Vermeer C, Grobbee DE, Schurgers LJ, Knapen MH, et al. (2004) Dietary intake of menaquinone is associated with a reduced risk of coronary heart disease: the Rotterdam Study. *J Nutr* 134: 3100-3105.
7. Gast GC, de Roos NM, Sluijs I, Bots ML, Beulens JW, et al. (2009) A high menaquinone intake reduces the incidence of coronary heart disease. *NutrMetabCardiovasc Dis* 19: 504-510.
8. Beulens JW, Bots ML, Atsma F, Bartelink ML, Prokop M, et al. (2009) High dietary menaquinone intake is associated with reduced coronary calcification. *Atherosclerosis* 203: 489-493.
9. Nimptsch K, Rohrmann S, Linseisen J (2008) Dietary intake of vitamin K and risk of prostate cancer in the Heidelberg cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC-Heidelberg). *Am J Clin Nutr* 87: 985-992.
10. Nimptsch K, Rohrmann S, Kaaks R, Linseisen J (2010) Dietary vitamin K intake in relation to cancer incidence and mortality: results from the Heidelberg cohort of the European Prospective Investigation into Cancer and Nutrition (EPIC-Heidelberg). *Am J Clin Nutr* 91: 1348-1358.
11. Theuwissen E, Magdeleyns EJ, Braam LA, Teunissen KJ, Knapen MH, et al. (2014) Vitamin K status in healthy volunteers. *Food Funct* 5: 229-234.
12. Sato T, Schurgers LJ, Uenishi K (2012) Comparison of menaquinone-4 and menaquinone-7 bioavailability in healthy women. *Nutr J* 11: 93.