

Pharmaceutical Misadventures in the Neonatal Intensive Care Unit

Steven M. Donn*

Department of Pediatrics, C.S. Mott Children's Hospital, University of Michigan, Ann Arbor Michigan, USA

DESCRIPTION

The newborn infant, especially those who are born prematurely, are among the most vulnerable patients in all of medicine. Organs and organ systems are incompletely formed and are still developing, metabolic pathways are also in a state of evolution, and both hepatic and renal functions are relatively inadequate compared to older children or adults. The developing brain and central nervous system may be affected and show evidence of dysfunction years later. Yet, infants in the milieu of the Neonatal Intensive Care Unit (NICU) are exposed to countless pharmaceutical agents of dubious efficacy and potentially dangerous toxicity without a sufficient evidence base to justify their use.

A recent study by Bamat and colleagues looked at medication use in preterm infants with severe bronchopulmonary dysplasia, the most severe long term respiratory morbidity in this population resulting from the combination of ventilator-induced lung injury, oxygen toxicity, and a developmental arrest of alveolarization. They performed a retrospective cohort study to identify the number of cumulative medication exposures and characterize the most frequently used agents. Astoundingly, in 43 hospitals surveyed, encompassing 3,252 patients, a median of 30 cumulative medication exposures per infant was found, with an interquartile range of 17-45. Diuretics, drugs of doubtful efficacy and significant toxicity, were the most frequently prescribed agents [1].

The history of neonatology is replete with examples of good ideas that turned into disaster. In 1959, the use of the antimicrobial, chloramphenicol was found to cause the "gray baby syndrome," characterized by severe hypotension, cyanosis, and even death. Some infants developed fatal aplastic anemia [2]. In the 1960's, Usher introduced the practice of using sodium bicarbonate therapy to maintain blood pH in preterm infants with respiratory distress syndrome. After some preliminary success, the dose of sodium bicarbonate was increased but led to a significant incidence of intraventricular hemorrhage [3]. The use of intravenous vitamin-E in the 1980s produced the "gaspings syndrome," resulting from the preservative benzyl alcohol. More than a dozen deaths were attributed to this. Affected infants developed metabolic acidosis which progressed to severe respiratory distress as well as seizures

and brain hemorrhages [4]. More recently, the prokinetic agent, cisapride, used to promote gastrointestinal motility, was found to be associated with prolongation of the QTc interval in very preterm infants, leading to serious cardiac arrhythmias and mortality and was withdrawn from the market [5].

These examples demonstrates the severity of unintended consequences in this population. The problem is underscored by the fact that most, pharmaceuticals administered in the NICU are off label (two-thirds to three-quarters) [6], and thus have not undergone adequate testing or gained approval for use by the US Food and Drug Administration. These infants are particularly susceptible to injury because of organ immaturity, especially the brain, and many pharmaceutical agents that are well tolerated in the adult population may have undesirable side effects in the preterm infant; many may not surface for years. Other potential dangers are unknown drug-drug interactions, and drug-nutrient interactions.

While neonatologists are intensivists and deal with a critically ill and unstable patient population and are quick to reach for therapeutic solutions, it would be the responsibility of all of us to remember the oath we took at the outset of our careers, "Primum non nocere." First, do no harm.

REFERENCES

1. Bamat NA, Kirpalani H, Feudtner C, Jensen EA, Laughon MM, Zhang H, et al. Medication use in infants with severe bronchopulmonary dysplasia admitted to United States children's hospitals. *J Perinatol.* 2019;39(9):1291-1299.
2. Choonara IA, Rieder MJ. Drug toxicity and adverse drug reactions in children: A brief historical review. *Paediatr Perinat Drug Ther.* 2002;5(1):12-18.
3. Dunn PM. Robert Usher. *Br Med J.* 2006;332(7556):1514.
4. Gershanik J, Boecler B, Ensley H, McCloskey S, George W. The gasping syndrome and benzyl alcohol poisoning. *N Engl J Med.* 1982;307(22):1384-1388.
5. Dubin A, Kikkert M, Mirmiran M, Ariagno R. Cisapride associated with QTc prolongation in very low birth weight preterm infants. *Pediatrics.* 2001;107(6):1313-1316.
6. Conroy S, McIntyre J, Choonara I. Unlicensed and off label drug use in neonates. *Arch Dis Child Fetal Neonatal Ed.* 1999;80(2):F142-F145.

Correspondence to: Steven M. Donn, Department of Pediatrics, C.S. Mott Children's Hospital, University of Michigan, Ann Arbor Michigan, USA, Tel: +1734763-4109; E-mail: smdonnm@med.umich.edu

Received: 16-Feb-2022, Manuscript No. PTCR-22-15875; **Editor assigned:** 21-Feb-2022, PreQC No. PTCR-22-15875 (PQ); **Reviewed:** 02-Mar-2022, QC No. PTCR-22-158755; **Revised:** 09-Mar-2022, Manuscript No. PTCR-22-15875 (R); **Published:** 16-Mar-2022, DOI: 10.35248/2161-0665.22.12.437.

Citation: Donn SM (2022) Pharmaceutical Misadventures in the Neonatal Intensive Care Unit. *Pediatr Ther.* 12:437.

Copyright: © 2022 Donn SM. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.