

# Physical Activity and Associated Factors among Patients Attending Hypertension Clinics in Mbarara City, Southwestern Uganda

Atwongyeire Night\*, Ssewanyana Anna Maria, Namukwaya Racheal, Mutesasira Edward, JJunju Fred, Nuwahereza Amon, Jean Damascene Niyonsenga, Zillah Whitehouse, Kazibwe Herman, Arubaku Wilfred

Department of Physical Medicine, Mbarara University of Science and Technology, Mbarara, Uganda

## ABSTRACT

**Background:** Physical activity has been shown to prevent mortality and morbidity among people with hypertension. Hypertension has been identified to affect about 33% of the adult population across the globe and 26.4% in Uganda. An increase in hypertension and its adverse outcomes have been observed and reported in Southwestern Uganda. This may be due to physical inactivity. However, there is limited evidence demonstrating physical activity among patients with hypertension within Southwestern Uganda. The current study investigated the extent of physical activity and associated factors among patients attending hypertension clinics in Mbarara City, Southwestern Uganda.

**Objective:** To investigate the extent of physical activity and associated factors among patients attending hypertension clinics in Mbarara City, Southwestern Uganda.

**Methods:** A descriptive, quantitative cross-sectional study was conducted. Participants' data was obtained using a structured researcher-administered questionnaire consisting of the International Physical Activity Questionnaire (IPAQ) long form, Motivation for Physical Activity Questionnaire (RM4-FM) and the Barrier to Being Active Quiz (BBAQ). Frequency distribution tables, Fischer's exact test and multivariate logistic regression were used to describe data and establish associations during data analysis. A p-value less than 0.05 with 95% confidence interval was considered to be statistically significant.

**Results:** Less than half of the participants (45.39%) were found to be physically active. Level of education, having heard of physical activity, place of residence, sedentary behaviour and social influence were associated with physical activity with the Fischer's exact test (p-value<0.05). On multivariate adjustment, only level of education (aOR=1.374; CI=1.055-1.790; p-value=0.018) and sedentary behaviour (aOR=0.276; CI=0.126-0.606; p-value=0.001) remained significant factors associated with physical activity. Majority of those who were active reported to be autonomously motivated. Most reported barriers were lack of skill, social influence and lack of willpower.

**Conclusion:** More than half of the patients with hypertension were physically inactive and almost half were inactive and sedentary. This is a double disaster especially for patients with hypertension; therefore, emphasis should be put on educating the public about the benefits of physical activity and availing a variety of physical activity options that engage all demographic groups.

**Keywords:** Physical activity; Hypertension; Sedentary behaviour

## INTRODUCTION

Across the globe, 7.6% of deaths due to cardiovascular diseases are attributed to physical inactivity [1]. Evidence also shows that

74.0% of the cardiovascular disease deaths associated with physical inactivity occurs in low and middle-income countries. Physical activity has been recommended in management of hypertension [1]. In 1968, one of the first clinical studies demonstrating the

**Correspondence to:** Atwongyeire Night, Department of Physical Medicine, Mbarara University of Science and Technology, Mbarara, Uganda, E-mail: atnightb@gmail.com

**Received:** 20-Jul-2023, Manuscript No. JPMR-23-25783; **Editor assigned:** 22-Jul-2023, Pre QC No. JPMR-23-25783(PQ); **Reviewed:** 08-Aug-2023, QC No. JPMR-23-25783; **Revised:** 17-Aug-2023, Manuscript No. JPMR-23-25783(R); **Published:** 25-Aug-2023, DOI: 10.35248/2329-9096.23.11.686

**Citation:** Night A, Maria SA, Racheal N, Edward M, Fred J, Amon N, et al (2023) Physical Activity and Associated Factors among Patients Attending Hypertension Clinics in Mbarara City, Southwestern Uganda. Int J Phys Med Rehabil. 11:686.

**Copyright:** © 2023 Night A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

antihypertensive effect of physical activity was published. The study found a 13.4 mmHg decrease in the mean systolic pressure and a 2 to 30 mmHg decrease in diastolic pressure, after the completion of the exercise training period [2]. More studies have recently been done to confirm the effect of physical activity on blood pressure control through various mechanisms including, decreasing sympathetic activity and reducing oxidative stress [3,4]. The notion that an individual can be both physically active and also sedentary has come as an eye-opener for most physical activity researchers, and some findings have examined the possible negative effect of sedentary behaviour over physical activity in physically active populations and sedentary behaviour has been reported to produce direct negative effects on outcomes of hypertension [5-7]. Evidence comes from who reported that each one hour of sedentary behaviour was associated with 0.06 mmHg increases in systolic pressure and 0.20 mmHg increase in diastolic pressure among hypertensives [8]. In addition, sedentary behaviour has been reported to compromise outcomes of hypertension leading to stroke and myocardial infarction among others [9]. Despite the known benefits of physical activity, physical inactivity especially among individuals with hypertension is still high across the globe ranging from 23% in adults to 81% in adolescents [10]. It is important to note that the prevalence of physical inactivity varies across various regions and countries and may be as high as 80% among some adult populations [11]. The major form of motivation to physical activity adherence has been reported to be autonomous motivation [12]. Additionally, found that autonomous regulation was the main predictor of exercise intensity, frequency and time among males and females [13]. The major barriers that have been reported include inadequate information about physical activity, fear of physical activity, health illiteracy and fear of changing lifestyle [14].

In East Africa, in a study in Kenya, it was found that majority (63.0%) of the patients with hypertension had low levels of physical activity [15]. In another study done in Kigali Rwanda, it was indicated that the major factors associated with physical activity among patients with hypertension were age, marital status, place of residence, and education level, among others, and the perceived motivators were perceived health benefits, self-efficacy and blood pressure control [16].

On that account, increasing incidences of adverse outcomes of hypertension like heart failure, renal failure and stroke among individuals with hypertension have been reported and observed in this region [17-19]. It is important to note that, in a recent study by at Mbarara Regional Referral hospital, hypertension was cited as a major contributing factor to stroke mortality [20]. These observations could be attributed to poor practice of lifestyle modifications such as physical inactivity among patients with hypertension. However, this argument cannot be fully supported since there is still limited evidence on physical activity among individuals with hypertension in this region. Relatedly, there is inadequate information on the factors associated with physical activity that might influence the extent of physical activity in this population. Therefore, this study sought to investigate the extent of physical activity and associated factors among patients attending hypertension clinics in Mbarara City, Southwestern Uganda.

## MATERIALS AND METHODS

### Study design

A descriptive, quantitative cross-sectional study using a researcher-administered questionnaire was conducted. Hypertensive patients were identified by the nurse and connected to the research members, and a consecutive sampling strategy was used to recruit participants in the study. Interviewer-administered questionnaires were adopted from the International Physical Activity questionnaire (IPAQ) long form to investigate the extent of physical activity, Motivation for Physical Activity Questionnaire (RM4-FM) to explore motivators to physical activity and the Barriers to Being Active Quiz (BBAQ) to explore barriers to physical activity.

### Study setting

The study was conducted at two hypertension clinics in Mbarara City, Southwestern Uganda. The two clinics were DMA Diagnostics and Laboratory Limited in Kamukuzi division and Mbarara Medical Specialist clinic located in Kakoba division. The participants got from these clinics were 133 patients and 99 respectively, obtained through proportionate sampling.

### Study variables

The level of physical activity (dependent variable) was analyzed by considering the metabolic equivalent of the task, METs-minutes per week and the days of physical activity. MET-minutes per participant were calculated using the formula (MET value) \* (time of activity in minutes per day \* number of days the activity was done in a week). The independent variables were the associated factors to physical activity. These included: Socio-demographic characteristics (including age, gender, place of residence, monthly income, occupation, marital status, level of education), duration of hypertension, awareness of physical activity, source of information about physical activity, sedentary behaviour, motivators for physical activity, barriers to physical activity and sedentary behaviour. Sedentary behaviour was measured from time spent sitting as average sitting time in the past seven days, assessed by the International Physical Activity Questionnaire (IPAQ). Motivators and barriers were measured using Motivation for Physical Activity Questionnaire (RM4-FM) and Barriers to Being Active Quiz (BBAQ)

### Data analysis

Given that the total study sample size was small, Fischer's exact test was used to test for association between physical activity and the independent variables. Multivariate logistic regression was also used to test for further association between physical activity and the independent variables. A p-value less than 0.05 was considered for statistical significance.

## RESULTS AND DISCUSSION

Overall, 141 participants participated in the study. Median age of the participants was 58 years (IQR=14). Table 1 shows that females accounted for 67.4% of the participants while males accounted for 32.6% and the age categories were: less than 35 years 3.6%, 35-49 years 18.4% and 50 years above 78.0%. The places of residence of the participants were: Mbarara 44.0%, Greater

Bushenyi 15.6%, Isingiro 13.5% and others 27.0%. The different occupations for the participants were: Farming 27.0%, business 23.4%, housewife 10.6%, retired 9.2% and others 29.8%. The monthly income of the participants was 0-500,000UGX 55.3%, 500,000-1M UGX 31.9% 1M UGX or more 12.8% while the marital status of the participants was: Married 73.8% single 6.4% divorced 5.7% and others 14.2%. The duration of hypertension among the participants was: less than one year 14.2%, 1-5 years 38.3% and more than 6 years 47.5%. The level of education of participants was; completed primary 22.0% secondary 19.9%, tertiary 34.0% and no formal education 24.1%. Those who had heard of physical activity were 85.8% and those who never heard of physical activity were 14.2%. Source of information

(from whom) included friends and family 17.7%, health workers 56.0%, mass media 12.1%. The overall physical activity referred to a combination of activities in the different domains. Table 2 shows that overall, of the 141 participants, 54.6% participants had low levels of physical activity, 40.4% achieved moderate levels of physical activity and 5.0% had high levels of physical activity. The factors found to be associated with physical activity on Fishers exact test included place of residence, social influence, level of education, information about physical activity and sedentary behaviour, as shown in Table 3. On bivariate and multivariate logistic regression, only two factors remained statistically significant and these were; sedentary behavior and level of education, as shown in Table 4.

**Table 1:** Demographic characteristics of study participants.

Characteristics	Frequencies	%
<b>Age group</b>		
less than 35 years	5	3.6
35-49 years	26	18.4
50 years and above	110	78
<b>Gender</b>		
Female	95	67.4
Male	46	32.6
<b>Place of residence</b>		
Greater Bushenyi	22	15.6
Isingiro	19	13.5
Mbarara	62	44
Others	38	27
<b>Occupation</b>		
Business	33	23.4
Farmer	38	27
Housewife	15	10.6
Others	42	29.8
Retired	13	9.2
<b>Monthly income</b>		
0-500000 UGX	78	55.3
500000-1M UGX	45	31.9
1MUGX or more	18	12.8
<b>Marital status</b>		
Divorced	8	5.7
Married	104	73.8

Other	20	14.2
Single	9	6.4
<b>Duration of hypertension</b>		
1-5 years	54	38.3
More than 6 years	67	47.5
less than one year	20	14.2
<b>Level of education</b>		
No formal education	34	24.1
Completed Primary	31	22
Completed Secondary	28	19.9
Completed Tertiary	48	34
<b>Heard of physical activity</b>		
No	20	14.2
Yes	121	85.8
<b>From whom</b>		
Friends and family	25	17.7
Health worker	79	56
Mass media	17	12.1
Never heard of physical activity	20	14.2
<b>Total</b>		
All	141	100

**Table 2:** Extent of physical activity among the participants.

Domains of physical activity	Frequencies	%
<b>Job-related physical activity</b>		
Low	130	92.2
Moderate	10	7.1
High	1	0.7
<b>Transportation physical activity</b>		
Low	131	92.9
Moderate	10	7.1
High	0	0
<b>Housework and house maintenance</b>		
Low	124	87.9
Moderate	16	11.4
High	1	0.7

Recreation, sport and leisure-time physical activity		
Low	125	88.7
Moderate	16	11.4
High	0	0
Overall physical activity score		
Low	77	54.6
Moderate	57	40.4
High	7	5
Total		
All	141	100

Table 3: Extent of physical activity and sociodemographic characteristics.

Characteristic	Frequencies			Total	%			Total	p-value
	Physical activity score				Physical activity score				
	Low	Moderate	High	All	Low	Moderate	High	All	
<b>Age group</b>									
less than 35 years	4	1	0	5	80	20	0	100	0.549
35-49 years	16	8	2	26	61.5	30.8	7.7	100	
50 years and above	57	48	5	110	51.8	43.6	4.5	100	
<b>Gender</b>									
Female	50	42	3	95	52.6	44.2	3.2	100	0.211
Male	27	15	4	46	58.7	32.6	8.7	100	
<b>Place of residence</b>									
Greater Bushenyi	7	15	0	22	31.8	68.2	0	100	0.016
Isingiro	15	3	1	19	78.9	15.8	5.3	100	
Mbarara	31	26	5	62	50	41.9	8.1	100	
Others	24	13	1	38	63.2	34.2	2.6	100	
<b>Occupation</b>									
Business	18	12	3	33	54.5	36.4	9.1	100	0.705
Farmer	20	17	1	38	52.6	44.7	2.6	100	
Housewife	9	6	0	15	60	40	0	100	
Others	25	14	3	42	59.5	33.3	7.1	100	
Retired	5	8	0	13	38.5	61.5	0	100	
<b>Monthly income</b>									
0-500000 UGX	46	31	1	78	59	39.7	1.3	100	0.166
500000-1M UGX	22	19	4	45	48.9	42.2	8.9	100	
1M UGX or more	9	7	2	18	50	38.9	11.1	100	
<b>Marital status</b>									
Divorced	2	6	0	8	25	75	0	100	0.376
Married	58	39	7	104	55.8	37.5	6.7	100	
Single	4	5	0	9	44.4	55.6	0	100	
Other	13	7	0	20	65	35	0	100	

Duration of hypertension								
less than one year	11	8	1	20	55	40	5	100
1-5 years	30	20	4	54	55.6	37	7.4	100
More than 6 years	36	29	2	67	53.7	43.3	3	100
Level of education								
No formal education	24	10	0	34	70.6	29.4	0	100
Completed Primary	21	10	0	31	67.7	32.3	0	100
Completed Secondary	12	14	2	28	42.9	50	7.1	100
Completed Tertiary	20	23	5	48	41.7	47.9	10.4	100
Heard of physical activity								
No	16	3	1	20	80	15	5	100
Yes	61	54	6	121	50.4	44.6	5	100
From whom								
Friends and family	11	14	0	25	44	56	0	25
Health worker	42	31	6	79	53.2	39.2	7.6	79
Mass media	8	9	0	17	47.1	52.9	0	17
Never heard of physical activity	16	3	1	20	80	15	5	20
Total All	77	57	7	141	54.6	40.4	5	

Note: Figures in the parentheses are column relative frequency; Others for place of residence include districts of Rukungiri, Ntungamo, Kiruhura, Kitagwenda; Others for occupation include teacher, health worker, office workers, security guards; The p-values were reported from the Fischer's exact test. Kabale, Kasese, Rwampara, Kamwenge, Ibanda.

Table 4: Significant associated factors to physical activity on bivariate and multivariate adjustment.

Characteristic	Bivariate analysis		Multivariate analysis	
	crude OR (cOR), (95% CI)	p-value	adjusted OR (aOR), (95% CI)	p-value
Place of residence				
Isingiro	-	-	-	-
Greater Bushenyi	0.964 (0.751-1.238)	0.773	-	-
Mbarara	-	-	-	-
Others	-	-	-	-
Level of education				
No formal education	-	-	-	-
Completed Primary	-	-	-	-
Completed	1.420 (1.106-1.824)	0.006	1.374 (1.055-1.790)	0.018
Secondary	-	-	-	-
Completed Tertiary	-	-	-	-
Heard of physical activity				
No	3.934 (1.243-12.452)	0.02	3.262 (0.953-11.166)	0.06
Yes	-	-	-	-
Sedentary behaviour				
More than 6 hours	0.262 (0.123-0.557)	0.001	0.276 (0.126-0.606)	0.001
Barrier				
Social influence	0.970 (0.851-1.106)	0.651	-	-

Notes: Others for place of residence include districts of Rukungiri, Ntungamo, Kiruhura, Kitagwenda, Kabale, Kasese, Rwampara, Kamwenge, Ibanda.

This study sought to establish the extent of physical activity and associated factors among patients attending hypertension clinics in Mbarara City, Southwestern Uganda. Physical activity is an effective way of managing hypertension. A participant is physically active if they achieve  $\geq 600$  METs in a period of 5 days or more per week. Therefore, an individual who had moderate or high levels of physical activity was considered to be physically active.

This study found that only 45.39% of the participants were physically active. These findings are in agreement with a study among hypertensive patients attending hospitals in the Central Gander Zone in Addis Ababa who found that (45%) of the participants were physically active [21]. These findings suggest that the majority of hypertensives are physically inactive. Therefore, interventions to increase physical activity among hypertensives should be emphasized. Few participants were active for transport compared to other domains. Creation of low traffic neighborhoods can improve walking and cycling as forms of active transport [22]. Therefore, policy formulation to include pedestrian and bicycle lanes during road infrastructure planning may be the solution to solve this transportation physical activity challenge.

Findings of this study reveal that participants who had higher levels of monthly income had high physical activity levels. Also found that participants with higher income levels were more physically active [23]. Poverty is more likely to be linked with poor health outcomes, even on the aspect of engaging in physical activity. Therefore, poverty eradication strategies and campaigns can be equally used as platforms for physical activity awareness and counselling.

This study found that those who had information about physical activity were more physically active and contrary to many studies, the study found that those who heard of physical activity from family and friends were more physically active compared to those who heard from health workers therefore health workers may need to assess available family support so as to achieve and sustain physical activity.

36.17% of participants were sedentary (spent more than 6 hours of daily sitting). Almost half of the participants who were physically inactive were also sedentary, reported similar findings [24]. This is a double disaster for people with hypertension [25]. Therefore, interventions to improve levels of physical activity should be put in place so as to address this issue.

The current study found level of education was associated with physical activity. Participants who had higher levels of education were more physically active. These results agree with a study by which reported that participants with hypertension who had completed high school and higher levels of education, were more physically active [26]. It has been noted that more educated participants are more aware of the normal values of blood pressure and their significance hence they partake in physical activity. It was also found that, with high education levels, one can easily obtain information, practice counselling, organise and carry out physical activities as per their schedule [21,27,28]. Higher levels of education have also been associated with better

economic outcomes, which consequently influence physical activity positively [29]. Contrasting results where participants with low levels of education were more physically active have been reported. This was explained in relation to the kind of jobs those with low levels of education engage in that is; more strenuous or physically demanding jobs [16,30].

Approximately, half of the physically inactive participants were also sedentary in the current study. A decrease in physical activity alongside an increase in sedentary behaviour has been reported in other studies [24,31,32]. A study among hypertensive subjects reported that each one hour of sedentary behaviour was associated with 0.06 mmHg increase in systolic blood pressure and 0.20 mmHg or diastolic blood pressure [8]. Further noted that higher periods of sedentary behaviour during television viewing are associated with higher risk of death among hypertensive individuals [33]. Therefore, being inactive and sedentary is a double disaster for ill health especially among people with chronic diseases such as hypertension, hence interventions to improve levels of physical activity should be put in place [24]

## CONCLUSION

Our study revealed that majority of hypertensive patients was physically inactive despite physical activity benefits. Almost half of the hypertensive patients were both physically inactive and sedentary, each of which has an independent negative effect on blood pressure control. Therefore, programs or interventions informing of the benefits of physical activity in management of hypertension should be greatly emphasized in the local settings. We recommend that, urban planners and ministry of transport provide for pedestrian and cyclists' lanes during road construction, and for ministry of health to establish cardiac rehabilitation programs in all government hospitals for all patients with hypertension and other cardiovascular diseases.

## STUDY LIMITATIONS

This study used subjective study tools therefore the results were subject to recall bias. The study tool adopted only assessed physical activity of individuals aged 18-69 years only therefore our results cannot be used to inform possible interventions and recommendations for improving physical activity among patients younger than 18 years and older than 69 years of age. The study sample size was small which makes our results unrepresentative to the general population.

## ACKNOWLEDGMENT

We want to acknowledge all those who have made this research project successful. We first thank God. Secondly, we thank our parents who have offered the necessary support. We also want to acknowledge our supervisor Dr. Arubaku Wilfred who guided us throughout the project. Many thanks to the MUST Research Ethics Committee for granting approval of our proposal, DMA Diagnostics and Laboratory Limited and Mbarara Medical Specialists Clinic for granting us opportunities to undertake research at their clinics. We also acknowledge Ms Eleanor Turyakira and Dr. Muwanguzi Moses for the support rendered to us during the data analysis process. Further thanks to the participants who participated in this study.

## DISCLOSURE

As the authors of this article, we disclose that there is no conflict of interest regarding this research.

## SOURCE OF FUNDING

The authors individually contributed to raising the funds that were used to conduct the study.

## REFERENCES

- Katzmarzyk PT, Friedenreich C, Shiroma EJ, Lee IM. Physical inactivity and non-communicable disease burden in low-income, middle-income and high-income countries. *Br J Sports Med.* 2022;56(2):101-106.
- Paffenbarger RS, Thorne MC, Wing AL. Chronic disease in former college students. VIII. Characteristics in youth predisposing to hypertension in later years. *American Am J Epidemiol.* 1968;88(1):25-32.
- Pinto A, di Raimondo D, Tuttolomondo A, Fernandez P, Arna V, Licata G. Twenty-four hour ambulatory blood pressure monitoring to evaluate effects on blood pressure of physical activity in hypertensive patients. *Clin J Sport Med.* 2006;16(3):238-243.
- Diaz KM, Shimbo D. Physical activity and the prevention of hypertension. *Curr Hypertens Rep.* 2013;15:659-668.
- Genin PM, Dessenne P, Finaud J, Pereira B, Dutheil F, Thivel D, et al. Effect of work-related sedentary time on overall health profile in active vs. inactive office workers. *Front Public Health.* 2018;6:279.
- Menotti A, Puddu PE, Lanti M, Maiani G, Catasta G, Alberti Fidanza A. Lifestyle habits and mortality from all and specific causes of death: 40-year follow-up in the Italian Rural Areas of the Seven Countries Study. *J Nutr Health Aging.* 2014;18:314-321.
- Lee PH, Wong FK. The association between time spent in sedentary behaviors and blood pressure: A systematic review and meta-analysis. *Sports Med.* 2015;45:867-880.
- Yu L, Liang Q, Zhou W, Huang X, Hu L, You C, et al. Sedentary behavior and the risk of cardiac-cerebral vascular diseases in southern China. *Medicine.* 2018;97(44).
- World Health Organization. Global action plan for the prevention and control of noncommunicable diseases 2013-2020. 2013.
- World Health Organization. Global status report on noncommunicable diseases 2014. 2014.
- Rodgers WM, Hall CR, Duncan LR, Pearson E, Milne MI. Becoming a regular exerciser: Examining change in behavioural regulations among exercise initiates. *Psychol Sport Exerc.* 2010;11(5):378-386.
- Duncan LR, Hall CR, Wilson PM, Jenny O. Exercise motivation: A cross-sectional analysis examining its relationships with frequency, intensity, and duration of exercise. *International Int J Behav Nutr Phys Act.* 2010;7:1-9.
- Lee GY, Kim KO, Ryu JH, Park SH, Chung HR, Butler M. Exploring perceived barriers to physical activity in Korean older patients with hypertension: Photovoice inquiry. *Int J Environ Res Public Health.* 2022;19(21):14020.
- Westertep KR. Assessment of physical activity: A critical appraisal. *Eur J Appl Physiol.* 2009;105:823-828.
- Mbijiwe J, Chege P, Munyaka A. Assessment of physical activity level and its effects on blood pressure control among hypertensive patients attending Kiambu Level Five Hospital, Kenya. *Int J Health Sci Res.* 9(8):406-410.
- Umuvandimwe B. Factors associated with participation in physical activity among adults with hypertension in Kigali, Rwanda.
- Kalima NA, Gabriel BK, Muhindo R, Muyingo A. Chronic kidney disease in patients admitted to the medical ward of Mbarara Regional Referral Hospital in southwestern Uganda: prevalence and associated factors. *IJMBR.* 2015;4(2):107-116.
- Bukenya D, van Hout MC, Shayo EH, Kitabye I, Junior BM, Kasidi JR, et al. Integrated healthcare services for HIV, diabetes mellitus and hypertension in selected health facilities in Kampala and Wakiso districts, Uganda: A qualitative methods study. *PLOS Glob Public Health.* 2022;2(2):e0000084.
- Okello S, Muhithi A, Mohamed SF, Ameh S, Ochimana C, Oluwasanu AO, et al. Hypertension prevalence, awareness, treatment, and control and predicted 10-year CVD risk: A cross-sectional study of seven Communities in East and West Africa (SevenCEWA). *BMC Public Health.* 2020;20:1-3.
- Olum S, Muyingo A, Wilson TL, Demaerschalk BM, Hoxworth JM, Zhang N, et al. Stroke mortality outcomes in Uganda. *J Stroke Cerebrovasc Dis.* 2021;30(5):105661.
- Smachew M, Melak MF, Atenafu A, Belew AK. Lifestyle modification practice and associated factors among diagnosed hypertensive patients in selected hospitals in central gondar zone. *Nutr Metab Insights.* 2022;15:11786388221088245.
- Gamage AU, Seneviratne RD. Physical inactivity, and its association with hypertension among employees in the district of Colombo. *BMC Public Health.* 2021;21(1):1-1.
- da Silva BP, Tosi MM, dos Santos NQ, de Oliveira DV, Fidelix YL, do Nascimento Júnior JR, et al. Nivel de atividade física de idosos hipertensos e sua associação com dados sociodemográficos e condições de saúde. *Revista Pesquisa em Fisioterapia.* 2022;12:e4398.
- Lavie CJ, Ozemek C, Kachur S. Promoting physical activity in primary and secondary prevention. *Eur Heart J.* 2019;40(43):3556-3558.
- di Pietro L, Al-Ansari SS, Biddle SJ, Borodulin K, Bull FC, Buman MP, et al. Advancing the global physical activity agenda: recommendations for future research by the 2020 WHO physical activity and sedentary behavior guidelines development group. *Int J Behav Nutr Phys Act.* 2020;17(1):1-1.
- Churilla JR, Ford ES. Comparing physical activity patterns of hypertensive and nonhypertensive US adults. *Am J Hypertens.* 2010;23(9):987-993.
- Rizvi F, Afzal M, Chaudhry MA, Baig A. Impact of education and physical activity on awareness and control of hypertension.
- Boshara R. Thrivers and Strugglers: A Growing Economic Divide. Bridges. 2015.
- Cavagnoli D. The labour supply curve: A pluralist approach to investigate its measurements. *Econ Labour Relat Rev.* 2012;23(3):71-87.
- NICE BG. Four commonly used methods to increase physical activity: brief interventions in primary care, exercise referral schemes, pedometers and community-based exercise programmes for walking and cycling. National Institute for Clinical Excellence. 2006.

31. Healy GN, Wijndaele K, Dunstan DW, Shaw JE, Salmon J, Zimmet PZ, et al. Objectively measured sedentary time, physical activity, and metabolic risk: the Australian Diabetes, Obesity and Lifestyle Study (AusDiab). *Diabetes Care*. 2008;31(2):369-371.
32. Du Y, Liu B, Sun Y, Snetselaar LG, Wallace RB, Bao W. Trends in adherence to the physical activity guidelines for Americans for aerobic activity and time spent on sedentary behavior among US adults, 2007 to 2016. *JAMA Netw Open*. 2019;2(7):e197597.
33. Lemes IR, Sui X, Fernandes RA, Blair SN, Turi-Lynch BC, Codogno JS, et al. Association of sedentary behavior and metabolic syndrome. *Public Health*. 2019;167:96-102.