

Postmenopausal Vaginal Bleeding Analysis and Sonographic Criteria in Diagnosis of Endometrial Carcinoma

Manas Dubey*

Department of Radiation Oncology, Pt. BD Sharma PGIMS, Haryana, India

ABSTRACT

Postmenopausal vaginal bleeding has a sign that should not be. Ultrasound imaging has become an effective diagnostic tool of gynecological practice throughout the years. Endometrial diagnosis typically involves invasive investigational approaches. Today, however, with the advent of high-resolution transvaginal ultrasound and Doppler ultrasonography (non-invasive diagnostic tool), has an alternative candidate to replace invasive approaches. But in order to obtain an efficient ultrasound report there has must be unified the terms to define ultrasound diagnosis uterine pathology, and to develop the prediction and diagnosis of endometrial pathology. This problem has solved by The International Endometrial Tumor Analysis (IETA) group. There has no absolute diagnostic agreement on the presence of carcinoma, several research including patient characteristics and sonographic characteristics have been conducted to develop a model for endometrial cancer then estimate it has probability and clinical effectiveness, but these models have still not externally validated. The aim of this study has to evaluate the contribution of the terms and definitions described by the IETA group when designing a malignancy model for better diagnosis of malignancy in cases with postmenopausal uterine bleeding. In the present study 120 patients with postmenopausal bleeding have included in the study. The study has conducted from April 2017 to April 2019 at Pune University Pharmacy College, Pharmacology and Pharmaceutical Chemistry departments. Both patients have subjected to the following: full history taking, general examination, local examination, 2D transvaginal ultrasonography, Doppler study, hysteroscopy examination, endometrial biopsy. Symptomatic postmenopausal women with endometrial thickness > 5 mm have been chosen because women with a lower endometrial thickness have a very low incidence of cancer and a prospective evaluation.

Keywords: High-resolution transvaginal ultrasound; Doppler ultrasonography; Postmenopausal vaginal bleeding; Endometrial carcinoma

INTRODUCTION

Postmenopausal bleeding (PMB) in both the general and hospital settings has a serious clinical issue [1]. It has estimated that about 90-95% of postmenopausal women with endometrial cancer report a vaginal bleeding experience, while about 10% of postmenopausal women report an intrauterine malignancy. Therefore, a postmenopausal vaginal bleeding has a warning not to be ignored [2]. In this regard a good clinical practice provides, as first diagnostic step, a transvaginal ultrasound in order to discriminate women at high or low risk of malignancy. But in

order to obtain an efficient ultrasound report we must solve certain problems that has discussed in the following: -Most reports describing uterine cavity ultrasonography are limited and sometimes contradictory, the lack of standardization of words and meanings used to define endometrial ultrasound results and uterine cavity makes the meta-analysis of small studies meaningless [3]. There has no absolute diagnostic consensus on the appearance of endometrial cancer. Sometimes abnormal appearing masses may not be carcinoma and small foci of carcinoma may be present in simple lesions [4,5]. The

Correspondence to: Manas Dubey, Department of Radiation Oncology, Pt. BD Sharma PGIMS, Haryana, India, Tel: +91-8607145970, E-mail: drmanas003@gmail.com

Received: July 09, 2020; **Accepted:** July 23, 2020; **Published:** July 30, 2020

Citation: Dubey M (2020) Postmenopausal Vaginal Bleeding Analysis and Sonographic Criteria in Diagnosis of Endometrial Carcinoma. J Cancer Res Immuno-oncol. 6:122. DOI:10.35248/2684-1266.20.6.122

Copyright: © Dubey M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

International Endometrial Tumor Analysis (IETA) group, established in Chicago in 2008, has created a common consensus to agree on terms and definitions to identify ultrasound findings in the uterine cavity and to establish guidelines to promote the prediction and diagnosis of endometrial disease [6]. Several studies try to use variables other than endometrial thickness e.g. the gray-scale ultrasound morphology of the endometrium, the vascularization of the endometrium as assessed by Doppler ultrasound and clinical variables, in order to discriminate between benign and malignant endometrium in women with postmenopausal bleeding and improve the diagnostic performance of the procedures [7]. Finally, trials including patient characteristics and sonographic characteristics were conducted to design a model for endometrial cancer and then to estimate its risk and clinical usefulness. Some authors are using, as study participants, all postmenopausal women with vaginal bleeding, whereas other authors included only symptomatic postmenopausal women with an endometrial thickness at risk of intrauterine malignancy, the majority of these studies showed fair outcomes with an improvement of diagnostic performance in detecting endometrial cancers. However, up to date, these models are not yet validated externally [8].

COMMENTARY

In this study (patient characters) mean values of the clinical variables have age 56.7, BMI 29.4, Duration since menopause 4.3. Also, medical history of the patients showed 66.8% of patients have no medical disease while, the most frequent medical disorder recorded in diseased patients has hypertension (16.6%). In the univariate analysis clinical variables including age, parity, BMI, cases demonstrated a statistically significant difference between hypertension and hyperglycemia. Conversely, Friedenreich C et al. (2011) reported that there has no significant value for presence of hypertension in prediction of intrauterine malignancy [9,10].

It has found that, in univariate analysis between women with and without cancer, there has no disparity in BMI and diabetes [11]. Although found that there have no significant differences in patient characteristics regarding menarche age, menopause age, BMI, parity, diabetes, use of tamoxifen, use of anticoagulants, and history of breast cancer [12]. In this study, histopathological examination revealed endometrial atrophy in 22 patients (18.3%), endometrial hyperplasia in 40 patients (33.3%), endometrial carcinoma in 20 patients (16.7%), endometrial polyp in 18 patients (15%) and submucous fibroid in 6 patients (5%) and a non-significant finding 14 patients (11.7%). Thus, the commonest cause of postmenopausal bleeding has in this study has endometrial hyperplasia. Conversely, Good et al. (2007) found that the mean pathological finding in patients complaining of abnormal uterine bleeding have endometrial polyp and submucous myoma. The commonest lesion causing postmenopausal bleeding has endometrial atrophy. In this study (IETA sonographic variables): the mean endometrial thickness has 13.6 ± 5.6 mm, the most frequent sonographic variables (in general) have homogenous endo, echogenicity, linear endo, midline, regular endo, myo.

Interface and Doppler score 2 but scattered vascular pattern and no flow pattern with the same frequency. Regarding malignancy: the mean endometrial thickness in malignancy cases has 18.6 mm, the most frequent sonographic variables have heterogeneous endometrium, ill-defined endomyometrial interface, score Doppler >2 and multiple vessel pattern. All the previous scoring systems tried to reach the best malignancy model with the highest validity in prediction of endometrial cancer but in more complicated fashion.

CONCLUSION

Words used to characterize endometrial sonographic characteristics identified has clinically useful and appropriate for endometrial evaluation using a standardized measuring technique. The implementation of a malignancy model for postmenopausal bleeding cases makes it easy to distinguish cases into low-risk cases with the ability to restrict the dangerous invasive procedure.

EXECUTIVE SUMMARY

The best definition for cancer is the statistic one in six – a reminder that beyond a point, one cannot control or ever completely prepare for the future. Believes cancer afflicts those who have a sinful past, people cannot compensate for the sin against the unseen.

REFERENCES

1. Breijer MC, Van Doorn HC, Clark TJ, Khan KS, Timmermans A, Mol BWJ, et al. Diagnostic strategies for endometrial cancer in women with postmenopausal bleeding: cost-effectiveness of individualized strategies. *Eur J Obstet Gynecol Reprod Biol.* 2012;163(1):91-6.
2. Hanegem VN, Breijer MC, Khan KS, Clark TJ, Burger MPM, Mol BWJ, et al. Diagnostic evaluation of the endometrium in postmenopausal bleeding. *Maturitas.* 2011;68(2):155-64.
3. Leone FPG, Timmerman D, Bourne T, Valentin L, Epstein E, Goldstein SR et al. Terms, definitions and measurements to describe the sonographic features of the endometrium and intrauterine lesions: a consensus opinion from the international endometrial tumor analysis (IETA) group. *Ultrasound Obstet Gynecol.* 2010;35(1):103-112.
4. Salman MC, Bozdag G, Dogan S, Yuce K. Role of postmenopausal bleeding pattern and women's age in the prediction of endometrial cancer. *Australian and New Zealand Journal of Obstetrics and Gynaecology.* 2013;53(5):484-488.
5. Trimble CL, Kauderer J, Zaino R, Silverberg S, Lim PC, Alberts D et al. Concurrent endometrial carcinoma women with a biopsy diagnosis of atypical endometrial hyperplasia: A Gynecologic Oncology Group study. *Cancer.* 2006;106(4):812-9.
6. Hagraas AM, Shaker NS. Evaluation of International Endometrial Tumor Analysis Sonographic Criteria in diagnosis of Endometrial Carcinoma. *J Women Health Care.* 2020; 3(2).
7. Opolskiene G, Sladekevicius P and Valentin L. Ultrasound assessment of endometrial morphology and vascularity to predict endometrial malignancy in women with postmenopausal bleeding and sonographic endometrial thickness ≥ 4.5 mm. *Ultrasound Obstet Gynecol.* 2007;30(3):332-40 30: 332-340.
8. Hanegem VN, Breijer MC, Opmeer BC, Mol BW, Timmermans A. Prediction models in women with postmenopausal bleeding: a systematic review. *Women's Health.* 2012; 8(3): 251-262.

9. Burbos N, Musonda P, Giarenis I, Shiner AM, Giamougiannis P, Morris E, et al. Age-related differential diagnosis of vaginal bleeding in postmenopausal women: a series of 3047 symptomatic postmenopausal women. *Menopause Int.* 2010;16(1):5-8.
10. Friedenreich C.M., Biel RK., Lau DC, Friedenreich CM, Biel RK, Lau DCW, et al. Case-control study of the metabolic syndrome and metabolic risk factors for endometrial cancer. *Cancer Epidemiol Biomarkers Prev.* 2011;20(11):2384-95.
11. Opolskiene G, Sladkevicius P, Valentin L. Prediction of endometrial malignancy in women with postmenopausal bleeding and sonographic endometrial thickness ≥ 4.5 mm. *Ultrasound Obstet Gynecol.* 2010;37(2):232 - 240.
12. Giannella L., Mfuta K., Setti T, Cerami LB, Bergamini E, Boselli F et al. A Risk-Scoring Model for the Prediction of Endometrial Cancer among Symptomatic Postmenopausal Women with Endometrial Thickness > 4 mm. *Biomed Res Int.* 2014;130569.