

**Review Article** 

**Open Access** 

# Potent Role of Antioxidant Molecules in Prevention and Management of Skin Cancer

Wahida Khan Chowdhury<sup>1</sup>, Shahida Arbee<sup>2</sup>, Sujan Debnath<sup>3</sup>, Shah Mehedi Bin Zahur<sup>4</sup>, Sharmim Akter<sup>5</sup>, A K M Rezaul Karim<sup>6</sup>, Mohammad Mohabbulla Mohib <sup>7,8</sup>, Abida Tisha<sup>8</sup>, Md Abu Taher Sagor<sup>8\*</sup> and Sarif Mohiuddin<sup>2</sup>

<sup>1</sup>Department of Dermatology, Shahabuddin Medical College and Hospital, Bangladesh

<sup>2</sup>Department of Anatomy and Physiology, Pioneer Dental College and Hospital, Bangladesh

<sup>3</sup>Department of Dental Public Health, Pioneer Dental College and Hospital, Bangladesh

<sup>4</sup>Infectious Disease Division, (icddr,b)s, Bangladesh

<sup>5</sup>Department of Medicine, Dhaka Central International Medical College and Hospital, Bangladesh

<sup>6</sup>Department of Pediatric Hematology and Oncology, Chittagong Medical College and Hospital, Chittagong-4203, Bangladesh

<sup>7</sup>Research Institute for Medicines (iMed.ULisboa) Faculty of Pharmacy, Universidade de Lisboa, Portugal

<sup>8</sup>Department of Pharmaceutical Sciences, School of Life Sciences, North South University, Bangladesh

\*Corresponding author: Md Abu Taher Sagor, Department of Pharmaceutical Sciences, North South University, Dhaka-1229, Bangladesh, Tel: 8801718130130; E-mail: sagor2008nsu@gmail.com

Received date: April 22, 2017; Accepted date: May 09, 2017; Published date: May 15, 2017

Copyright: © 2017 Chowdhury WK, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

#### Abstract

Evidence reported that cancers are spreading every nook and corner of the world at an alarming rate. Skin diseases like chronic skin inflammations, psoriasis and skin cancers have also been burning topic in today. Protections of a biological system are often hampered while skin gets damaged. Factors like UV, radiation, viruses, chronic diseases, genetic predispositions, food habits and environmental exposures might lead to skin cancers. In addition to these, urbanization and globalization may also contaminate the environment that may eventually modify several biological and genetic functions. USA, Europe and Australia are in the most dangerous zone to be exposed. We basically performed detailed search of PubMed, Google Scholar and Science Direct for literature search and collecting related information. On the other hand, experiments suggested that antioxidant components such as phenolic acid derivatives, flavonoids and flavonol found to be preventive against cancer cell proliferations. Moreover, antioxidants have been also evaluated as a protective agent against chronic inflammatory diseases as well. These molecules may participate as an additional therapy which could exert synergistic effects while applying with other chemotherapeutic agents. Our literature findings and hypothetical figure may establish a good correlation between skin cancer and antioxidant therapy. Therefore this study will be focused on skin cancer biology and some possible management strategies using antioxidant phyto-nutrients.

**Keywords:** Skin; Radiation; Cancer; Chemotherapy; Free radicals; Antioxidants

#### Introduction

Skin is the most outer part of the body which also known as the largest organ of the body. It protects internal organs, heals automatically and lasts a lifetime [1]. However, mammalian skin is composed of generally epidermis and dermis. Skin helps in several communications like absorption, excretion, heat regulation and water resistance. In addition, these layers prevent and protects from injuries, pathogens, heat and other harmful exposures [2,3].

According to National Center for Health Statistics, around 1,685,210 new cancer population and 595,690 cancer deaths are projected to occur in United States during 2016 only [4]. Currently, cancers have been a real burden and the cost of this life threatening situations has become unattainable for the lower and middle class people; as a result, affected subjects lose their property as well as savings. Similarly, it is also a major challenge to the health care providers to ensure the effective treatment for everyone though researchers are working very hard to discover competent remedies to minimize this concern [5]. However, chronic exposures, human papiloma virus, UV-radiation, and dietary intake have been identified as the major culprit of skin cancer. Several reports found that using moisturizing cream may increase the risk of developing skin cancers through UV [6,7]. Melanoma, squamous-cell skin cancer and basal-cell skin cancer are observed mostly in today's medical investigations. Basal cells constantly divide to form daughter cells to replace the squamous cells. This type of cancer is observed through skin cells in the lowest layer of the epidermis which is the most seen skin carcinoma [8]. Squamouscell skin cancers are the second most prominent owing to UV exposures; hence, clinical appearance is highly variable and often has the tendency to spread towards other organs [9]. On the other hand, excess sun burn in the childhood and adulthood may cause Melanoma [10] which further linked with melanoma once there is any mutations in the germ line p16 are occurred [11]. Several sign and symptoms have been reported on these such as burning, itching, moderate to severe pain, ulcer, edema, redness and often bleeding [12].

Free radical often reacts with cell membrane, endoplasmic reticulum, necessary enzymes and other genetic materials result in cellular necrosis. In addition, necrotized cells often secret various potent pro-inflammatory markers like Tumor necrosis factors,

activator proteins, Toll like receptors and interferons [13,14]. Several investigations reported that oxygen mediated free radicals often react with DNA and increase the chance of cancer [15,16]. On top of that, oxidative stress is considered as one of the key regulators in skin carcinogenesis, and thereby identifying nontoxic strong antioxidants to diminish skin cancer is an important area of investigation. Antioxidants strongly prevent both photo-carcinogenesis and skin tumor progression in animal model, in part, by reducing free radicals and reactive oxygen species. These molecules have also been proven highly potent by inhibiting endogenous TNF-a in animal skin, a prime mediator in skin tumor initiator. Furthermore, evidences often suggested that antioxidant inhibit mitogenic and cell survival signaling and initiate apoptosis for cell suppression. Finally, phenolic acid derivatives and flavonoids effectively modulate cell-cycle regulators and check points toward reserve of proliferation, and often arrest growth in G0-G1 phases of a cell cycle [17-19]. Therefore, this study will describe the probable molecular reasons for developing skin cancer along with that a series of therapeutic approaches based on antioxidant molecules will be exposed to prevent skin carcinoma.

## **Skin Cancer**

Cancer, one of the most complicated system in a biological subject which deals with several factors such as alteration of DNA bases, mutation in genetic materials, mutation or blocking of tumor suppressive gene p53 [20,21]. As a result, normal cell of the body lost their control and start replicating their progenitor cells for an enormous numbers which finally act as immortal cells [22]. Every cell has some predetermined functions and they remain in one place for their own survival and biological importance but in cancer, these cells have the ability to travel or spread toward other organs and take control over there. These cells then become malignant tumor and may be threat for a subject [23]. In today's world, men are more prone to prostate cancer, and women are mostly vulnerable towards breast and ovarian cancers but it is skin cancer which affects both the genders [24]. Merkel cell carcinoma, Kaposi sarcoma, Cutaneous (skin) lymphoma, Skin adnexal tumors and different types of sarcomas are rarely observed on skin [25,26]. The most common reason for skin cancer is either sunburn or excess UV exposure, however, family history also plays a major role [27]. Several reports have been suggested where excess UV can cause skin carcinoma in the laboratory animals [28,29]. So far UVA, UVB and UVC have been investigated where UVA and UVB found to have mostly responsible for skin carcinoma, on the contrary, UVC can't reach to the earth due to protection of Ozone layer. Extreme Sun light/UV may penetrate to the skin nucleus and alter/damage genetic code which further leads to abnormal cell replication [30,31]. Often, the Sun penetrates skin cells deeply that these cells are to be damaged. In the affected areas, chemokines and harmful cytokines are accumulated resulting inflammation that may eventually trigger skin cancers [32]. Environmental toxins like smoking, high oxygen atmosphere, household spills and chemical exposures may contain several free radicals and once they enter inside a biological subject, they produce ionization radiation [33]. Inside the body, once ionizing radiation hits a molecule in any cell, an electron may be replaced which ultimately lead to the formation of a highly reactive free radical. The generation of unwanted high levels of free radicals is the mechanism by which ionizing radiation alter necessary proteins, genetic codes and leads to the necrosis of a cell [34]. Skin has been a major target of oxidative stress due to highly reactive free radicals that originate in the

Page 2 of 7

environment and in the skin itself that further is responsible for the development of skin carcinoma [35].

## Limitations of Chemotherapies

The prevalence of cancer in USA and other western countries is very high and often found tumor who lead normal life [36]. At the initial stage of the tumor progression the treatment on any cancer is easier and to some extend very much curable, however once it reaches towards the metastasis it become more complex and it most cases the treatment seems to be more resistant and the chances of cure become very narrow. Currently, three types of treatment protocol are now being used in the clinic to treat cancer. Radiation which kills individual cells, full or partial removal of tumor cells by surgery and chemotherapies which work against particular cells [37]. In contrast, the treatment of cancer is becoming very difficult owing to following different patterns by the tumor cell. Tumor cells are very distinct and always vary within individuals [38,39]. In addition, it is really difficult to deal with cancer subjects because cancer is a constellation of at least 200 other diseases and it has been often reported that cancer cells are not homogenize [24]. It is always suggested that cancer therapies should optimize most tumor cells and it must be monitored for minimal effect to the normal cells [40]. Unfortunately, all cancer therapies produce severe side effects, adverse effects and are very costly. In majority if the cases, single drug therapy is not sufficient to treat carcinoma. Multiple therapies with several cycles are necessary to optimize the cancer cells. Alopecia, bone marrow depression, severe vomiting, fatigue, rash and heart failure are associated with chemotherapies [41,42]. Most of the anti-tumor drugs are not well studied and many of these are not approved by FDA. Anti-cancer molecules are often contraindicated during pregnancy and several studies suggested that these drugs may interact with foetus and may cause death [43,44]. Besides, several deaths have been reported while given chemotherapies on many clinical evaluations [45,46]. Hepatic and kidney failure have been documented when platinum based chemotherapy was provided [47,48]. Cardiac arrest or arrhythmia have been documented when monoclonal antibodies were administrated to the cancer subjects and sometimes the subjects must be admitted in ICU during prior treatment which increase overall cost [49-51].

# An Overview of Antioxidants

Oxidative stress mostly found to be free radical-mediated which further correlates with the response of cellular metabolism and catabolism [52]. In addition, free radicals are mainly generated from mitochondria during ATP production and cellular response in a biological subject. Sometimes drug molecules produce free radicals when given in higher dosage [53]. There are numerous evidences where free radical-mediated oxidative stress linked with many diseases like cardiovascular dysfunctions [54], hepatic damages [55], renal injury [56], diabetes [57], neuro-degeneration [58], aging [59] and other life threatening events [60]. Free radicals are often blamed for acute and chronic inflammation [61] which further drawn with cancers [62,63]. Several approaches against cancer cells have been undertaken targeting reactive oxygen species and often found good results while taken with other synthetic drugs [64]. On the other hand, antioxidants are the molecules which fight against oxidative stress. These are also known as 'free radical scavengers'. Basically, polyphenols and flavonoids are considered as anti-oxidants which exert several protective mechanisms and found to be effective among in vivo, in vitro and human subjects [65]. Foods and supplements based

#### Page 3 of 7

antioxidants have been proved beneficial against free radical mediated damages, at the same time they protect nucleus and necessary enzymes [57,66]. There are multiple studies have been undertaken to investigate anti-cancer activities of antioxidant molecules and some breakthrough outcomes were successfully documented [67,68].

## Role of Antioxidant on Skin Cancers

Skin, the biggest organ of the body is protected by several layers. Multiple protective mechanisms have been proposed so far through animal and human studies. Chronic UV exposure-induced skin diseases are mainly caused by the excessive oxidative stress that further leads to inflammation and finally results in DNA and necessary protein damages [35]. DNA protection using antioxidants therapy has been a good choice for clinical trials. Phenolic acid and flavonoids are being reported extremely well against chemoprevention [69]. It is reported that in the TPA-induced skin inflammation model citrus peel extract prevented expression of inducible nitric oxide synthase, cyclooxygenase-2, ornithine decarboxylase, and vascular endothelial growth factor in mouse skin. The study also noticed inhibitory effects of citrus on 7,12-dimethylbenez [a] anthracene (DMBA)/TPA-induced skin tumour formation and reduced tumor incidence, tumor weight and tumor multiplicity of papillomas at 20 weeks treatment [70]. The blocking properties of naringin were observed when in BALB/c mice skin irradiated with UVB. The expressions of interleukin-1 $\beta$  (IL-1 $\beta$ ), interleukin-6 (IL-6), interleukin-8 (IL-8) and cyclooxygenase-2 (COX-2) were significantly lowered in Ha CaT cells by primarily blocking MAPK pathways [71]. Src family kinases (SFKs), basically known as oncogenes, are recently drawing attention for their association in melanoma skin cancers; on the other hand, naringin may be an effective molecule for the treatment of melanoma as a natural inhibitor of c-Src [72]. Retinoids such as 13-cis-retinoic acid and retinyl palmitate exhibit a good preventive effect in chemically induced papillomas and carcinomas of the skin in female Swiss mice after three weeks of induction of cancer. The study showed inhibitory effects of retinoids on skin carcinoma [73]. Topical application of vitamin E in a dose-dependent manner has been shown to reduce the incidence of UV-induced skin cancer in mice by exhibiting antioxidant mechanism. Study shows that dimer and trimer products may contribute in avoidance of UV-induced photo-damage in animal [74]. Vitamin C was also found as photo-protectant while applied to mice and pig skin before exposure to ultraviolet radiation (Figure 1) [75,76].



A potent antioxidant molecule, Coenzyme Q 10, generally exerts several mechanisms in cancer, inflammation and aging. When Coenzyme Q10 applied on human dermal and epidermal cells, it protects against cell death induced by reactive oxygen species in keratinocytes [77]. A population-based case-control study was performed to investigate the correlation between citrus peel use and black tea intake on squamous cell carcinoma of the skin tissue. Preventive effects of citrus peel and black tea were statistically noticed on human subjects [78]. A study was undertaken with a large number of healthy women whom solar stimulated UV were given on the skin and  $\beta$ -carotene (120 mg) was provided. The study found to be

protective by inducing skin lycopene and reducing oxidative markers [79].

The hypothetical pathway explains that there are various extracellular effects can activate the signaling of the growth factor. One of the major reasons of skin cancer is the UV-B from UV radiation. UV-B can influence the mutation of K-RAS which is a upstream signaling kinase of MAPK family. Even though K-RAS signaling is essential for normal cell proliferation and apoptosis but when mutated it is a key player of activating certain signaling pathway that leads to the binding of different transcription factor with DNA that will cause the activation of different cytokines which will participate in the

## Page 4 of 7

progression of cancer cell. It has been most commonly known that chronic inflammation leads to the progression of cancer. In skin cancer many studies suggested that activation of TNFR1 by its binding with TNF- $\alpha$  facilitate the synthesis of different inflammatory cytokines. All these factors together participate in the progression of tumor cell. White people are more vulnerable towards the skin cancer than the black and mixed race people. One of the possible reason is the production of melanin is really low in the skin of white people compared to black. However, certain polyphenols and flavons can

prevent the mutation from taking place. The underlining reason is yet unknown, due to the lack of study in this particular area it is yet to be discover, but one hypothesis is the hyper activation of the growth factor can be prevented by polyphenols and flavons. They also prevent the phosphorylation of certain kinases, which leads to the apoptosis of those cells. All these activity made them a potential molecule against skin cancer (Table 1).

Subjects	Outcomes of the study	References
Model: Solar simulated irradiation of skin Disease induced by: Radiation Treatment: Ferulic acid Dose: 0.5% Ferulic acid	Caspase 3, 7 density were lowered	[80]
Model: Human keratinocytes Disease induced by: Se methyl selenocysteine Treatment: Cuso <sub>4</sub> Dose: N/A	Decreased DNA synthesis, cell growth inhibition, DNA synthesis and Blockade of the cell cycle at the S/G2-M phase and cell death by necrosis.	[81]
Model: CD-1 mice Disease induced by: Genetic Treatment: Lupeol Dose: 1–2 mg/mouse	Decreased protein expression of ODC, cox-2, nitric oxide synthase, and blocked NF-κB and phosphatidyl inositol 3-kinase (PI3K)/Akt signaling in tumor promotion	[82]
Model: Heterozygous DBA/2 MnSOD KO mice Disease induced by: 12-O-tetradecanoylphorbol-13- acetate Treatment: SOD mimetic (MnTE-2-PyP5+) Dose: 5 ng/mouse	suppressed mitosis without interfering with apoptosis, decreased protein carbonyls and reduced the activity of activator protein-1, and prevented the level of proliferating cellular nuclear antigen	[83]
Model: Human Disease induced by: UV light irradiation Treatment: Lutein and zeaxanthin Dose: Lutein 5 mg/zeaxanthin 0.3 mg 2/day	Changed in lipid peroxidation and photo-protective activity in the skin.	[84]
Model: Mouse Disease induced by: Solar ultraviolet (UV) radiation Treatment: (-)-Epigallocatechin-3-gallate (EGCG) Dose: 1 mg/cm <sup>2</sup> skin area	inhibited UVB-induced oxidative stress, significant prevention of UVB-induced depletion of antioxidant enzymes, and Inhibition of a single UVB irradiation-induced phosphorylation of ERK1/2 and MAPK family.	[7]
Model: Skh:2 Hairless Mice Disease induced by: Ultraviolet Irradiation Treatment: Vitamin E Dose: 5% oral and topical RRR-α-tocopheryl acetate	showed no signs of toxicity and had significantly less acute and chronic skin damage induced by UV irradiation, and reduced inflammation and pigmentation.	[29]
Model: Human Disease induced by: Acute UV radiation Treatment: Antioxidant complex – vitamins (lycopene, β-carotene, α-tocopherol), selenium Dose: 3 mg of tomato lycopene, 3 mg of natural α- and b-carotene 5 mg of natural α-tocopherol and 37.5 mg of organic selenium	reduction of the UV-induced erythemas, and Parallel reduction of the lipoperoxide level was observed.	[85]

Citation: Chowdhury WA, Arbee S, Debnath S, Bin Zahur SM, Akter S, et al. (2017) Potent Role of Antioxidant Molecules in Prevention and Management of Skin Cancer. J Clin Exp Dermatol Res 8: 393. doi:10.4172/2155-9554.1000393

Model: Mice Disease induced by: UV-induced skin cancer Treatment: Vitamin E and epigallocatechin gallate Dose: Olive oil on the dorsal skin with a moist cotton swab	Delayed the onset of and to reduce the incidence rate of skin cancer development.	[28]
Model: SK-mel28 and A375 human melanoma cells Disease induced by: Genetically induced Treatment: Resveratrol Dose: 30, 60, or 100 mM all-trans-resveratrol	Induced phosphorylation of ERK1/2 in A375	[86]
Model: Healthy women Disease induced by: Solar stimulated light Treatment: β-carotene Dose: 120 mg	Skin lycopene found to be increased and reduced oxidative damages.	[79]
Model: Human Disease induced by: UV light Treatment: Carotenoids and carotenoids plus vitamin E Dose: 25 mg total carotenoids/d and 335 mg (500 IU) RRR-α-tocopherol/d	Dorsal black skin was significantly diminished	[87]
Model: Cell culture Disease induced by: UV or ozone Treatment: Vitamin E Dose: N/A	Combated oxidative stress induced by UV	[88]
Model: Hairless mice Disease induced by: UV exposure Treatment: β-carotene Dose: 10% carotenoids	Reduced skin tumor progression	[89]

 Table 1: Role of antioxidant molecules on various cancer induced skin cells.

# **Conclusion and Future Directions**

Evidences suggested that oxidative stress plays a pivotal role in mutation of the different genes, and the expression of many cytokines and chemokines leads to the progression of tumor. It also prevents mitochondrial biogenesis as well, resulting cellular reproduction gets hampered; on the other hand, antioxidants found to be very useful for biogenesis. Although antioxidant molecules have been highly appreciated in the treatment of skin diseases like skin cancers, negative findings are reported too. Both animal and small clinical trials are showing good activities when antioxidant molecules are applied on various skin cancers. Additional larger and wide randomized clinical trials are needed to provide clear scientific reports about the potential benefits of taking antioxidant supplements during skin cancer treatment.

## Funding

This work was not funded directly or indirectly from any organization or institution.

# **Conflict of Interest**

The authors declare no conflict of interest.

#### References

- Wahida Khan C, AbidaTisha, Sharmim A, Shah Mehedi Bin Z, Nahid H, et al. (2017) The Role of Arsenic on Skin Diseases, Hair Fall and Inflammation: An Immunological Review and Case Studies. J Clin Exp Dermatol Res 8: 2.
- de Golian E, Kwong BY, Swetter SM, Pugliese SB (2016) Cutaneous Complications of Targeted Melanoma Therapy. Current treatment options in oncology 17: 57.
- 3. Leung DY (2013) New insights into atopic dermatitis: role of skin barrier and immune dysregulation. Allergol Int 62: 151-161.
- 4. Siegel RL, Miller KD, Jemal A (2016) Cancer statistics, 2016. CA: A Cancer. J Clin 66: 7-30.
- Gordon LG, Brynes J, Baade PD, Neale RE, Whiteman DC, et al. (2017) Cost-effectiveness analysis of a skin awareness intervention for early detection of skin cancer targeting men over 50 years. Value Health 20: 593-601.
- 6. Lu YP, Lou YR, Xie JG, Peng Q, Shih WJ, et al. (2009) Tumorigenic effect of some commonly used moisturizing creams when applied topically to UVB-pretreated high-risk mice. J Invest Dermatol 129: 468-475.
- Vayalil PK, Elmets CA, Katiyar SK (2003) Treatment of green tea polyphenols in hydrophilic cream prevents UVB-induced oxidation of lipids and proteins, depletion of antioxidant enzymes and phosphorylation of MAPK proteins in SKH-1 hairless mouse skin. Carcinogenesis 24: 927-936.

Page 6 of 7

- 8. Saleh GM, Desai P, Collin JR, Ives A (2017) Incidence of eyelid basal cell carcinoma in England: 2000-2010. Br J Ophthalmol 101: 209-212.
- Latil M, Nassar D, Beck B, Boumahdi S, Wang L, et al. (2017) Cell-Type-Specific Chromatin States Differentially Prime Squamous Cell Carcinoma Tumor-Initiating Cells for Epithelial to Mesenchymal Transition. Cell stem cell 20: 191-204. e195.
- Autier P, Epimel D, Eortc (1998) Melanoma Cooperative Group JF: Influence of sun exposures during childhood and during adulthood on melanoma risk. Int J Cancer 77: 533-537.
- Hussussian CJ, Struewing JP, Goldstein AM, Higgins PA, Ally DS, et al. (1994) Germline p16 mutations in familial melanoma. Nat Genet 8: 15-21.
- 12. Esteva A, Kuprel B, Novoa RA, Ko J (2017) Dermatologist-level classification of skin cancer with deep neural networks. Nature. 542: 115-118.
- 13. Halliwell B, Gutteridge JM (2015) Free radicals in biology and medicine. 5th edition, Oxford University Press, USA.
- Reuter S, Gupta SC, Chaturvedi MM, Aggarwal BB (2010) Oxidative stress, inflammation, and cancer: how are they linked? Free Radic Biol Med 49: 1603-1616.
- Valko M, Izakovic M, Mazur M, Rhodes CJ, Telser J, et al. (2004) Role of oxygen radicals in DNA damage and cancer incidence. Mol Cell Biochem 266: 37-56.
- Pryor WA (1982) Free radical biology: xenobiotics, cancer, and aging. Ann N Y Acad Sci 393: 1-22.
- 17. Singh RP, Agarwal R (2002) Flavonoid antioxidant silymarin and skin cancer. Antioxid Redox Signal 4: 655-663.
- Zi X, Agarwal R (1999) Modulation of mitogen-activated protein kinase activation and cell cycle regulators by the potent skin cancer preventive agent silymarin. Biochemical and biophysical research communications 263: 528-536.
- 19. Katiyar SK (2002) Treatment of silymarin, a plant flavonoid, prevents ultraviolet light-induced immune suppression and oxidative stress in mouse skin. Int J Oncol 21: 1213-1222.
- 20. Errington TM, Iorns E, Gunn W, Tan FE, Lomax J, et al. (2014) An open investigation of the reproducibility of cancer biology research. Elife 10: 3.
- 21. Vander Heiden MG, DeBerardinis RJ (2017) Understanding the Intersections between Metabolism and Cancer Biology. Cell 168: 657-669.
- 22. Weinberg R (2013) The biology of cancer. 2nd edition, Garland science.
- 23. Greenberg ES, Chong KK, Huynh KT, Tanaka R, Hoon DS, et al. (2014) Epigenetic biomarkers in skin cancer. Cancer Lett 342: 170-177.
- 24. Ferlay J, Soerjomataram I, Dikshit R, Eser S, Mathers C, et al. (2015) Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. Int J Cancer 136: E359-E386.
- 25. Johnson MC, Myers AN (2017) Cytology of Skin Neoplasms. Vet Clin North Am Small Anim Pract 47: 85-110.
- 26. Agbai ON, Buster K, Sanchez M, Hernandez C, Kundu RV, et al. (2014) Skin cancer and photoprotection in people of color: a review and recommendations for physicians and the public. J Am Acad Dermatol 70: 748-762.
- 27. Stang A, Jöckel KH (2016) Does skin cancer screening save lives? A detailed analysis of mortality time trends in Schleswig-Holstein and Germany. Cancer 122: 432-437.
- Ichihashi M, Ahmed NU, Budiyanto A, Wu A, Bito T, et al. (2000) Preventive effect of antioxidant on ultraviolet-induced skin cancer in mice. J Dermatol Sci 23 Suppl 1: S45-50.
- Burke KE, Clive J, Combs GF Jr, Commisso J, Keen CL, et al. (2000) Effects of topical and oral vitamin E on pigmentation and skin cancer induced by ultraviolet irradiation in Skh:2 hairless mice. Nutr Cancer 38: 87-97.
- 30. Torres A (2017) Response to 'Avoidance of sun exposure as a risk factor for major causes of death: a competing risk analysis of the Melanoma in Southern Sweden cohort'. Journal of internal medicine 281: 217-218.

- Hault K, Rönsch H, Beissert S, Knuschke P, Bauer A (2016) Knowledge of outdoor workers on the effects of natural UV radiation and methods of protection against exposure. J Eur Acad Dermatol Venereol 30: 34-37.
- 32. Lund AW, Medler TR, Leachman SA, Coussens LM (2016) Lymphatic Vessels, Inflammation, and Immunity in Skin Cancer. Cancer Discov 6: 22-35.
- 33. Morgan WF (2003) Non-targeted and delayed effects of exposure to ionizing radiation: II. Radiation-induced genomic instability and bystander effects in vivo, clastogenic factors and transgenerational effects. Radiat Res 159: 581-596.
- Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, et al. (2007) Free radicals and antioxidants in normal physiological functions and human disease. Int J Biochem Cell Biol 39: 44-84.
- 35. Trouba KJ, Hamadeh HK, Amin RP, Germolec DR (2002) Oxidative stress and its role in skin disease. Antioxid Redox Signal 4: 665-673.
- 36. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet Tieulent J, et al. (2015) Global cancer statistics, 2012. CA Cancer J Clin 65: 87-108.
- Lalla RV, Bowen J, Barasch A, Elting L, Epstein J, et al. (2014) MASCC/ ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. Cancer 120: 1453-61
- Hosoya N, Miyagawa K (2014)Targeting DNA damage response in cancer therapy. Cancer Sci 105: 370-388.
- Suda K, Mitsudomi T (2014) Successes and limitations of targeted cancer therapy in lung cancer. In: Successes and Limitations of Targeted Cancer Therapy. Prog Tumor Res 41: 62-77.
- 40. Qiu H, Fang X, Luo Q, Ouyang G (2015) Cancer stem cells: a potential target for cancer therapy. Cell Mol Life Sci 72: 3411-3424.
- Burstein HJ (2000) Side effects of chemotherapy. Journal of Clinical Oncology 18: 693-693.
- 42. de Boer Dennert M, De Wit R, Schmitz P, Djontono J, v Beurden V, et al. (1997) Patient perceptions of the side-effects of chemotherapy: the influence of 5HT3 antagonists. Br J Cancer 76: 1055.
- **43**. Lynch CD, Lee MJ, Del Priore G (2013) Chemotherapy in Pregnancy. Clinical Pharmacology During Pregnancy. 201.
- Koren G, Carey N, Maxwell C, Nulman I, Gagnon R, et al. (2013) Cancer chemotherapy and pregnancy. Journal of Obstetrics and Gynaecology Canada 35: 263-278.
- 45. Leduc C, Quoix E (2017) Programmed death of chemotherapy in nonsmall-cell lung cancer? The Lancet. 389: 227-228.
- Vacchelli E, Aranda F, Eggermont A, Galon J, Sautès-Fridman C, et al. (2014) Trial Watch: Chemotherapy with immunogenic cell death inducers. Oncoimmunology 3: e27878.
- 47. Chu E, DeVita Jr VT, DeVita Jr VT (2016) Physicians' Cancer Chemotherapy Drug Manual Jones & Bartlett Publishers.
- 48. Group IPFS (2016) Prognostic factors in patients with metastatic germ cell tumors who experienced treatment failure with cisplatin-based first-line chemotherapy. J Clin Oncol 28: 4906-11.
- Guglin M, Aljayeh M, Saiyad S, Ali R, Curtis AB, et al. (2009) Introducing a new entity: chemotherapy-induced arrhythmia. Europace 11: 1579-1586.
- Chaudhary P, Gajra A (2010) Cardiovascular effects of EGFR (epidermal growth factor receptor) monoclonal antibodies. Cardiovasc Hematol Agents Med Chem 8: 156-163.
- 51. Hasselbalch B, Lassen U, Hansen S, Holmberg M, Sørensen M, et al. Cetuximab, bevacizumab, and irinotecan for patients with primary glioblastoma and progression after radiation therapy and temozolomide: a phase II trial. Neuro Oncol 12: 508-16.
- 52. Chowdhury MRH, Sagor MAT, Tabassum N, Potol MA, Hossain H, et al. (2015) Supplementation of Citrus maxima peel powder prevented oxidative stress, fibrosis, and hepatic damage in carbon tetrachloride (CCl4) treated rats. Evidence-Based Complementary and Alternative Medicine 598179.
- 53. Sagor MAT, Mohib M, Tabassum N, Ahmed I, Reza H, et al. (2016) Fresh seed supplementation of syzygium cumini attenuated oxidative stress,

inflammation, fibrosis, iron overload, hepatic dysfunction and renal injury in acetaminophen induced rats. J Drug Metab Toxicol 7: 2.

- 54. Sagor MAT, Tabassum N, Potol MA, Alam MA (2015) Xanthine oxidase inhibitor, allopurinol, prevented oxidative stress, fibrosis, and myocardial damage in isoproterenol induced aged rats. Oxidative medicine and cellular longevity. 478039.
- 55. Sagor AT, Chowdhury MRH, Tabassum N, Hossain H, Rahman MM, et al. (2015) Supplementation of fresh ucche (Momordica charantia L. var. muricata Willd) prevented oxidative stress, fibrosis and hepatic damage in CCl 4 treated rats. BMC complementary and alternative medicine 15: 115.
- 56. Alam MA, Chowdhury MRH, Jain P, Sagor MAT, Reza HM, et al. (2015) DPP-4 inhibitor sitagliptin prevents inflammation and oxidative stress of heart and kidney in two kidney and one clip (2K1C) rats. Diabetol Metab Syndr 7: 107.
- Mohib MM, Rabby SF, Paran TZ, Hasan MM, Ahmed I, et al. (2016) Protective role of green tea on diabetic nephropathy—A review. Cogent Biology 2: 1248166
- 58. Al-Amin MM, Akhter S, Hasan AT, Alam T, Hasan SN, et al. (2015) The antioxidant effect of astaxanthin is higher in young mice than aged: a region specific study on brain. Metab Brain Dis 30: 1237-1246.
- 59. Abu Taher Sagor M (2016) Angiotensin-II, a potent peptide, participates in the development of hepatic dysfunctions. Immunology ,Endocrine & Metabolic Agents in Medicinal Chemistry 16: 1-17.
- 60. Mohib MM, Hasan I, Chowdhury WK, Chowdhury NU, Mohiuddin S, et al. (2016) Role of angiotensin II in hepatic inflammation through MAPK pathway: a review. J Hep 2: 2
- 61. Abu Taher Sagor M, Mahmud Reza H, Tabassum N, Sikder B, Ulla A (2016) Supplementation of Rosemary Leaves (Rosmarinus officinalis) Powder Attenuates Oxidative Stress, Inflammation and Fibrosis in Carbon Tetrachloride (CCl4) Treated Rats. Current Nutrition & Food Science 12: 288-295.
- 62. Valko M, Rhodes C, Moncol J, Izakovic M, Mazur M, et al. (2006) Free radicals, metals and antioxidants in oxidative stress-induced cancer. Chem Biol Interact 160: 1-40.
- 63. Dreher D, Junod AF (1996) Role of oxygen free radicals in cancer development. Eur J Cancer 32:30-38.
- 64. Trachootham D, Alexandre J, Huang P (2009) Targeting cancer cells by ROS-mediated mechanisms: a radical therapeutic approach? Nature reviews Drug discovery 8: 579-591.
- 65. Diplock A, Charuleux JL, Crozier-Willi G, Kok F, Rice-Evans C, et al. (1998) Functional food science and defence against reactive oxidative species. Br J Nutr 80: S77-S112.
- 66. Abu Taher S, Hasan Mahmud R, Nabila T, Biswajit S, Anayt U, et al. (2016) Supplementation of rosemary leaves (Rosmarinus officinalis) powder attenuates oxidative stress, inflammation and fibrosis in carbon tetrachloride (CCl4) treated rats. Current Nutrition & Food Science 12:1-8.
- 67. Kornhauser A, Lambert LA, Wamer WG, Wei RR, Lavu S, et al. (1995) Antioxidants and cancer prevention. In: Nutrients in Cancer Prevention and Treatment. edn. Springer 83-100.
- Huang MT, Ho CT, Lee CY (1992) Phenolic compounds in food and their effects on health II: antioxidants and cancer prevention. ACS Publications 507.
- 69. Nichols JA, Katiyar SK (2010) Skin photoprotection by natural polyphenols: anti-inflammatory, antioxidant and DNA repair mechanisms. Arch Dermatol Res 302: 71-83.
- Pan MH, Li S, Lai CS, Miyauchi Y, Suzawa M (2012) Inhibition of citrus flavonoids on 12-O-tetradecanoylphorbol 13-acetate-induced skin inflammation and tumorigenesis in mice. Food Science and Human Wellness 1:65-73.

 Ren X, Shi Y, Zhao D, Xu M, Li X (2016) Naringin protects ultraviolet Binduced skin damage by regulating p38 MAPK signal pathway. J Dermatol Sci 82: 106-114.

Page 7 of 7

- Guo B, Zhang Y, Hui Q, Wang H, Tao K (2016) Naringin suppresses the metabolism of A375 cells by inhibiting the phosphorylation of c-Src. Tumour Biol 37: 3841-3850.
- 73. Abdel Galil A, Wrba H, El Mofty M (1984) Prevention of 3methylcholanthrene-induced skin tumors in mice by simultaneous application of 13-cis-retinoic acid and retinyl palmitate (vitamin A palmitate). Experimental pathology 25: 97-102.
- 74. Krol E, Kramer Stickland KA, Liebler DC(2000) Photoprotective actions of topically applied vitamin E. Drug Metab Rev 32: 413-420.
- 75. Pandel R, Poljšak B, Godic A, Dahmane R (2013) Skin photoaging and the role of antioxidants in its prevention. ISRN dermatology. 930164.
- 76. Elmore A (2004) Final report of the safety assessment of L-Ascorbic Acid, Calcium Ascorbate, Magnesium Ascorbate, Magnesium Ascorbyl Phosphate, Sodium Ascorbate, and Sodium Ascorbyl Phosphate as used in cosmetics. Int J Toxicol 24: 51-111.
- 77. Muta Takada K, Terada T, Yamanishi H, Ashida Y, Inomata S, et al. (2009) Coenzyme Q10 protects against oxidative stressinduced cell death and enhances the synthesis of basement membrane components in dermal and epidermal cells. Biofactors 35: 435-441.
- 78. Hakim IA, Harris RB (2001) Joint effects of citrus peel use and black tea intake on the risk of squamous cell carcinoma of the skin. BMC dermatology 1: 3.
- Ribaya Mercado JD, Garmyn M, Gilchrest BA, Russell RM (1995) Skin lycopene is destroyed preferentially over beta-carotene during ultraviolet irradiation in humans. J Nutr 125: 1854.
- Lin FH, Lin JY, Gupta RD, Tournas JA, Burch JA, et al. (2005) Ferulic acid stabilizes a solution of vitamins C and E and doubles its photoprotection of skin. J Invest Dermatol 125: 826-832.
- 81. Tapiero H, Townsend D, Tew K (2003) The antioxidant role of selenium and seleno-compounds. Biomed Pharmacother 57: 134-144.
- Saleem M, Afaq F, Adhami VM, Mukhtar H (2004) Lupeol modulates NF-κB and PI3K/Akt pathways and inhibits skin cancer in CD-1 mice. Oncogene 23: 5203-5214.
- Zhao Y, Chaiswing L, Oberley TD, Batinic Haberle I, Clair WS, et al. (2005) A mechanism-based antioxidant approach for the reduction of skin carcinogenesis. Cancer Res 65: 1401-1405.
- 84. Palombo P, Fabrizi G, Ruocco V, Ruocco E, Fluhr J, et al. (2007) Beneficial long-term effects of combined oral/topical antioxidant treatment with the carotenoids lutein and zeaxanthin on human skin: a double-blind, placebo-controlled study. Skin Pharmacol Physiol 20: 199-210.
- Cesarini J, Michel L, Maurette J, Adhoute H, Bejot M, et al. (2003) Immediate effects of UV radiation on the skin: modification by an antioxidant complex containing carotenoids. Photodermatol Photoimmunol Photomed 19: 182-189.
- Niles RM, McFarland M, Weimer MB, Redkar A, Fu YM, et al. (2003) Resveratrol is a potent inducer of apoptosis in human melanoma cells. Cancer Lett 190: 157-163.
- 87. Stahl W, Heinrich U, Jungmann H, Sies H, Tronnier H(2000) Carotenoids and carotenoids plus vitamin E protect against ultraviolet light–induced erythema in humans. Am J Clin Nutr 71: 795-798.
- Packer L, Weber SU, Rimbach G (2001) Molecular aspects of αtocotrienol antioxidant action and cell signalling. J Nutr 131: 369S-373S.
- Mathews Roth MM, Krinsky NI (1987) Carotenoids Affect Development of UVB Induced Skin Cancer. Photochem Photobiol 46: 507-509.