

# Pregnancy Nutritional Status in Ethiopia: A case study of Pregnant Women in Shashemene District, Southern Oromia Region Ethiopia

Yoseph Gela\*

Addis Ababa University Main campus college of Development study, Addis Ababa, Ethiopia

**Keywords:** Nutritional Status, Pregnant Women, Protein, Vitamins, Energy, Southern Oromia Region, Ethiopia

## OPINION

Inadequate quality and quantity diet is one of the major reasons for high levels of malnutrition in pregnant women [1]. Across-sectional survey was conducted in Shashemene District, Southern Oromia Region [2]. A two-stage cluster sampling technique was used to select a representative sample of 15190 pregnant women aged 20-50 years from four rural villages Energy and nutrient intakes from foods were calculated from one-day weighed food records on a sub-sample (n = 83) [3]. The result of the study showed that the intakes of most nutrients were lower than the recommended intake. The energy intake of the study participants both in 2nd and 3rd trimesters of pregnancy were 2,308 kcal and 1,420.5 kcal compared to the recommended 2,340 kcal and 2,452 kcal, respectively [4].

Except iron, almost all micronutrient intakes were lower than the recommended intake. Vitamin A intake was 3 µg compared with the recommended 800 µg, while protein intake of the study respondents in 2nd and 3rd trimester of pregnancy was 45.9 g and 31.5 g, respectively, compared with the recommended 71 g. Risk factors for under nutrition were multiple pregnancy and no consumption of cereal-based foods [5]. This study revealed that the energy and nutrient intake of the pregnant women in study area was below the recommended intakes [6]. Furthermore, the situation might be aggravated by high phytate content food consumption reported [7,8].

Nutritional status of pregnant women in study area was not adequate to support the increased energy and nutrient requirement of the participants [9].

## REFERENCES

1. Brown, L. S. 2011. Nutrition Requirement during Pregnancy. 2015.
2. Okwu GN, Ukoha AI, Nwachukwu N, Agha NC. Studies on the predisposing factors of protein energy malnutrition among pregnant women in a Nigerian community. *Online J of Health and Allied Sci.* 2008;6(3).
3. West Jr KP. Extent of vitamin A deficiency among preschool children and women of reproductive age. *J Nutr.* 2002;132(9):2857S-66S.
4. Teller CH, Yimer G. Levels and determinants of malnutrition in adolescent and adult women in southern Ethiopia. *Ethiopian J Health Develop.* 2000;14(1):57-66.
5. Karim R, Bhat D, Troy L, Lamstein S, Levinson FJ. Determinants of food consumption during pregnancy in rural Bangladesh: examination of evaluative data from the Bangladesh Integrated Nutrition Project. *Friedman School of Nutri Sci and Policy*; 2002.
6. <https://ideas.repec.org/p/fsn/wpaper/11.html>. 2014.
7. Ali YG. Pregnancy Nutritional Status in Ethiopia. *J Food Technol Nutri Sci. SRC/JFTN/108.* 2020.
8. Ethiopian Health and Nutrition Research Institute (EHNRI). Food Composition Table for Use in Ethiopia. Addis Ababa: EHNRI, 34 (199 Ethiopia. 8).
9. Umata M, West CE, Fufa H. Content of zinc, iron, calcium and their absorption inhibitors in foods commonly consumed in *J Food Composition Anal.* 2005;18(8):803-17.

\*Correspondence to: Yoseph Gela, Addis Ababa University Main campus college of Development study, Addis Ababa, Ethiopia, Tel: 251913372078; E-mail: yodapartner@gmail.com

Received: April 18, 2021; Accepted: May 11, 2021; Published: May 18, 2021

Citation: Yoseph Gela (2021) Pregnancy Nutritional Status in Ethiopia: A case study of Pregnant Women in Shashemene District, Southern Oromia Region Ethiopia. *J Women's Health Care* 10:529. doi:10.35248/2167-0420.21.10.529.

Copyright: © 2021 Gela Y, et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.