

Prevalence of Complications Associated with Alzheimer Disease patients in Aseer Region, Saudi Arabia

Adel Ali Alhazzani^{1*}, Mohammed Saeed Alqahtani², Ahmed A. Awwadh³, Turki Ali Alyami⁴, Mohammad Saad Alshomrani⁵, Mushary Saeed Alqahtani⁶

¹Department of Medicine, College of Medicine, King Saud university, Riyadh, Saudi Arabia;

²Neurology Resident, Armed Forces Hospital–Southern Region/King Fahad Hospital-Jeddah, Saudi Arabia;

³Emergency Medicine Resident, Aseer Central Hospital, Abha, Saudi Arabia; ⁴Khamis Mushayt Maternity And Children Hospital, Saudi Arabia; ⁵Anesthesia Resident, King Sakman Armed Forces Hospital, Tabuk, Saudi Arabia; ⁶ Department of Medicine, College of Medicine, King Khalid University, Abha, Saudi Arabia

ABSTRACT

Background: Alzheimer's disease (AD) is neurodegenerative disorder that develops over a period of years that differs from normal aging. One of the most important risk factors is increasing age, and the majority of people with Alzheimer's are 65 years and older. AD reduces life expectancy and is one of the major causes of physical disability, institutionalization, and low quality of life among the elderly. AD is highly related to functional disability and institutionalization. There are many factors associated with AD including physical and behavioural complications.

Aim: To assess epidemiological pattern and complications of AD among patients in Aseer region, Saudi Arabia.

Methodology: A descriptive cross-sectional study included 110 Alzheimer patients (66 males and 44 females) registered at Aseer Central Hospital, Southern of Saudi Arabia. A pre-structured questionnaire was used for data collection that included patients' personal characteristics and frequency of exposure to complications associated with Alzheimer disease.

Results: About 72% of the patients aged 70 years or older and 60% were males. Almost all of the patients were citizens 97.3% and 62.7% were married. Exact of 56.4% of the patients were illiterate and only few 4.5% were university graduated. Pneumonia was the most frequent complication followed by getting lost, fall down, and bone fracture. Getting lost was significantly more among males than females ($P=0.007$), while pneumonia was significantly more among patients treated in governmental hospitals ($P=0.003$). On the other hand, bone fractures and falling down did not differ significantly according to patients' personal characteristics.

Conclusions: The most frequent complications associated with Alzheimer Disease in our study population were pneumonia, getting lost, falling down and bone fractures. Risk factors associated with these complications include male gender for getting lost. Health care providers are advised to provide close care to Alzheimer disease patients.

Keywords: Alzheimer's disease; Care givers; Complications; Epidemiology; Patient characteristics; Disease burden

INTRODUCTION

Alzheimer's disease (AD) is a type of dementia that causes decline in cognitive function and subsequently behavioural changes. [1] Symptoms usually develop slowly and progress over time; yet severe enough to interfere with daily tasks which differentiate it from normal aging. Alzheimer's is the most common type of dementia, accounts for 60 % to 80 % of dementia cases. [2, 3] One of the most important risk factors is increasing age, and the

majority of people with Alzheimer's are 65 years and older. [4] Nearly 200,000 Americans under the age of 65 have younger-onset Alzheimer's disease (also known as early-onset Alzheimer's). [5] Memory loss, in particular recent and short term memory, is the first sign which is mild, but as disease progress decline in other cognitive functions, communication, visuospatial orientation and ambulation [6] Alzheimer's is the sixth leading cause of death in the United States. On average, a person with Alzheimer's lives four to eight years after diagnosis, but can be as long as 20 years,

Correspondence to: Adel Alhazzani, Department of Medicine, College of Medicine, King Saud University, Riyadh, Saudi Arabia, Tel: +966 556644455; E-Mail: aalhazzani2@ksu.edu.sa

Received: February 19, 2021; **Accepted:** March 05, 2021; **Published:** March 12, 2021

Citation: Alhazzani AA, Alqahtani MA, Awwadh MA, Alyami TA, Alshomrani MS, Saeed MAlqahtani MS (2021) Prevalence of complications associated with Alzheimer Disease Patients in Aseer Region, Saudi Arabia. Int J Phys Med Rehabil. 9:592.

Copyright: © 2021 Adel AA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

depending on other factors. [7].

Regarding disease burden at the individual level, AD reduces life expectancy and is one of the major causes of physical disability, institutionalization, and low quality of life among the elderly. [8] AD is highly related to functional disability and institutionalization. It is reported that among individuals over 60 years of age dementia contributes 11.2 % of the years lived with disability, compared with 9.5 % for stroke, 8.9 % for musculoskeletal disorders, and 5.0 % for cardiovascular disease. [9].

Alzheimer's has no curative therapy, but therapeutic interventions can be considered to temporarily delay the progress of dementia symptoms and improve quality of life for those with Alzheimer's and their caregivers [10-12].

METHODOLOGY

A descriptive cross-sectional approach including 110 patients with Alzheimer's disease at Aseer Central Hospital during the period from January to July 2017 was conducted. Data were collected directly from the patients and caregivers using pre-structured questionnaire. Questionnaire was developed by the authors with the help of literature and experts consultation. Data collected included patients demographic data such as age, gender, education level, marital status, and relation of the caregiver with the patient. Data regarding disease were also extracted including disease duration, hospitalizations, and disease complications. Medical care setting (private or governmental) besides patients' satisfaction regarding the provided care were also covered in the study questionnaire.

Data analysis

After data were extracted, it was revised, coded and fed to statistical software IBM SPSS version 22(SPSS, Inc. Chicago, IL). All statistical analysis was done using two tailed test. P value less than 0.05 was considered to be statistically significant. Descriptive analysis based on frequency and percent distribution was done for all variables including patients' demographic data, medical care data, and disease complications. Univariate relations between patients' bio-demographic data and disease complications were assessed using exact probability test due to low expected frequencies in the cross tabulations.

RESULTS

A sample of 110 cases of AD was included in the study. About 72% of the patients aged 70 years or older and 60% were males. Almost all of the patients were Saudis 97.3% except to and 62.7% of them were married. As for educational level, 56.4% of the patients were illiterate and only 4.5% were university graduated. More than 50% of the cases had the disease for less than two years and 33.6% were diseased for 3-5 years. Regarding hospital admission, 30% of the patients were not admitted to any hospital during the last year while 50% were admitted 1-4 times. About 69% of the patients' caregivers were their siblings and only 11.8% were spouse.

Considering patients medical care, 73.6% of the patients had their medical care in governmental hospitals in Aseer region while 38.2% needed to receive their health services outside the Aseer region. As for patients' satisfaction regarding the provided health care, 53.6% of the patients were unsatisfied while 18.2% were satisfied regarding the health care they received.

Regarding the complications associated with AD among the pa-

tients, pneumonia was the most frequent complication (60.9%) followed by getting lost (60.9%), fall down (50.9%), and bone fracture (18.2%) (Figure 1).




Figure 1: AD consequences among patients in Aseer region, Saudi Arabia.

Demonstrates the relation the complications with the patients' characteristics. Pneumonia was recorded among 62.6% of the Saudi cases compared to none of the non-Saudis with recorded statistical significance ($P=.028$). Also 69.1% of the patients who received their medical care in the governmental hospitals had pneumonia compared to 37.9% of those who received the care in private hospitals ($P=.003$). As for satisfaction relation with pneumonia, it was recorded among 72.9% of the unsatisfied cases compared to 29% of patients with neutral attitude regarding the health care ($P=.001$). As for bone fracture correlates, it was recorded among 66.7% of the non-Saudi patients compared to 16.8% of Saudis with recorded statistical significance ($P=.027$). None of the other patients' characteristics were significantly related with fractures. Considering fall down, it was reported for 52.3% of the Saudi patients compared to none of the others ($P=.074$). Also fall down was recorded for 58.1% of the illiterate patients compared to 41.7% of educated group ($P=.088$). With consideration to getting lost, it was reported among 71.2% of the male patients compared to 45.5% of the females ($P=.007$). None of the other patients' characteristics were significantly related with getting lost.

DISCUSSION

Alzheimer's disease (AD) is a degenerative disease, which is characterized by being worse with time. Alzheimer's disease may begin 20 years or more before being symptomatic, [13-15] with small changes in the brain which usually patient unaware. Only after years of brain changes do individuals experience noticeable symptoms, such as memory loss and language problems. Symptoms occur because nerve cells (neurons) in parts of the brain involved in thinking, learning and memory (cognitive function) have been damaged or destroyed. [16] Over time, symptoms progress and affect individuals' ability to perform everyday activities. At this point, the individual is said to have dementia due to Alzheimer's disease, or Alzheimer's dementia. [17, 18].

The current study aimed to assess the epidemiology and consequences in AD among patients in Aseer region, Saudi Arabia. The study revealed that more than two thirds of the patients aged 70 years or more. Also about two thirds were males and married. Regarding disease duration of the included patients, the current study revealed that more than half of the cases had the disease for less than two years and nearly half of these patients were ad-

mitted to hospital one to four times during the last year. As for care giver, more than two thirds were patients siblings and only very few % (11%) were spouse. Alzheimer's Association in 2011 reported that over 5.4 million people in the United States have AD, including 5.2 million people aging 65 years or older. With the advanced ages, it is estimated that this number will increase by 50% with over 7.7 million people in that age range affected by AD by the year 2030, and will almost triple of 11 to 16 million by the year 2050. [19] Other studies revealed that the most significant risk factors for late-onset Alzheimer's are older age, carrying the e4 form of the APOE gene and having a family history of Alzheimer's. The vast majority of people with Alzheimer's dementia at age of 65 or older. Also other researchers confirmed that most affected patients were males. [20-22].

With regard to complications recorded among AD patients, the current study revealed that pneumonia was the most frequent complication which was recorded among nearly two thirds of the cases followed with getting lost and falling down but the least recorded was having bone fractures. Recurrent pneumonia explains the high admission rate to hospitals during the last year with some cases with fractures. The study also revealed that pneumonia was significantly associated with patient nationality as it was more recorded among Saudi patients because they are the main bulk of the sample. Also pneumonia was significantly more among patients who received the medical care in the governmental hospitals which may be explained by that the medical care of lower quality and some of the governmental hospitals don't have the trained staff to deal with AD cases. Also pneumonia was highly recorded among patient with poor satisfaction regarding the provided medical care (most probably those who were treated at governmental hospitals). Fall down was more recorded among illiterate AD patients who were cared by parents or paid persons. Getting lost was more among male patients who may be allowed to be outdoor due to gender nature making them more liable for getting lost.

CONCLUSION

In our study population of representing sample of Alzheimer Disease patients, most frequent complications was development of pneumonia which needed hospitalization followed by getting lost, falling down and bone fractures. Risk factors associated with these complications include male gender for getting lost, lower educational level and frequent hospital admissions for pneumonia. Health care providers are advised to be aware of the disease nature and the physiological and behavioural changes associated with the disease to be aware of all possible health related risks.

REFERENCES

- Hendrie HC. Epidemiology of dementia and Alzheimer's disease. *Am J Geriatr Psychiatry*. 1998; 6: S3-18.
- Sosa-Ortiz AL, Acosta-Castillo I, Prince MJ. Epidemiology of dementias and Alzheimer's disease. *Archives of medical research*. 2012; 43: 600-608.
- Scazufca M, Menezes PR, Vallada HP, Crepaldi AL, Pastor-Valero M, Coutinho LMS, et al. High prevalence of dementia among older adults from poor socioeconomic backgrounds in São Paulo, Brazil. *Int Psychogeriatr*. 2008; 20: 394-405.
- Plassman BL, Langa KM, Fisher GG, Heeringa SG, Weir DR, Ofstedal MB, et al. Prevalence of dementia in the United States: the aging, demographics, and memory study. *Neuroepidemiology*. 2007; 29: 125-132.
- Mattson MP. Pathways towards and away from Alzheimer's disease. *Nature*. 2004; 430: 631-639.
- Alzheimer's Association. 2016 Alzheimer's disease facts and figures. *Alzheimer's & Dementia*. 2016; 12: 459-509.
- Kahle-Wroblewski K, Ye W, Henley D, Hake AM, Siemers E, Chen YF, et al. Assessing quality of life in Alzheimer's disease: Implications for clinical trials. *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*. 2017; 6: 82-90.
- Shah H, Albanese E, Duggan C, Rudan I, Langa KM, Carrillo MC, et al. Research priorities to reduce the global burden of dementia by 2025. *The Lancet Neurology*. 2016; 15: 1285-1294.
- Schelterns P, Feldman H. Treatment of Alzheimer's disease; current status and new perspectives. *The Lancet Neurology*. 2003; 2: 539-547.
- Neugroschl J, Sano M. Current treatment and recent clinical research in Alzheimer's disease. *Mount Sinai Journal of Medicine: A Journal of Translational and Personalized Medicine: A Journal of Translational and Personalized Medicine*. 2010; 77: 03-16.
- Leon R, Garcia AG, Marco Contelles J. Recent advances in the multitarget directed ligands approach for the treatment of Alzheimer's disease. *Medicinal research reviews*. 2013; 33: 139-189.
- Villemagne VL, Burnham S, Bourgeat P, Brown B, Ellis KA, Salvado O, et al. Amyloid β deposition, neurodegeneration, and cognitive decline in sporadic Alzheimer's disease: A prospective cohort study. *Lancet Neurol*. 2013; 12: 357-367.
- Reiman EM, Quiroz YT, Fleisher AS, Chen K, Velez-Pardos C, Jimenez-Del-Rio M, et al. Brain imaging and fluid biomarker analysis in young adults at genetic risk for autosomal dominant Alzheimer's disease in the presenilin 1 E280A kindred: A case-control study. *Lancet Neurol*. 2012; 11: 1048-1056.
- Jack CR, Lowe VJ, Weigand SD, Wiste HJ, Senjem ML, Knopman DS, et al. Serial PiB and MRI in normal, mild cognitive impairment and Alzheimer's disease: Implications for sequence of pathological events in Alzheimer's disease. *Brain*. 2009; 132: 1355-1365.
- Bateman RJ, Xiong C, Benzinger TL, Fagan AM, Goate A, Fox NC, et al. Clinical and biomarker changes in dominantly inherited Alzheimer's disease. *N Engl J Med*. 2012; 367: 795-804.
- Gordon BA, Blazey TM, Su Y, Hari-Raj A, Dincer A, Flores S, et al. Spatial patterns of neuroimaging biomarker change in individuals from families with autosomal dominant Alzheimer's disease: A longitudinal study. *Lancet Neurol*. 2018; 17: 241-250.
- Braak H, Thal DR, Ghebremedhin E, Del Tredici K. Stages of the pathologic process in Alzheimer disease: age categories from 1 to 100 years. *J Neuropathol Exp Neurol*. 2011; 70: 960-969.
- Alzheimer's Association. 2011 Alzheimer's disease facts and figures. *Alzheimer's & Dementia*. 2011; 7: 208-244.

19. Hebert LE, Bienias JL, Aggarwal NT, Wilson RS, Bennett DA, Shah RC, et al. Change in risk of Alzheimer disease over time. *Neurology*. 2010; 75: 786-791.
20. Hebert LE, Weuve J, Scherr PA, Evans DA. Alzheimer disease in the United States (2010-2050) estimated using the 2010 Census. *Neurology*. 2013; 80: 1778-1783.
21. Saunders AM, Strittmatter WJ, Schmechel D, George-Hyslop PH, Pericak-Vance MA, Joo SH, et al. Association of apolipoprotein E allele epsilon 4 with late-onset familial and sporadic Alzheimer's disease. *Neurology*. 1993; 43: 1467-1472.
22. Farrer LA, Cupples LA, Haines JL, Hyman B, Kukull WA, Mayeux R, et al. Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease: A meta-analysis. *JAMA*. 1997; 278: 1349-1356.