

Proteomics Analysis for Amino Acid Misincorporation Detection: Mini Review

Joana Tavares, Filipe Assis-Santos and Manuel AS Santos*

Department of Medical Sciences, iBiMED- Institute of Biomedicine, University of Aveiro, 3910-193 Aveiro, Portugal

Abstract

Protein biosynthesis is a highly accurate biological process essential for life. Amino acid misincorporation errors (mistranslation) normally occur at low levels, but can increase sharply upon amino acid starvation, exposure to drugs, oxidative stress and other physiological perturbations. These processes disrupt protein function and are normally regarded as being deleterious, however, recent work has shown that they can also be regulated to produce advantageous phenotypes in both prokaryotes and eukaryotes. The biology of such unexpected adaptive mistranslation is poorly understood due to technical difficulties in the identification and quantification of amino acid misincorporations. In this mini-review, we describe proteome scale methodologies involving the use of mass-spectrometry and bioinformatics tools to directly detect and quantify mistranslation events and also indirect functional methods that permit sensitive, flexible and low-cost analysis of site specific amino acid variation.

Keywords: Amino acid misincorporation; Direct tools; Indirect tools; Mistranslation; Mass spectrometry; Reporter proteins

Introduction

The very high energetic and functional costs of protein synthesis require almost instantaneous tight regulation of its rate to guarantee cellular homeostasis during developmental programs, nutritional stress or other physiological perturbations. Fidelity of protein synthesis is equally important as it ensures functional activity and minimizes costly refolding or degradation of aberrant proteins. However, high synthesis rates impose constraints on accuracy that lead to an average error rate of 1 amino acid in every 1000 to 10000, implying that at least 15% of proteins have an average of 1 misincorporated amino acid [1-4]. Protein quality control systems, which in eukaryotes involve the ubiquitin-proteasome system (UPS), the unfolded protein response (UPR) and autophagy [5-7], ensure that levels of protein mistranslation are compatible with life. Mistranslation is produced by erroneous mischarging of transfer RNA (error rate $>10^{-4}$) and tRNA-mRNA mispairing in the ribosome (error *in vitro* 10^{-4}) [8,9]. However, these errors do not occur at the same level for all amino acids or with similar frequencies at different codon sites; chemically similar amino acids are more often mischarged on tRNAs, and near cognate codons (codons that differ in one base only) lead to higher error rates than noncognate codons. For example, the small difference of a single methyl group between Ile and Val leads to isoleucyl-tRNA synthetase (IleRS) mischarging of tRNA^{Ile} with Val at a rate of ~1% [10]. In *E. coli* and mammalian cells, Asn and His starvation results in Lys and Gln misincorporation, respectively [11], as low levels of charged tRNA^{Asn} and tRNA^{His} allow charged tRNA^{Lys} and tRNA^{Gln} to read codons that differ in the third base [12,13].

Most surprisingly, several microorganisms use protein mistranslation to functionally diversify their proteome and expand adaptation capacity – adaptive mistranslation [14]. For instance, *Mycoplasma* spp have frequent point mutations and deletions in the editing site of ThrRS, LeuRS and PheRS [15,16] and poorly discriminate mischarged non-cognate tRNAs [17]. AsnRS and GlnRS are absent in *Mycobacterium* species, instead they produce charged tRNAs through natural Glu-tRNA^{Gln} and Asp-tRNA^{Asn} misacylation, followed by amino acid transamidation of the tRNA by the protein complex GatCAB [18]. Mutations in GatCAB subunits found in natural isolates of *M. tuberculosis* or specific physiological conditions increase Gln-to-Glu and Asn-to-Asn mistranslation [19]. Similarly, the fungal pathogen *Candida albicans* mistranslates naturally at a very

high level by misincorporating both Ser (97%) and Leu (3%) at CUG codons [20]. Moreover, the MetRS loses its natural charging specificity through phosphorylation (humans), temperature dependent structural shift (archeon *Aeropyrum pernix*) and removal of a succinyllysine modification (*E. coli*), it also charges various noncognate tRNAs, leading to Met misincorporation into proteins at multiple codon sites [21-23]. These counterintuitive selective advantages produced by mistranslation [24,25] have direct phenotypic consequences. Indeed, in *Mycobacterium tuberculosis* PBE increase resistance to rifampicin, in *Mycoplasma* spp mistranslation contributes to evasion of the host's immune system by increasing cell surface variability (antigenic variation) and in *C. albicans* mistranslation increases survival in the mouse gut and accelerates evolution of resistance to fluconazole. In humans and other organisms Met misincorporation, which can be reversibly oxidized and reduced, protects proteins against oxidative stress with minimal disruption to their structure [20,25-28]. The above cases of regulated mistranslation contrast sharply with random amino acid misincorporation events induced by stress or by damage to the protein synthesis machinery. The latter normally disrupts protein structure, increases protein degradation and activates the proteotoxic stress network leading to cell death. It can also be a serious problem in the production of therapeutics. Overexpression of recombinant proteins in *E. coli*, mammalian cells or other heterologous hosts can result in high levels of amino acid misincorporation reducing the therapeutic value of the recombinant therapeutics. For example, monoclonal antibody production in batch cultures of Chinese hamster ovary (CHO) cells often results in Ser-to-Asn, Meta- and ortho-Tyr-to-Phe misincorporation [29]. Similar misincorporations are observed during production of IGF-1, Interleukin 2, human haemoglobin and other proteins in *E. coli* [30-32], demonstrating that the identification and quantification of mistranslated proteins is relevant to the biotech

*Corresponding authors: Manuel AS Santos, Department of Medical Sciences, iBiMED- Institute of Biomedicine, University of Aveiro, 3910-193 Aveiro, Portugal, Tel: +351234247248; E-mail: msantos@ua.pt

Received January 22, 2018; Accepted February 01, 2018; Published February 10, 2018

Citation: Tavares J, Assis-Santos F, Santos MAS (2018) Proteomics Analysis for Amino Acid Misincorporation Detection: Mini Review. J Proteomics Bioinform 11: 045-050. doi: [10.4172/jpb.1000464](https://doi.org/10.4172/jpb.1000464)

Copyright: © 2018 Tavares J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

and pharmaceutical industries. Detection and quantification of amino acid misincorporation rates remain a significant challenge since the abundance of error-free proteins is considerably higher than that of aberrant proteins. A variety of mass spectrometry methods (MS) have been developed to directly identify and quantify amino acids misincorporated into proteins, but sensitivity, mass-spectra complexity and a lack of statistical and computational methods to filter rare mistranslated peptides continue to be important challenges that must be overcome. Fluorescent, luminescent, radioactive and other methods based on specific gain of function mutations or detection of radioactive peptides, respectively, have also been developed and are frequently used due to their low cost and relatively high reproducibility and robustness, however these methods only identify misincorporated amino acids in specific proteins or at specific protein sites and largely underestimate global misincorporation rates. We describe below the methods in some detail identifying their advantages and disadvantages.

Indirect Measurements of Amino Acid Misincorporation Rates

The indirect methods fall into three groups (Table 1): 1) use of a specific amino acid not observed in a reporter protein; 2) measurement of changes in the isoelectric point of a protein; 3) use of biochemical or genetic reporter systems. These methods detect a specific type of amino acid misincorporation at a time, normally a heterologous recombinant

protein expressed in a specific host cell. Low frequency amino acid misincorporations are normally underrepresented or are not detected, and there is also significant error detection bias due to instability of the mistranslated recombinant proteins [33]. Loftfield and Vanderjagt were among the first scientists to detect amino acid misincorporations in proteins by incubating rabbit reticulocytes with radioactive valine and analyzing tryptic fragments of the purified haemoglobin. They estimated a misreading probability per codon of approximately $2-6 \times 10^{-4}$ [34]. Other groups have taken advantage of radiolabeled amino acids to detect the incorporation of ^{35}S -cysteine into the cysteine-free flagellin of *E. coli* (Figure 1A) [2,35,36] and also at Arg codon sites of an *E. coli* ribosomal protein [2] by growing cells in a medium supplied with a radioactive amino acid that was not present in a protein of interest. Others estimated the error frequency during translation by detecting differences in the isoelectric point (IP) of selected proteins under normal conditions and during amino acid starvation for Asn and Lys [1,12,13]. Such IP differences result in the separation of mistranslated and wild type protein isoforms in two-dimensional (2D) gel electrophoresis (Figure 1B). During Asn starvation, Lys errors at Asn codons altered the isoelectric point of the MS2 coat protein and the measured error frequency reached 5×10^{-3} [12,13] Khazaie and collaborators evaluated radioactive phage proteins produced by *E. coli* using 2D gel electrophoresis and were able to detect multiple protein isoforms labeled with misincorporated His and Trp at similar error rates [36].

Detection method	Principle	Type of error	Error rate/codon	Organism/Cell	Ref.
Indirect Methods					
Detection of a specific amino acid not found in the protein	Radioactively labeled Val	Ile → Val	$2.0 - 6.0 \times 10^{-4}$	Rabbit	[34]
Detection of a specific amino acid not found in the protein	^{35}S -Cys incorporation in Cys-free flagellin	Arg → Cys	1.0×10^{-4}	<i>E. coli</i>	[35]
Detection of a specific amino acid not found in the protein	Radioactively labeled Cys and interference with protease digestion at Arg residues	Arg → Cys Trp → Cys	10^{-3} $3.0 - 4.0 \times 10^{-3}$	<i>E. coli</i>	[2]
Detection of a specific amino acid not found in the protein	Detection of radioactive His, Lys, Trp-free phage proteins	Lys, Leu, Trp misincorporation	$3.0 - 4.0 \times 10^{-4}$	<i>E. coli</i>	[36]
Electrophoretic heterogeneity in the protein	Change of the isoelectric point pattern	Asn → Lys	5.0×10^{-3}	<i>E. coli</i>	[13]
Electrophoretic heterogeneity in the protein	Change of the isoelectric point pattern	Several misincorporations	4.0×10^{-4}	<i>E. coli</i>	[1]
Reporter system	Restoration of β-lactamase activity	Gly → Ser	10^{-3}	<i>E. coli</i>	[37]
Reporter system	Restoration of CAT activity	Tyr → His	0.5×10^{-5}	<i>S. cerevisiae</i>	[38]
Reporter system	Restoration of F-luc activity	Lys misincorporation	$2.0 \times 10^{-4} - 3.6 \times 10^{-3}$	<i>E. coli</i>	[8]
Reporter system	Restoration of R-luc activity	Gln → Glu Asn → Asp	0.20% 0.80%	<i>M. segmentis</i>	[25]
Reporter system	Restoration of GFP fluorescence	Ser/Leu ambiguous codon	0 - 98%	<i>C. albicans</i>	[20]
Reporter system	Restoration of GFP fluorescence	Pro → Ala	~6%	<i>S. cerevisiae</i>	[53]
Reporter system	Restoration of mCherry fluorescence	Glu → Met	ND	Human cells	[42]
Direct Methods					
Reporter protein	MRM mode MS/MS	Leu → Ser	1 - 2 %	<i>S. cerevisiae</i>	[43]
Reporter protein	MS/MS of modified side-chains	Asn → Asp	~10%	<i>E. coli</i>	[45]
Reporter protein	MS/MS and modified database	Ser → Asn	0.01 - 0.2%	<i>E. coli</i> ; CHO and NS0 cell lines	[44]
Reporter protein	MS/MS and precursor ion exclusion approach	Several misincorporations	0.0010%	<i>E. coli</i> ; mammalian cells	[47]
Reporter protein	MRM mode MS/MS	Phe → Tyr	0.02%	<i>S. cerevisiae</i>	[52]
Reporter protein	MS/MS and data processing with SPIDER algorithm	Pro → Ala	ND	<i>S. cerevisiae</i>	[53]
Detection of single amino acid misincorporations	Shotgun MS analysis and dynamic SILAC approach	Asp → Glu Glu → Asp	$\sim 1 \times 10^{-3}$ $\sim 0.5 \times 10^{-4}$	Mice	[54]
Detection of single amino acid misincorporations	Systematic shotgun MS analysis and Super-SILAC approach	Leu → Nva	~0.14%	<i>E. coli</i>	[55]

Table 1: Methods for detection of mistranslation.

Gain of function reporter systems are also widely used to determine amino acid misincorporation levels at specific codons. The most commonly used recombinant reporters are GFP, luciferases and β -gal that have loss of function mutations at critical functional residues and enzymatic activity can only be restored through misincorporation of the wild type amino acid (Figure 1C). Recovery of function can be quantified by measuring fluorescent, luminescent or chromogenic signals of those proteins providing an indirect measure of error rates, i.e., misincorporation frequency can be estimated as the ratio of mutant versus wild type enzymatic activity.

Toth and collaborators were the first to use a beta-lactamase gene with a mutation in a AGC (Ser) codon change to GGA/GGC – (Gly), which inactivates beta-lactamase activity. Expression of this gene in mutant *E. coli* with compromised translational fidelity restored beta-lactamase activity providing an indirect measure of Ser misincorporation at the mutant Gly sites [37]. Similar strategies using mutant chloramphenicol acetyl transferase (CAT_{III}) were used to determine mistranslation levels in *S. cerevisiae* since His misincorporation at Tyr codon sites restores CAT_{III} activity [38]. These reporter systems are sensitive but have the disadvantage of only measuring one type of misincorporation at a time and underestimate amino acid misincorporation rates. Kramer and Farabaugh used a dual reporter system consisting of a fusion of a mutant form of the firefly luciferase (F-luc) and *Renilla* luciferase (R-luc), whose luminescence can be measured independently. These two proteins are expressed as a single polypeptide, but their activities are independent, therefore, any difference in the activity of F-luc relative to R-luc reflects changes in protein activity. The F-luc has a series of inactivating mutations at a codon (L529) site that encodes an essential lysine residue,



permitting monitoring gain of function activity of F-luc caused by Lys misincorporation at the various mutant codons [8]. This approach was used and adapted by other groups and became widely used by the scientific community [25,39]. Javid and co-workers inactivated R-luc, instead of F-luc, by mutating the active site to detect and quantify Glu and Asp misincorporations at Gln and Asn codons in mycobacteria [25]. The above approaches have the caveat of requiring the addition of substrate or cofactors to cell lysates to measure enzymatic activity. To circumvent this limitation, a fluorescent reporter assay based on the expression of recombinant GFP capable of reporting amino acid misincorporation at a specific site was developed [20,40,41]. Bezerra et al. constructed a codon-optimized yEGFP gene, in which the UUG Leu codon at position 201 was mutated to an atypical CUG Ser codon in *C. albicans*. Leu misincorporation at the CUG position resulted in a gain of function of the fluorescent protein that was therefore able to report accurate levels of Leu misincorporation in that pathogenic fungus [20]. Recently, a dual fluorescent reporter (EGFP-mCherry fusion protein) was developed for the detection and quantification of Met mistranslation in mammalian cells. This reporter is a mCherry gene containing a mutation at position 72 (Met72) fused to a wild-type EGFP gene, which is used to determine the expression level of the reporter [42]. The advantage of these dual reporter systems relative to single protein reports is that they overcome biases in the determination of enzyme activity levels caused by variation in protein expression. The other advantage is that fluorescence can be detected by microscopy, flow cytometry and other methods allowing for a broader range of applications.

Direct Measurements of Amino Acid Misincorporation Rates

Advances in tandem mass spectrometry (MS/MS) has allowed researchers to directly measure mistranslation. Here, we differentiate MS/MS analysis of reporter proteins and MS/MS analysis of a total proteome (Table 1 and Figure 2).

MS/MS of reporter proteins

MS/MS analysis of reporter proteins purified to near homogeneity reduces the complexity of the MS spectra search space, increases the probability of detecting mistranslated peptides and the reproducibility



of the measurements. Deep quantitative MS/MS analysis of the reporter peptides against an appropriate database can detect amino-acid substitutions and locate the specific sites where they occur along the primary structure. This highly sensitive approach permits identification and quantification of a wide variety of amino acid misincorporations, but does not produce a comprehensive view of mistranslation rates because single proteins rarely contain the 61 sense codons of the genetic code; codon context effects affect mistranslation rate and each protein only contains a small number of possible contexts of each codon [43,44]. In 2007, Silva and collaborators constructed a reporter system to detect Ser misincorporation at Leu CUG codon sites in *C. albicans*. The target protein was expressed, purified by affinity chromatography, digested and the peptides were separated chromatographically before mass spectrometric analyses in multiple reaction monitoring (MRM) mode [43]. Another group quantified Asp misincorporation in recombinant Dihydrofolate Reductase (DHFR) that was digested and methylated before MS/MS analysis. The level of amino acid misincorporation was calculated as the ratio of absolute intensities of the mistranslated peptide to the WT peptide peaks [45]. Several mistranslation errors were identified using this approach in different organisms and cell lines, namely CHO cells, which are frequently used for recombinant expression of target proteins [44,46-49]. This strategy was also used to demonstrate that the tyrosyl-tRNA synthetase is highly error prone [50,51] and that the editing site of some aminoacyl-tRNA synthetases controls the sensitivity of amino acid stress responses [52].

Hoffman and collaborators combined direct and indirect methods to detect Ala substitutions in a reporter protein for MS/MS analysis (Two Tel2-Interacting protein - Tti2) and in *EGFP* reporter protein. Comparison of the MS/MS and *EGFP* reporter data showed similar levels of Ala misincorporation at Pro codons, indicating that both methods are reliable and can be used alone or in combination [53].

Proteome scale MS/MS analysis

Single amino acid substitutions in the whole proteome of an organism or cell line can be detected using a systematic shotgun mass spectrometry-based proteomics approach. In this type of approach, complex protein samples are digested with a protease, usually trypsin, and digested peptides are fractionated on HPLC columns, prior to tandem mass spectrometry analysis. Identification of proteins is achieved by searching the MS/MS spectra against a protein database, in which spectra derived from peptide fragmentation is compared with theoretical spectra generated from *in silico* digestion of the protein database. Recently, quantification of single amino acid misincorporations was achieved using SILAC methods, which use stable isotopes to label amino acids in cell culture. Tuorto et al. identified high frequency of Asp to Glu substitutions and vice-versa in the proteome of mutant mice bone marrow using dynamic SILAC techniques [54]. Dynamic SILAC is a technique that allows detection of variances in *de novo* protein synthesis since labeled amino acids are present in the growth medium for a short period of time. On the other hand, Cveticic et al. detected norvaline misincorporation in the *E. coli* proteome using Super-SILAC labeling [55]. In this work, labeled proteins from different samples were mixed and used as a spike-in standard, allowing a more accurate quantification of unlabeled samples [56]. Along with the SILAC approaches, a spectral counting strategy has been also used to quantify mistranslation [55]. This approach combines the number of identified MS/MS spectra from the same protein throughout multiple LC-MS runs to assess relative protein abundance.

A handful of researchers have been able to identify a small number of misincorporations from large MS spectra databases

using prior knowledge of the misincorporated amino acid and codon sites, demonstrating that identification and quantification of misincorporated amino acids by MS/MS is feasible. However, the high percentage of false positives observed in those data sets highlights the need to optimize the methodologies used in sample preparation, MS/MS and bioinformatics. A successful proteomics study should be well designed, selecting the sample preparation approach that best fits the experiment to improve detection of low abundance proteins and peptides [57]. Indeed, the major challenge in detecting mistranslated peptides is their low abundance against a background of wild-type peptides. Additionally, amino acid misincorporations detected by MS/MS can be misinterpreted as post-translational modifications, such as phosphorylation, glycosylation, nitration and acylation, reducing the reliability and reproducibility of this approach [47]. Bioinformatic tools are also critical in this field. Some researchers use algorithms that search against a database containing all possible modified peptides of their target proteins [45]. However, these databases provide a biased and limited view of the mistranslated peptides, forcing spectra to be identified as peptides present in the manipulated database [58]. To overcome these limitations, one could use error-tolerance searches, such as *de novo* approaches, at the amino acid sequence level since *de novo* sequencing identifies peptide sequences by analyzing MS/MS spectra. Other algorithms combine *de novo* sequencing with database search of the candidate peptides looking for homology [59]. This kind of strategy could help in the detection of misincorporations at a global proteome scale, decreasing the number of false positives.

Conclusion

Several approaches are now available to identify and quantify protein biosynthesis errors, particularly those allying MS sensitivity approaches and reporter protein systems. Despite the technical difficulties in the identification and quantification of mistranslation errors at the proteome scale, recent works show that errors are possible to quantify and it is likely that future improvements in mass spectrometry and bioinformatics will allow us to tackle the relevance of protein synthesis errors for aging and human disease in a robust and comprehensive manner. Such improvements should also allow the development of methods and host cells essential for the production of high quality recombinant therapeutics.

Acknowledgments

This work was supported by the Portuguese Foundation for Science and Technology through COMPETE2020 and FEDER funds (project FCT-PTDC/BEX-BCM/2121/2014) and Integrated Projects FEDER/CENTRO-45-2015-01.

References

1. Ellis N, Gallant J (1982) An estimate of the global error frequency in translation. *MGG Mol Gen Genet* 188: 169-172.
2. Bouadloun F, Donner D, Kurland CG (1983) Codon-specific missense errors in vivo. *EMBO J* 2: 1351-1356.
3. Parker J (1989) Errors and alternatives in reading the universal genetic code. *Microbiol Rev* 53: 273-98.
4. Ogle JM, Ramakrishnan V (2005) Structural insights into translational fidelity. *Annu Rev Biochem* 74: 129-177.
5. Frydman J (2001) Folding of newly translated proteins in vivo: The Role of Molecular Chaperones. *Annu Rev Biochem* 70: 603-647.
6. Ciechanover A (1998) The ubiquitin-proteasome pathway: on protein death and cell life. *EMBO J* 17: 7151-7160.
7. Liliienbaum A (2013) Relationship between the proteasomal system and autophagy. *Int J Biochem Mol Biol* 4: 1-26.
8. Kramer EB, Farabaugh PJ (2007) The frequency of translational misreading errors in *E. coli* is largely determined by tRNA competition. *RNA* 13: 87-96.

9. Ling J, Söll D (2010) Severe oxidative stress induces protein mistranslation through impairment of an aminoacyl-tRNA synthetase editing site. *Proc Natl Acad Sci USA* 107: 4028-4033.
10. Hale SP, Auld DS, Schmidt E, Schimmel P (1997) Discrete Determinants in Transfer RNA for Editing and Aminoacylation. *Science* 276: 1250-1252.
11. Parker J, Pollard JW, Friesen JD, Stanners CP (1978) Stuttering: High-level mistranslation in animal and bacterial cells. *Proc Natl Acad Sci USA* 75: 1091-1095.
12. Parker J, Friesen JD (1980) "Two out of three" codon reading leading to mistranslation in vivo. *MGG Mol Gen Genet* 177(3): 439-445.
13. Parker J, Johnston TC, Borgia PT (1980) Mistranslation in cells infected with the bacteriophage MS2: Direct evidence of Lys for Asn substitution. *MGG Mol Gen Genet* 180: 275-281.
14. Pan T (2013) Adaptive translation as a mechanism of stress response and adaptation. *Annu Rev Genet* 47: 121-137.
15. Li L, Boniecki MT, Jaffe JD, Imai BS, Yau PM, et al. (2011) Naturally occurring aminoacyl-tRNA synthetases editing-domain mutations that cause mistranslation in *Mycoplasma* parasites. *Proc Natl Acad Sci USA* 108: 9378-9383.
16. Li L, Palencia A, Lukk T, Li Z, Luthey-Schulten ZA, et al. (2013) Leucyl-tRNA synthetase editing domain functions as a molecular rheostat to control codon ambiguity in *Mycoplasma* pathogens. *Proc Natl Acad Sci USA* 110: 3817-3822.
17. Yadavalli SS, Ibba M (2013) Selection of tRNA charging quality control mechanisms that increase mistranslation of the genetic code. *Nucleic Acids Res* 41: 1104-1112.
18. Nair N, Raff H, Islam MT, Feen M, Sheppard K, et al. (2016) The *Bacillus subtilis* and *Bacillus halodurans* Aspartyl-tRNA Synthetases Retain Recognition of tRNA Asn. *J Mol Biol* 428: 618-630.
19. Su HW, Zhu JH, Li H, Cai RJ, Ealand C, et al. (2016) The essential mycobacterial amidotransferase GatCAB is a modulator of specific translational fidelity. *Nat Microbiol* 1: 16147.
20. Bezerra AR, Simões J, Lee W, Rung J, Weil T, et al. (2013) Reversion of a fungal genetic code alteration links proteome instability with genomic and phenotypic diversification. *Proc Natl Acad Sci USA* 110: 11079-11084.
21. Lee JY, Kim DG, Kim BG, Yang WS, Hong J, et al. (2014) Promiscuous methionyl-tRNA synthetase mediates adaptive mistranslation to protect cells against oxidative stress. *J Cell Sci* 127: 4234-4245.
22. Schwartz MH, Pan T (2016) Temperature dependent mistranslation in a hyperthermophile adapts proteins to lower temperatures. *Nucleic Acids Res* 44: 294-303.
23. Schwartz MH, Waldbauer JR, Zhang L, Pan T, (2016) Global tRNA misacylation induced by anaerobiosis and antibiotic exposure broadly increases stress resistance in *Escherichia coli*. *Nucleic Acids Res* 44: 10292-10303.
24. Santos MAS, Cheesman C, Costa V, Moradas-Ferreira P, Tuite MF, et al. (1999) Selective advantages created by codon ambiguity allowed for the evolution of an alternative genetic code in *Candida* spp. *Mol Microbiol* 31: 937-947.
25. Javid B, Sorrentino F, Toosky M, Zheng W, Pinkham JT, et al. (2014) Mycobacterial mistranslation is necessary and sufficient for rifampicin phenotypic resistance. *Proc Natl Acad Sci USA* 111: 1132-1137.
26. Finlay BB, McFadden G (2006) Anti-immunology: evasion of the host immune system by bacterial and viral pathogens. *Cell* 124: 767-782.
27. Levine RL, Mosoni L, Berlett BS, Stadtman ER, (1996) Methionine residues as endogenous antioxidants in proteins. *Proc Natl Acad Sci USA* 93: 15036-15040.
28. Luo S, Levine RL (2009) Methionine in proteins defends against oxidative stress. *FASEB J* 23: 464-472.
29. Popp O, Larraillat V, Kettenberger H, Gorr IH, Hilger M, et al. (2015) Molecular polygamy: The promiscuity of l-phenylalanyl-tRNA-synthetase triggers misincorporation of meta- and ortho-tyrosine in monoclonal antibodies expressed by Chinese hamster ovary cells. *Biotechnol Bioeng* 112: 1187-1199.
30. Seetharam R, Heeren RA, Wong EY, Braford SR, Klein BK, et al. (1988) Mistranslation in IGF-1 during over-expression of the protein in *Escherichia coli* using a synthetic gene containing low frequency codons. *Biochem Biophys Res Commun* 155: 518-523.
31. Tsai LB, Lu HS, Kenney WC, Curless CC, Klein ML, et al. (1988) Control of misincorporation of de novo synthesized norleucine into recombinant interleukin-2 in *E. coli*. *Biochem Biophys Res Commun* 156: 733-739.
32. Apostol I, Levine J, Lippincott J, Leach J, Hess E, et al. (1997) Incorporation of norvaline at leucine positions in recombinant human hemoglobin expressed in *Escherichia coli*. *J Biol Chem* 272: 28980-28988.
33. Goldberg AL (1972) Correlation between Rates of Degradation of Bacterial Proteins in Vivo and Their Sensitivity to Proteases. *Proc Natl Acad Sci USA* 69: 2640-2644.
34. Lofffield RB, Vanderjagt D (1972) The Frequency of Errors in Protein Biosynthesis. *Biochem J* 128: 1353-1356.
35. Edelmann P, Gallant J (1977) Mistranslation in *E. coli*. *Cell* 10: 131-137.
36. Khazaie K, Buchanan JH, Rosenberger RF (1984) The accuracy of Qb RNA translation. 1. Errors during the synthesis of Qb proteins by intact *Escherichia coli* cells. *Eur J Biochem* 144: 485-489.
37. Toth MJ, Murgola EJ, Schimmel P (1988) Evidence for a unique first position codon-anticodon mismatch in vivo. *J Mol Biol* 201: 451-454.
38. Stansfield I, Jones KM, Herbert P, Lewendon A, Shaw WV, et al. (1998) Missense translation errors in *Saccharomyces cerevisiae*. *J Mol Biol* 282: 13-24.
39. Moreno-Martinez E, Vallieres C, Holland SL, Avery SV (2015) Novel, Synergistic Antifungal Combinations that Target Translation Fidelity. *Sci Rep* 5.
40. Ilegems E, Pick HM, Vogel H (2002) Monitoring mis-acylated tRNA suppression efficiency in mammalian cells via EGFP fluorescence recovery. *Nucleic Acids Res* 30: 128.
41. Meyerovich M, Mamou G, Ben-Yehuda S (2010) Visualizing high error levels during gene expression in living bacterial cells. *Proc Natl Acad Sci USA* 107: 11543-11548.
42. Gomes AC, Kordala AJ, Strack R, Wang X, Geslain R, et al. (2016) A dual fluorescent reporter for the investigation of methionine mistranslation in live cells. *RNA* 22: 467-476.
43. Silva RM, Paredes JA, Moura GR (2007) Genetic Code Alteration Plays a Critical Role in the Evolution of *Candida* Genes Quantitative Analysis of Single Amino Acid Changes using a 4000 QTRAP system. *Applied Biosystems* 33.
44. Yu XC, Borisov OV, Alvarez M, Michels DA, Wang YJ, et al. (2009) Identification of codon-specific serine to asparagine mistranslation in recombinant monoclonal antibodies by high-resolution mass spectrometry. *Anal Chem* 81: 9282-9290.
45. Ruan L, Zhou C, Jin E, Kucharavy A, Zhang Y, et al. (2017) Cytosolic proteostasis through importing of misfolded proteins into mitochondria. *Nature* 543: 443-446.
46. Guo D, Gao A, Michels DA, Feeney L, Eng M, et al. (2010) Mechanisms of unintended amino acid sequence changes in recombinant monoclonal antibodies expressed in Chinese Hamster Ovary (CHO) cells. *Biotechnol Bioeng* 107: 163-171.
47. Zhang Z, Shah B, Bondarenko PV (2013) G/U and certain wobble position mismatches as possible main causes of amino acid misincorporations. *Biochemistry* 52: 8165-8176.
48. Harris RP, Mattocks J, Green PS, Moffatt F, Kilby PM, et al. (2012) Determination and control of low-level amino acid misincorporation in human thioredoxin protein produced in a recombinant *Escherichia coli* production system. *Biotechnol Bioeng* 109: 1987-1995.
49. Ragionieri L, Vitorino R, Frommlet J, Oliveira JL, Gaspar P, et al. (2014) Improving the accuracy of recombinant protein production through integration of bioinformatics, statistical and mass spectrometry methodologies. *FEBS J* 1-19.
50. Feeney L, Carvalhal V, Yu XC, Chan B, Michels DA, et al. (2013) Eliminating tyrosine sequence variants in CHO cell lines producing recombinant monoclonal antibodies. *Biotechnol Bioeng* 110: 1087-1097.
51. Raina M, Moghal A, Kano A, Jerums A, Schnier PD, et al. (2014) Reduced Amino Acid Specificity of Mammalian Tyrosyl-tRNA Synthetase Is Associated with Elevated Mistranslation of Tyr Codons. *J Biol Chem* 289: 17780-17790.
52. Mohler K, Mann R, Bullwinkle TJ, Hopkins K, Hwang L, et al. (2017) Editing of misaminoacylated tRNA controls the sensitivity of amino acid stress responses in *Saccharomyces cerevisiae*. *Nucleic Acids Res* 45: 3985-3996.

53. Hoffman KS, Berg MD, Shilton BH, Brandl CJ, O'Donoghue P, et al. (2017) Genetic selection for mistranslation rescues a defective co-chaperone in yeast. *Nucleic Acids Res* 45: 3407-3421.
54. Tuorto F, Herbst F, Alerasool N, Bender S, Popp O, et al. (2015) The tRNA methyltransferase Dnmt2 is required for accurate polypeptide synthesis during haematopoiesis. *Embo J* 34: 2350-2362.
55. Cvetesic N, Semanjski M, Soufi B, Krug K, Gruic-Sovulj I, et al. (2016) Proteome-wide measurement of non-canonical bacterial mistranslation by quantitative mass spectrometry of protein modifications. *Sci Rep* 6: 28631.
56. Shenoy A, Geiger T (2014) Super-SILAC: Current trends and future perspectives. *Expert Review of Proteomics* 12: 13-19.
57. Eriksson J, Fenyö D (2007) Improving the success rate of proteome analysis by modeling protein-abundance distributions and experimental designs. *Nature Biotechnology* 25: 651-655.
58. Sticker A, Martens L, Clement L (2017) Mass spectrometrists should search only for peptides they care about. *Nature Methods* 14: 643-644.
59. Ma B, Johnson R (2012) De Novo Sequencing and Homology Searching. *Mol Cell Proteomics* 11: O111.014902.