OPEN ACCESS Freely available online

Journal of Cell Science & Therapy

ISSN: 2157-7013

Short Communication

Repair Mechanisms and Their Role in Maintaining Genome Stability and Therapeutic Research

Nivara Colsten*

Department of Genetic Cell Medicine, Horizon Biomedical University, Stockholm, Sweden

DESCRIPTION

Cells are continuously exposed to conditions that may affect the stability of their genetic material. Internal biological processes, environmental influences, and cellular activities can produce changes in Deoxyribonucleic Acid (DNA) that require correction to preserve normal function. Cellular repair mechanisms are essential systems that identify and correct genetic damage, allowing cells to maintain accurate information required for survival and reproduction. Research into these mechanisms has become an important area within cell science and therapy because disturbances in DNA repair processes are associated with various diseases and cellular abnormalities [1]. DNA contains the instructions required for controlling cellular activities, including growth, division, metabolism, and specialized functions [2]. Maintaining the accuracy of this information is necessary for healthy cell behavior. Throughout the life of a cell, DNA may experience different forms of damage, including small chemical changes, breaks in DNA strands, and errors that occur during replication. Cells possess multiple repair systems that recognize these changes and restore genetic stability.

DNA damage can occur through both internal and external sources. Normal cellular metabolism can produce molecules capable of affecting DNA structure, while environmental factors such as radiation and certain chemical exposures may also contribute to genetic alterations [3]. Cells continuously monitor DNA integrity and activate repair responses when damage is detected. One important repair process is nucleotide excision repair, which removes damaged sections of DNA and replaces them with correctly formed sequences. This mechanism is particularly important for correcting damage caused by ultraviolet exposure and certain chemical factors. Research into nucleotide excision repair has improved understanding of how cells protect genetic information from environmental influences.

Base excision repair is another essential mechanism involved in maintaining DNA stability. This pathway corrects small changes affecting individual DNA components [4]. Specialized enzymes identify damaged sections, remove affected molecules, and restore the correct genetic structure. Scientists study this process to understand how cells manage everyday molecular damage. Double-strand DNA breaks represent a more serious form of genetic damage because both strands of the DNA molecule are affected. Cells respond to these breaks through repair pathways that restore DNA structure and prevent harmful changes [5]. Two major repair methods involved in this process are homologous recombination and non-homologous end joining. These mechanisms allow cells to correct damage while preserving genetic information.

Homologous recombination uses similar genetic sequences as a reference during repair. This process provides a highly controlled method for restoring damaged DNA regions. It is especially active during specific stages of the cell cycle when accurate repair is required [6]. Researchers study homologous recombination because changes in this pathway are linked with certain inherited disorders and cancer development. Non-homologous end joining repairs DNA breaks by directly reconnecting damaged DNA ends. Although this process is faster than some other repair mechanisms, it requires careful regulation to maintain genetic accuracy. Scientists continue examining how cells select and coordinate different repair pathways depending on the type of damage present.

DNA repair is closely connected with cell cycle regulation. Before cells divide, they must ensure that their genetic material is accurately copied and maintained. Checkpoint systems monitor DNA conditions and delay cell division when damage is detected. These control systems provide time for repair activities to occur before genetic information is passed to new cells. Inherited disorders caused by DNA repair defects provide additional information about these biological systems. Certain genetic conditions occur when specific repair mechanisms do not function properly [7]. Studying these disorders helps scientists understand the importance of DNA maintenance and the effects of repair pathway disruption.

The connection between DNA repair and aging has become an important area of cell science research. Over time, cells may

Correspondence to: Nivara Colsten, Department of Genetic Cell Medicine, Horizon Biomedical University, Stockholm, Sweden, E-mail: nivara.colsten@horizonbiomed.se

Received: 02-Jan-2026, Manuscript No. JCEST-26-43540; **Editor assigned:** 05-Jan-2026, PreQC No. JCEST-26-43540(PQ); **Reviewed:** 19-Jan-2026, QC No. JCEST-26-43540; **Revised:** 26-Jan-2026, Manuscript No. JCEST-26-43540 (R); **Published:** 02-Feb-2026, DOI: 10.35248/2157-7013.26.17.559

Citation: Colsten N (2026). Repair Mechanisms and Their Role in Maintaining Genome Stability and Therapeutic Research. J Cell Sci Therapy. 17:559.

Copyright: © 2026 Colsten N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

experience increasing levels of genetic damage due to repeated exposure to internal and external factors. Although repair systems continuously correct many forms of damage, changes in repair efficiency may influence cellular aging and tissue function. Scientists are investigating how DNA maintenance processes affect age-related changes in different organs [8]. Mitochondrial DNA repair has received increasing scientific interest because mitochondria contain their own genetic material. Damage to mitochondrial DNA can influence energy production and cellular performance. Researchers are studying how mitochondria maintain genetic stability and how changes in mitochondrial repair processes affect human health [9]. Cancer treatment research has also examined how tumor cells use DNA repair mechanisms. Some cancer cells develop strong repair abilities that allow them to survive conditions that damage genetic material. Scientists are studying these characteristics to understand how tumors respond to different treatments and how repair pathways influence treatment outcomes [10].

CONCLUSION

The study of DNA repair proteins has provided valuable information about molecular interactions inside cells. These proteins detect damaged DNA, organize repair activities, and coordinate responses with other cellular systems. Researchers continue identifying how different repair proteins function together and how changes in these proteins affect cellular health. The study of DNA repair also contributes to understanding cellular responses after injury. When tissues experience damage, cells activate repair systems to restore genetic stability and support recovery. The efficiency of these processes influences how cells respond to challenging conditions and maintain their functions.

REFERENCE

1. Aguilera P, Whalen J, Minguet C, Churikov D, Freudenreich C, Simon MN, et al. The nuclear pore complex prevents sister chromatid recombination during replicative senescence. Nat Commun. 2020;11(1):160.
2. Berti M, Cortez D, and Lopes M. The plasticity of DNA replication forks in response to clinically relevant genotoxic stress. Nat Rev Mol Cell Biol. 2020;21(10):633-651.
3. Chakraborty S, Schirmeisen K, Lambert SA. The multifaceted functions of homologous recombination in dealing with replication-associated DNA damages. DNA Repair (Amst). 2023;129:103548.
4. Dovrat D, Dahan D, Sherman S, Tsirkas I, Elia N, Aharoni A. A live-cell imaging approach for measuring DNA replication rates. Cell Rep. 2018;24(1):252-258.
5. Masnovo C, Paleiov Z, Dovrat D, Baxter LK, Movafaghi S, Aharoni A, et al. Stabilization of expandable DNA repeats by the replication factor Mcm10 promotes cell viability. Nat Commun. 2024;15(1):10532.
6. Mirkin EV, Mirkin SM. Replication fork stalling at natural impediments. Microbiol Mol Biol Rev. 2007;71:13-35.
7. Semlow DR, Walter JC. Mechanisms of vertebrate DNA interstrand cross-link repair. Annu Rev Biochem. 2021;90:107-135.
8. Vanoli F, Fumasoni M, Szakal B, Maloisel L, Branzei D. Replication and recombination factors contributing to recombination-dependent bypass of DNA lesions by template switch. PLoS Genet. 2010;6(11):e1001205.
9. Mouron S, Rodriguez-Acebes S, Martinez-Jimenez MI, Garcia-Gomez S, Chocron S, Blanco L, et al. Repriming of DNA synthesis at stalled replication forks by human primpol. Nat Struct Mol Biol. 2013;20(12):1383-1389.
10. Adolph MB, Cortez D. Mechanisms and regulation of replication fork reversal. DNA Repair. 2024:141:103731.