

# Role of Gut-Brain Axis in the Aetiology of Neurodevelopmental Disorders with Reference to Autism

Afaf El-Ansary<sup>1,2\*</sup>, Ghada H. Shaker<sup>2</sup> and Maha Zaki Rizk<sup>3</sup>

<sup>1</sup>Department of Biochemistry, College of Science, King Saud University, Saudi Arabia

<sup>2</sup>Department of Microbiology and Immunology, College of Pharmacy, Zagazig University, Egypt

<sup>3</sup>Therapeutic Chemistry Department, National Research Center, Dokki, Giza, Egypt

## Abstract

Neurodevelopmental disorders, especially in children, result in brain and nervous system damage. These may result from environmental contaminants, intrauterine environment, infectious diseases or exposure to nanoparticles that cross the blood brain barrier. Gut microbiota directly influence the immune system, nervous system and brain development during microbial colonisation of the newborn (microbiota gut-brain axis) and are controlled and modulated by different endogenous and exogenous factors. Of these factors feeding with human milk creates a healthy microbiota in the infant gut and reduces incidence and severity of infections and promotes normal gastrointestinal function. In addition there is a direct correlation between maternal vaginal and intestinal bacteria, gut microbiota composition, and increased rates of obesity, metabolic and neuropathological disorders such as autism. Gut-brain factors secondary to alterations in gut microbiome by antibiotics or diet may influence brain function in patients with Autism Spectral Disorders (ASD). Children with ASD ingest food products that provide high carbohydrates for bacterial fermentation to produce propionic acid through the bacterial strain *Clostridium difficile*, which is associated with diarrhoea. Treatment strategies to reduce *Clostridium difficile* include probiotics, prebiotics, faecal transplantation and hyperbaric oxygen therapy. Studies of microbiota-gut-brain axis could provide a deeper understanding of the relationship between the intestinal bacteria and their hosts which could help to suggest potential therapeutic strategies through affecting the composition of gut microbiota.

## Introduction

It is quite known that defects in brain function especially in children usually may result in neuro-developmental disorders such as intellectual disability, Attention-Deficit/Hyperactivity Disorder (ADHD), autism, and learning disabilities which is reflected in disabilities to communicate, move or behave. These symptoms usually change with age, although some children may develop permanent disabilities. Diagnosis and treatment of neurodevelopmental disorders often involves a combination of professional therapy, pharmaceuticals, and home- and school-based programs, though, achievement of successful results is difficult [1].

It was previously reported that the child's developing brain and nervous system are susceptible to damage as a result of exposure to environmental pollutants such as lead [2-4], methyl mercury [5] and Polychlorinated Biphenyls (PCBs) [6]. These developmental disorders include reduced cognitive development, lowered intelligence and behavioural deficits and brain trauma. The latter occurs in over 400,000 injuries per year in the US alone, without clarifying the number that may further produce developmental sequelae. It may be subdivided into two major categories, first, injury occurring in infancy or childhood and second, congenital injury (uncomplicated premature birth) resulting from asphyxia (obstruction of the trachea), hypoxia (lack of oxygen to the brain) or the mechanical trauma of the birth process itself [7].

It should be pointed out that fetal development is affected by the intrauterine environment and any disruptions in the latter may eventually lead to various learning, behavioural, and neurological disorders in childhood, as well as complex diseases such as obesity, stress and cardiovascular problems later in life [8], in addition to certain infectious diseases such as schizophrenia [9], or congenital toxoplasmosis. This latter parasite may result in formation of cysts in the brain and other organs, and even though there is a marked maternal IgG immune response, the parasite was found to continue proliferation in the brain [10]. Other diseases include congenital syphilis and

measles which may progress to neurosyphilis and subacute sclerosing panencephalitis respectively in addition to multiple other symptoms.

Furthermore, since the placenta is at a literal interface between maternal and fetal cells, maternal and fetal cells reside in the placenta and also maternal or intrauterine environment are necessarily conveyed to the developing embryo via the placenta. Consequently, the placenta is likely to play a critical role in modulating immune protection and the availability of nutrients and endocrine factors to the offspring. However, factors as autoimmunity, growth restriction and hypoxia implicate the role of the placenta and its involvement in development of neurological complications [11]. In this concern, early prenatal insults are usually involved in the occurrence of neuro-developmental disorders such as schizophrenia, autism and cerebral palsy.

Most recently, exposure to nanoparticles have been shown to accumulate in organs, cross the Blood-Brain Barrier (BBB) and placenta, and have the potential to elicit Developmental Neurotoxicity (DNT).

Another factor that contributes to brain development and behaviour, and also influences the nervous system, is the gut microbiota especially during microbial colonisation of the new born. Studies of microbiota-gut-brain axis could provide a deeper understanding of the

**\*Corresponding author:** Afaf El-Ansary, Department of Biochemistry, College of Science, King Saud University, P.O Box 22452, Zip Code 11495. Riyadh, Saudi Arabia, E-mail: [elansary@KSU.EDU.SA](mailto:elansary@KSU.EDU.SA)

**Received** April 03, 2013; **Accepted** June 04, 2013; **Published** June 06, 2013

**Citation:** El-Ansary A, Shaker GH, Rizk MZ (2013) Role of Gut-Brain Axis in the Aetiology of Neurodevelopmental Disorders with Reference to Autism. J Clin Toxicol S6: 005. doi:10.4172/2161-0495.S6-005

**Copyright:** © 2013 El-Ansary A, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

relationship between the intestinal bacteria and their hosts which could help to suggest potential therapeutic strategies through affecting the composition of gut microbiota.

This information initiate our interest to review all studies related to the role of gut microbiota, gut-brain axis and microbiome-host interaction in the aetiology of different neurodevelopmental disorders with special reference to autism. Understanding these aspects could help in early diagnosis, treatment or prevention of neurodevelopmental disorders.

## Gut Microbiota

### Normal gut microbiota

Bacterial species diversity in the gut largely derives from colonic transit time and the availability of different carbon substrates and energy sources which is the reason for the marked differences in species-level diversity found between individuals. However, the core members of microbiota are presented by, *Ruminococcus*, *Eubacterium* and *Dorea* (phylum Firmicutes); *Bacteroides* and *Alistipes* (phylum Bacteroidetes); and *Bifidobacterium* (phylum Actinobacteria) [12]. The composition and metabolic activities of majority of these members of gut microbiota depend on carbohydrate availability as the main nutritional factor and thus utilize saccharolytic metabolisms as the predominant pathway [13,14]. Most (95%) of the Firmicutes sequences examined by were members of the Clostridia class, which contains a substantial number of butyrate- producing bacteria that compose the clostridial clusters IV, XIVa and XVI [15]. *Roseburia intestinalis* and *Eubacterium rectale* have been reported to play dominant roles in butyrate synthesis, which is essential for the maintenance and protection of the normal colonic epithelium, whereas another butyrate producer, *Faecalibacterium prausnitzii*, is only weakly correlated with fecal butyrate concentrations [16]. On the other hand and differently, *Bifidobacterium* has been reported to produce lactate and acetate whereas *R. bromii* produces acetate, ethanol and hydrogen [17]. Schwiertz et al. [18] analyzed the fecal Short Chain Fatty Acid (SCFA) concentrations of lean and obese individuals and reported a 20% higher level of SCFAs in obese individuals, with the largest increase in propionate (41%), followed by butyrate (29%).

Moreover, Zhang et al. [19] hypothesized that in the gastrointestinal tracts of obese individuals, the coexistence of hydrogen-producing bacteria with relatively high numbers of hydrogen-utilizing methanogenic archaea could lead to an interspecies hydrogen transfer between bacterial and archaeal species. This may force the large intestine to an increase in the energy uptake in these individuals. It should be noted, however, that the “energy harvest” hypothesis suggests a protective effect of high intakes of dietary fibres (the main source of SCFAs) for enhancing weight loss or maintenance of a healthier body and thus reduces obesity.

### At-birth gut microbiota

Microbial colonization commences immediately after birth, and all infants are initially colonized by *Escherichia coli* and streptococci. The anaerobic genera *Bacteroides*, *Bifidobacterium* and *Clostridium* are established by the end of the first week of life. During the first months and years of life the neonatal gastrointestinal tract is colonized with an adult-type pattern of indigenous gut microflora finally comprising approximately  $10^{14}$  microorganisms, that is 10 times more than the number of eukaryotic cells in the adult body [20]. The development of the neonates gut microbiota is also controlled and modulated by different interacting mechanisms such as, genetic endowment, intrinsic

biological regulatory functions, environment influences and last but not least, the diet influence. Considered together with other endogenous and exogenous factors the type of feeding may interfere greatly in the regulation of the intestinal microbiota. The bacterial microbiota differs among formula-fed and breast-fed infants. In the former *Atopobium spp.* was found in significant counts and the numbers of *Bifidobacterium* dropped followed by increasing numbers in *Bacteroides* population. Moreover, under formula feeding the infants microbiota was more diverse [21]. Breast-fed infants harbour a fecal microbiota by more than two times increase in numbers of *Bifidobacterium* cells and also lacobacilli when compared to formula-fed infants [22]. The gut microbiota including *Bifidobacteria* constantly helps in successful maturation of the gut mucosal adaptive immune system [23-25].

It was previously reported that the mode of delivery strongly influences microbial colonization of infants including the gut [26]. Vaginally delivered infants acquire bacterial communities resembling mother's vaginal microbiota dominated by *Lactobacillus*, *Prevotella*, or *Sneathia spp.*, while caesarean delivery (C-section) infants harbor bacterial communities similar to those found on skin, dominated by *Staph.*, *Corynebacteria*, and *Propionibacterium spp.* In addition, the mode of delivery may have, possibly via gut microbiota development, significant effects on immunological functions in the infant since the total number of immunoglobulins IgA-, IgG- and IgM-secreting cells was found to be lower in infants born by vaginal delivery than in those born by C-section, possibly reflecting excessive antigen exposure across the vulnerable gut barrier. Also autism risk is influenced by the mode of delivery; a previous study has shown that C-section may double the risk of autism [27].

Furthermore, the size of healthy neonates vaginally born at term greatly affects the composition of gut microbiota and in turn the development of the immune system. The prevalence of Gram-negative Proteobacteria was higher in neonates born with Large Gestational Age (LGA), whereas Gram-positive Firmicutes was more prevalent in neonates born with appropriate gestational age (AGA). For this reason, appropriate care with pregnant woman and newborns should be considered as a preventive strategy of children diseases [28].

**Functions of human milk bacteria in the infant gut:** Human milk bacteria play a vital role in reducing incidence of infection breast-fed infants. This may occur by different mechanisms such as improvement of the intestinal barrier function by increasing mucine production and reducing intestinal permeability, competitive exclusion [29], or production of antimicrobial compounds [30-32]. The role of different bacterial strains in milk was previously reported. In this connection, administration of a human milk *Lactobacillus* strain to infants during 6 months led to 46%, 27%, and 30% reductions in the incidence rates of gastrointestinal infections, upper respiratory tract infections, and total number of infections, respectively [33]. Hospital environment resulting in undesired pathogens to infants or oral colonization by methicillin-resistant *S. aureus* in high-risk newborns may be inhibited by commensal coagulase-negative staphylococci and viridans streptococci provided by breast milk [34]. In fact, some *Staphylococcus epidermidis* strains that play such role have been postulated as a future strategy to eradicate such pathogens from the mucosal surfaces [35,36]. Breast milk bacteria may also participate in the correct maturation of the infant immune system since it was previously reported that some strains are able to modulate both natural and acquired immune responses in mice and humans with flexibility depending on the conditions found in the gut environment [37-39]. As an example, *Lactobacillus salivarius* CECT 5713 and *Lactobacillus fermentum* CECT 5716 enhanced macrophage production of Th1 cytokines, such as IL-2 and IL-12 and

the inflammatory mediator TNF-  $\alpha$ , in the absence of an inflammatory stimulus.

The glycobiome of some lactobacilli and bifidobacteria, including those of species isolated from human milk, may help to achieve a specific “healthy” microbiota in the infant gut [40,41]. These microorganisms are metabolically active in the infant gut by increasing the production of functional metabolites such as butyrate, which is the main energy source for colonocytes and a relevant compound in the modulation of intestinal function through the breakdown of sugars and proteins [42,43]. Taking in account that transit of food through the gastrointestinal tract is shorter in infants than in adults and, that the pH of the infant’s stomach is higher than that of the adult, human milk lactobacilli strains may improve the intestinal habit, with an increase in fecal moisture, and in stool frequency and volume.

### Development of the Microbiome

Diversity in the Gastrointestinal (GI) bacterial strains increases rapidly over the first few years of life [44,45]. The relatively few species GI strains that are first detected in infants, acquired from the mothers’ vagina and skin, are replaced by other strains of less certain origin [46-48]. However, the reason for this diversity is unknown: it is possible that new bacteria are incorporated at a constant rate as they are experienced in the environment, or that growing a larger gastrointestinal tract provide more distinct niches for bacteria, or a larger habitat for them to live in. Another alternative is that increasing functional complexity produces taxonomic complexity, until states of equilibrium are reached. Even though, it was found that within a single baby, the consortia of bacterial taxa is not random, at any given time point, indicating that the microbes depend on each other within the consortium. Therefore, during infancy groups of microbes rapidly colonize and may change in response to events such as illness [45]. This pattern of microbial diversity provides an efficient means for adaptation to the changing circumstances of development over an individual’s lifetime such as changes in lifestyle, illness, puberty, and others. Interestingly, human family members tend to have more similar microbiota.

Due to the function of gut flora in promoting normal gastrointestinal function, protecting from infection, regulating metabolism and comprising more than 75% of our immune system, so, dysregulated gut flora has been linked to diseases ranging from autism and depression to autoimmune conditions like Hashimoto’s, inflammatory bowel disease and type 1 diabetes. This probably explains why babies born via caesarean sections may have increased susceptibility to gut infections, asthma and allergies later in life [49].

### Factors affecting gut microbiota during development

The diversity in microbiota among children was shown in a recent study which compared the fecal microbiota of European children (EU) with that of African children from Burkina Faso (BF) in Central Africa. The results revealed significant differences in both biodiversity and richness of microbiota to the favour of BF children ( $P < 0.01$ ) [50]. African children fed on high carbohydrates and low protein diet, showed significantly higher levels of SCFAs ( $P < 0.001$ ) and a significant enrichment in Bacteroidetes and depletion in Firmicutes ( $P < 0.001$ ), with a unique abundance of bacteria from the genus *Prevotella* and *Xylanibacter* which were completely lacking in the EU children while *Enterobacteriaceae* (Shigella and Escherichia) were significantly lacking in African compared to EU children ( $P < 0.05$ ) [50]. Of somewhat greater surprise was the observation, that Gram-negative bacteria (mainly Bacteroidetes) were more abundant (58.5%) than Gram-positive bacteria (37.4%) in the BF population, whereas Gram-positive

(mainly Firmicutes) were more abundant than Gram-negative bacteria (70.4% vs 29.1% respectively) in the EU population, resulting in a Gram-positive to Gram negative ratio of 37 to 59 in the BF population compared to 70 to 29 in the EU population [50]. These observations regarding the effect of diet on gut microbiota was supported by Wu et al. [51] who investigated the association between dietary variables and gut microbiota in 98 individuals and demonstrated a strong correlation between long-term diet and enterotype. The Bacteroides enterotype was highly associated with the intake of animal protein and saturated fats, suggesting that meat consumption, as typified by a Western diet, characterize this enterotype. This could help to suggest that early dietary and gut microbiological environments have a more complex effect on the metabolic programming of a child than previously anticipated.

It was documented that obesity is greatly contributed the shift of children gut microbiota towards pathogenic composition. In a recent study done by Karlsson et al. [52], twenty 4–5 year old overweight or obese children were compared to twenty children of the same age but with normal body mass index. The burden of the Gram-negative family Enterobacteriaceae was significantly higher in the obese/overweight children and the levels of *Desulfovibrio* and *Akkermansia muciniphila*-like bacteria were significantly lower in the obese/overweight children. No significant differences were found in content of Lactobacillus, Bifidobacterium or the *Bacteroides fragilis* group. It was also observed that the diversity of the dominating bacterial community tended to be less diverse in the obese/overweight group, although the difference was not statistically significant.

A previous study has shown for the first time in human that differences in the gut microbiota may precede overweight development [53]. It was shown that *Bifidobacterium spp.* number was higher in children who exhibited a normal-weight at seven years than in children developing overweight. More importantly they observed that the *Staphylococcus aureus* counting was lower in children who maintain a normal-weight than in children becoming overweight several years later. This could provide evidence that the gut microbiota composition in children could be associated with weight gain and point out the putative role of the Bifidobacteria and Staphylococcus in that context. This is consistent with the recent finding of Barros et al. [54] showing 58% higher prevalence of obesity in young adult Brazilians born by CS than in young adults born vaginally. Because CS-born individuals do not make contact at birth with maternal vaginal and intestinal bacteria, this could lead to long-term changes in the gut microbiota that could contribute to obesity. The size of an infant at birth, a measure of gestational growth, has been recognized for many years as a biomarker of future risk of morbidity. Both being born Small for Gestational Age (SGA) and being born Large for Gestational Age (LGA), are associated with increased rates of obesity and metabolic disorder, as well as a number of mental disorders including attention deficit/hyperactivity disorder, autism, anxiety, and depression [55]. This could be related to the transfer of altered microbiota from pregnant mothers to infants which lead to an increased risk of abnormal gestational weight [56], and thus the composition and development of infant gut microbiota are influenced by Body Mass Index (BMI), weight, and weight gain of mothers during pregnancy.

It could be suggested that a balance between microbial groups present in the human gut is crucial for maintaining health. When this balance is disturbed, the host microbe relationship can progress toward a disease state. Altered intestinal colonization by commensal microorganisms as well as high inter-individual variability and reduced microbial diversity has been reported in preterm infants increasing the risk to develop later disease [57,58].



## Gut –brain axis and aetiology of neuro-developmental disorders

It is well known that gut microbiota can affect the development [59] and function [60] of the central nervous system, thereby, leading to the recent interesting concept of the microbiota gut–brain axis [61] (Figure 1).

Many studies using animal models of different behavioural disorders such as autism, anxiety, cognitive disability and depression proved that microbiota composition greatly influences brain function. Neuroactive compounds in the intestinal lumen can cross the blood-brain barrier and induce many cognitive and behavioural disturbances [62].

The composition of the intestinal microbiota is extremely relevant in neurogastroenterology, as a science deals with the gut–brain axis interactions. Several neuropathological diseases are thought to be associated with the gut microbiota. Autism as a neurodevelopmental disorder often involves GI symptoms. Recent studies related to faecal microbial profiles of autistic patients, indicated 10-fold higher counts of *Clostridium* spp. compared with healthy controls [63]. *Clostridium* is known to produce neurotoxins, which could contribute to the development of autistic behaviours. Higher urinary levels of hippurate, phenylacetylglutamine and tryptophan/nicotinic acid metabolism have been reported in autistic children as an aspect of metabolic alteration in gut host–microbial co-metabolism [64,65].

Although many studies have demonstrated altered gut microbiota composition in children with autism compared with control healthy subjects [66-70], such data should be interpreted with care, as autistic patients have a higher incidence of antibiotic usage and often have different diets compared with neurotypical individuals, both of which can alter the composition of the gut microbiota. Interestingly, a recent study also highlights alterations in the faecal concentrations of the short-chain fatty acids in children with autism [71] suggesting that production of such neuroactive microbial metabolites could be related to the mechanism by which bacteria may alter brain function.

Recently, intracerebroventricular or oral administration of neurotoxic doses of Propionic Acid (PPA) to animals was effective in inducing autistic features [72,73]. It is currently unclear whether the doses of propionic acid used in animal studies reflect the potential alterations in short-chain fatty acids observed in autistic individuals [74].

Interestingly, there has been some transient success in using the


antibiotic vancomycin in treating some of the symptoms of autism [75]. Although such studies are effective, it needs replication in a greater numbers of patients and controlled clinical trials using more sophisticated bacterial analyses are recommended to assess whether autism is associated with alterations in the gut microbiota and whether such alterations play a part in the gastrointestinal, behavioural and cognitive symptoms seen in autistic children.

Recently, there is a growing interest suggesting that dietary factors might worsen and, in some cases, improve the symptoms of autism. It is well known that SCFAs, such as PPA, are produced by many intestinal bacteria through the breakdown of dietary carbohydrates and amino acids [76]. Special attention is given to *Clostridia* species as the most infectious causes of ASDs [77]. *Clostridia* species, as anaerobic, gram-positive and PPA producers [78], are major bacteria that colonize the gut in early life. It is well documented that spore-forming anaerobes and microaerophilic bacteria, particularly from *Clostridia* species, are elevated in patients with autism [79].

Additionally, species of *Desulfovibrio*, a gram-negative, non-spore former were recently isolated from the stool of patients with autism, and, to a lesser extent, non-affected siblings. *Desulfovibrio*, in addition to PPA production, is resistant to most common antibiotics and produces the gasotransmitter and potential mitochondrial toxin, hydrogen sulfide. Eradication of these organisms with oritavancin and aztreonam was recently suggested as a possible treatment of ASDs.

Furthermore, ASDs often show comorbidity with a variety of gastrointestinal disorders, such as alterations in gut motility, leaky gut, bacterial dysbiosis, impaired carbohydrate digestion/ absorption, reflux esophagitis [80,81,66]. An association between long-term antibiotic use, hospitalization, abdominal discomfort and the onset of ASD symptoms after normal or near-normal development has also been reported [82-84]. These findings raise the possibility that gut-born factors secondary to alteration of the gut microbiome by antibiotics or diet may affect brain function in patients with autism. Moreover, a compromised gut–blood barrier in case of acquired colitis or impaired colonocyte energy metabolism [85], which use SCFAs as an energy substrate may contribute for greater systemic and brain access for PPA. PPA is also known to have a number of direct effects on gut physiology. As reviewed by MacFabe et al. [80], PPA increases the contraction of colonic smooth muscle, dilates colonic arteries, and increases serotonin release from gut chromaffin cells, and decrease gastric motility, which could be easily related to the gastrointestinal abnormalities frequently observed in many autistic patients. This could explain the observations of some parents of autism that gastrointestinal and behavioural symptoms increase when their children fed high carbohydrate diet or any food that contain PPA as preservative or eradication of PPA-producing bacteria using broad spectrum antibiotics [84,86].

In a recent study done by El-Ansary et al. [73], orally administered PPA was highly potent to induce oxidative stress (lipid peroxidation), coupled with a decrease in Glutathione (GSH) and Glutathione Peroxidase (GPX) and catalase activities. Impaired energy metabolism was also ascertained through the decrease of lactate dehydrogenase and activation of Creatine Kinase (CK). Elevated IL-6, TNF $\alpha$ , IFN $\gamma$  and heat shock protein 70 (HSP70) confirmed the neuroinflammatory effect of PPA. Moreover, elevation of caspase3 and DNA fragmentation proved the pro-apoptotic and neurotoxic effect of PPA to rat pups received 250mg/kg body weight for 3 days. Their study proved the involvement of PPA in inducing persistent autistic features in rat pups. In fact, El-Ansary et al. [74] previously provided plausible links that related the occurrence of lower PPA in the plasma of autistic patients to elevated



levels of PA in their brain. They attributed the remarkably lower plasma PPA in autistic patients to the high rate of blood to brain influx. In fact, and compared to other fatty acids, PPA was previously reported to cross the Blood Brain Barrier (BBB) with a brain uptake index of 43.53 and a low Km value of 2.03 [87]. Since the lower the Km, the higher the affinity of the transporters for the substrates, then an uptake index of 43.53% and a Km value of 2.03 are enough to facilitate the cross of PPA into the brain cell, which could explain the elevation of this SCFA in the brain homogenates of the treated rats.

In an attempt to prove the relationship between unbalanced gut microbiota and the etiology of autistic features and to confirm the critical role of *Clostridium difficile* as PPA producer, a comparative study of the effect of clindamycin-induced *Clostridium difficile* growth and orally administered PPA was done by El-Ansary et al. [88]. Both treatments were effective to induce biochemical autistic features (Oxidative stress, mitochondrial dysfunction, neuroinflammation, pro-apoptotic) with direct orally administered PPA being more potent compared to the indirect effect, through induction of PPA bacterial producers among which is *Clostridium difficile*.

Clearly, gut microbiota not only exert a local effect on the GI tract but also impact remote organs such as the brain through chemical signaling.

## Treatment Strategy to Reduce *Clostridium Difficile*

### Probiotics and prebiotics

The antibiotic-associated diarrhoea is mostly due to *C. difficile*, pathogenic bacteria recently reported as etiological factor in the pathophysiology of autism [70]. A randomised double-blind placebo-controlled trial done by Hickson et al. [89] recorded that consumption of a probiotic drink containing *L. casei*, *L. bulgaricus*, and *S. thermophilus* can reduce *C. difficile* associated Diarrhoea (CDD). Among the randomized patients, 138 received the Lactobacillus and Bifidobacterium strains as probiotic in combination with the antibiotic and the other half received the antibiotic therapy alone for 20 days compared with placebo group. On basis of diarrhoea development, 2.9% of patients' present *C. difficile* associated toxins in their faecal samples versus 7.9% in placebo control. After complete analysis of patient samples, 46% of probiotic patients were toxin-positive compared with 78% of the placebo group. Based on these records, probiotics could be suggested as treatment strategy for autistic patients.

A prebiotic is defined as selectively fermented ingredients that induce specific changes, both in the composition and/or activity in the gut microbiota that confers benefits upon host health [90,91]. In recent years, there is a dramatically increasing interest in the use of prebiotics as functional foods in order to modulate the composition of gut microbiota [92,93].

### Faecal transplantation

One of the most important techniques recently considered in treating *C. difficile* infection is faecal transplantation. This treatment strategy aims to replace the gut microbiota of a diseased individual by transplanting the microbiota from a healthy donor [94]. Meta analyses have recently reported a 90% successful trials when faecal transplantation is used to treat refractory *C. difficile* infection [95,96] showing that this methodology has potent and reproducible efficacy when broad-spectrum antibiotics, as traditional therapeutic option have failed to treat disease [95]. Recent studies, certainly show that faecal transplants can be effective even when samples that have been previously frozen were used or when the transplant is self-administered

suggest that it will be possible to simplify donor recruitment and sample processing steps without reducing the potency [97,98].

The mechanism of action of faecal transplantation has not been established. However, patients with recurrent *C. difficile* Infection (CDI) have been found to have decreased bacterial diversity in their stool microbiome [99,100]. By repopulating the gastrointestinal tract with a healthy microbiome, stool transplantation could be effective in restoring resistance to *C. difficile* growth [95,101]. Although fecal transplantation is considered as successful strategy to treat dysbiosis, but it is not widely used because of the time required to identify a suitable donor, the risk of introducing pathogenic bacteria, and a general recipient dislike [102]. Thus, the development of animal model that have many features of fecal transplantation in humans with recurrent *C. difficile* disease could help to understand the basic mechanisms of successful fecal transplantation and also to develop standardized bacteriotherapy [103].

### Hyperbaric Oxygen Therapy (HBOT)

HBOT has been used to decrease the amount of abnormal bacteria in the gut and therefore can function as an antibiotic [104]. In animal studies, HBOT was effective in reducing intestinal bacterial counts after bacteria overgrowth in the distal ileum associated with bile duct ligation [105]. It also shows bactericidal activity against many pathogenic bacteria, including *Pseudomonas* [106] *Salmonella* and *Proteus*, *Staphylococcus* [107], *Mycobacterium tuberculosis* [108], and anaerobic bacteria such as *Clostridia* [109].

Based on the fact that oxygen-dependent killing of *Staphylococcus aureus* by phagocytic leukocytes has been shown to increase by HBOT in animals [110], and that HBOT has also been shown to inhibit the growth of some yeast [111] and to possess virucidal activity against some enveloped viruses [112], HBOT might lead to an improvement in the dysbiosis found in some autistic patients by reducing counts of abnormal pathogens. However, many of the studies had limitations which may have contributed to inconsistent findings across them, including the use of many different standardized and non-standardized instruments, making it difficult to directly compare the results of studies or to know if there are specific areas of behaviour in which HBOT is most effective [113].

In a recent study done by Chiranjit et al. [114], use of HBOT for children appears generally safe, even at pressures up to 2.0 atm for 2 h per day for 40 sessions the atmospheric pressure has a significant impact on the bacterial colonization of the gut and on the ecology of the gut microflora.

## Conclusion

The gut microbiota, gut-brain axis and microbiome-host interaction play a significant role in aetiology of different neurodevelopmental disorders, especially autism.

Microbial colonization commences immediately after birth and carbohydrate availability is the most important nutritional factor which could control the composition and metabolic activities of microbiota and bacterial species diversity.

We can hypothesize that understanding these aspects could help in early diagnosis, treatment or prevention of neurodevelopmental disorders such as autism.

## References

1. Rossignol DA (2009) Novel and emerging treatments for autism spectrum disorders: a systematic review. *Ann Clin Psychiatry* 21: 213-236.

2. Jusko TA, Henderson CR, Lanphear BP, Cory-Slechta DA, Parsons PJ, et al. (2008) Blood lead concentrations < 10 microg/dL and child intelligence at 6 years of age. *Environ Health Perspect* 116: 243-248.
3. Lanphear BP, Hornung R, Khoury J, Yolton K, Baghurst P, et al. (2005) Low-level environmental lead exposure and children's intellectual function: an international pooled analysis. *Environmental Health Perspectives* 113: 894-899.
4. Schnaas L, Rothenberg SJ, Flores MF, Martinez S, Hernandez C, et al. (2006) Reduced intellectual development in children with prenatal lead exposure. *Environ Health Perspect* 114: 791-797.
5. Harada M, Akagi H, Tsuda T, Kizaki T, Ohno H (1999) Methylmercury level in umbilical cords from patients with congenital Minamata disease. *Sci Total Environ* 234: 59-62.
6. Jacobson JL, Jacobson SW (2003) Prenatal exposure to polychlorinated biphenyls and attention at school age. *J Pediatr* 143: 780-788.
7. Murray RM, Lewis SW (1987) Is schizophrenia a neurodevelopmental disorder? *Br Med J (Clin Res Ed)* 295: 681-682.
8. Schuurmans C, Kurrasch DM (2013) Neurodevelopmental consequences of maternal distress: what do we really know? *Clin Genet* 83: 108-117.
9. Kinney DK, Hintz K, Shearer EM, Barch DH, Riffin C, et al. (2010) A unifying hypothesis of schizophrenia: abnormal immune system development may help explain roles of prenatal hazards, post-pubertal onset, stress, genes, climate, infections, and brain dysfunction. *Med Hypotheses* 74: 555-63.
10. Ferguson DJ, Bowker C, Jeffery KJ, Chamberlain P, Squier W (2013) Congenital toxoplasmosis: continued parasite proliferation in the fetal brain despite maternal immunological control in other tissues. *Clin Infect Dis* 56: 204-208.
11. Hsiao EY, Patterson PH (2012) Placental regulation of maternal-fetal interactions and brain development. *Dev Neurobiol* 72: 1317-1326.
12. Tap J, Mondot S, Levenez F, Pelletier E, Caron C, et al. (2009) Towards the human intestinal microbiota phylogenetic core. *Environ Microbiol* 11: 2574-2584.
13. Macfarlane GT, Macfarlane S (2011) Fermentation in the human large intestine: its physiologic consequences and the potential contribution of prebiotics. *J Clin Gastroenterol* 45 Suppl: S120-127.
14. Smith AR, Macfarlane S, Furrie E, Ahmed S, Bahrami B, et al. (2011) Microbiological and immunological effects of enteral feeding on the upper gastrointestinal tract. *J Med Microbiol* 60: 359-365.
15. Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, et al. (2005) Diversity of the human intestinal microbial flora. *Science* 308: 1635-1638.
16. Duncan SH, Belenguer A, Holtrop G, Johnstone AM, Flint HJ, et al. (2007) Reduced dietary intake of carbohydrates by obese subjects results in decreased concentrations of butyrate and butyrate-producing bacteria in feces. *Appl Environ Microbiol* 73: 1073-1078.
17. Furrie E, Macfarlane S, Kennedy A, Cummings JH, Walsh SV, et al. (2005) Synbiotic therapy (Bifidobacterium longum/Synergy 1) initiates resolution of inflammation in patients with active ulcerative colitis: a randomised controlled pilot trial. *Gut* 54: 242-249.
18. Schwierdt A, Taras D, Schäfer K, Beijer S, Bos NA, et al. (2010) Microbiota and SCFA in lean and overweight healthy subjects. *Obesity (Silver Spring)* 18: 190-195.
19. Zhang H, DiBaise JK, Zuccolo A, Kudrna D, Braidotti M, et al. (2009) Human gut microbiota in obesity and after gastric bypass. *Proc Natl Acad Sci U S A* 106: 2365-2370.
20. Berg RD (1996) The indigenous gastrointestinal microflora. *Trends Microbiol* 4: 430-435.
21. Bezirtzoglou E, Tsiotsias A, Welling GW (2011) Microbiota profile in feces of breast- and formula-fed newborns by using fluorescence in situ hybridization (FISH). *Anaerobe* 17: 478-482.
22. Harmsen HJ, Wildeboer-Veloo AC, Raangs GC, Wagendorp AA, Klijn N, et al. (2000) Analysis of intestinal flora development in breast-fed and formula-fed infants by using molecular identification and detection methods. *J Pediatr Gastroenterol Nutr* 30: 61-67.
23. Brandtzaeg P (1996) Development of the mucosal immune system in humans. In: Bindels JG, Goedhart AC, Visser HKA (editors) *Recent developments in infant nutrition*. 1. UK: Kluwer Academic Publishers 349-376.
24. Avershina E, Storrø O, Øien T, Johnsen R, Wilson R, et al. (2013) Bifidobacterial succession and correlation networks in a large unselected cohort of mothers and their children. *Appl Environ Microbiol* 79: 497-507.
25. Turroni F, Peano C, Pass DA, Foroni E, Severgnini M, et al. (2012) Diversity of bifidobacteria within the infant gut microbiota. *PLoS One* 7: e36957.
26. Dominguez-Bello MG, Costello EK, Contreras M, Magris M, Hidalgo G, et al. (2010) Delivery mode shapes the acquisition and structure of the initial microbiota across multiple body habitats in newborns. *Proc Natl Acad Sci U S A* 107: 11971-11975.
27. Glasson EJ, Bower C, Petterson B, de Klerk N, Chaney G, et al. (2004) Perinatal factors and the development of autism: a population study. *Arch Gen Psychiatry* 61: 618-627.
28. Karlsson CL, Molin G, Cilio CM, Ahrné S (2011) The pioneer gut microbiota in human neonates vaginally born at term-a pilot study. *Pediatr Res* 70: 282-286.
29. Olivares M, Díaz-Ropero MP, Martín R, Rodríguez JM, Xaus J (2006) Antimicrobial potential of four Lactobacillus strains isolated from breast milk. *J Appl Microbiol* 101: 72-79.
30. Beasley SS, Saris PE (2004) Nisin-producing Lactococcus lactis strains isolated from human milk. *Appl Environ Microbiol* 70: 5051-5053.
31. Martín R, Olivares M, Marín ML, Fernández L, Xaus J, et al. (2005) Probiotic potential of 3 Lactobacilli strains isolated from breast milk. *J Hum Lact* 21: 8-17.
32. Martín R, Jiménez E, Olivares M, Marín ML, Fernández L, et al. (2006) Lactobacillus salivarius CECT 5713, a potential probiotic strain isolated from infant feces and breast milk of a mother-child pair. *Int J Food Microbiol* 112: 35-43.
33. Maldonado J, Cañabate F, Sempere L, Vela F, Sánchez AR, et al. (2012) Human milk probiotic Lactobacillus fermentum CECT5716 reduces the incidence of gastrointestinal and upper respiratory tract infections in infants. *J Pediatr Gastroenterol Nutr* 54: 55-61.
34. Uehara Y, Kikuchi K, Nakamura T, Nakama H, Agematsu K, et al. (2001) H2O2 produced by viridans group streptococci may contribute to inhibition of methicillin-resistant Staphylococcus aureus colonization of oral cavities in newborns. *Clinical Infectious Diseases* 32: 1408-1413.
35. Iwase T, Uehara Y, Shinji H, Tajima A, Seo H, et al. (2010) Staphylococcus epidermidis Esp inhibits Staphylococcus aureus biofilm formation and nasal colonization. *Nature* 465: 346-349.
36. Park B, Iwase T, Liu GY (2011) Intranasal application of S. epidermidis prevents colonization by methicillin-resistant Staphylococcus aureus in mice. *PLoS One* 6: e25880.
37. Díaz-Ropero MP, Martín R, Sierra S, Lara-Villoslada F, Rodríguez JM, et al. (2007) Two Lactobacillus strains, isolated from breast milk, differently modulate the immune response. *J Appl Microbiol* 102: 337-343.
38. Olivares M, Díaz-Ropero MP, Gómez N, Lara-Villoslada F, Sierra S, et al. (2006) The consumption of two new probiotic strains, Lactobacillus gasseri CECT 5714 and Lactobacillus coryniformis CECT 5711, boosts the immune system of healthy humans. *Int Microbiol* 9: 47-52.
39. Olivares M, Díaz-Ropero MP, Sierra S, Lara-Villoslada F, Fonollá J, et al. (2007) Oral intake of Lactobacillus fermentum CECT5716 enhances the effects of influenza vaccination. *Nutrition* 23: 254-260.
40. Zivkovic AM, German JB, Lebrilla CB, Mills DA (2011) Human milk glycomiome and its impact on the infant gastrointestinal microbiota. *Proc Natl Acad Sci U S A* 108 Suppl 1: 4653-4658.
41. Asakuma S, Hatakeyama E, Urashima T, Yoshida E, Katayama T, et al. (2011) Physiology of consumption of human milk oligosaccharides by infant gut-associated bifidobacteria. *J Biol Chem* 286: 34583-34592.
42. Maldonado J, Lara-Villoslada F, Sierra S, Sempere L, Gómez M, et al. (2010) Safety and tolerance of the human milk probiotic strain Lactobacillus salivarius CECT5713 in 6-month-old children. *Nutrition* 26: 1082-1087.
43. Gil-Campos M, López MÁ, Rodríguez-Benítez MV, Romero J, Roncero I, et al. (2012) Lactobacillus fermentum CECT 5716 is safe and well tolerated in infants of 1-6 months of age: a randomized controlled trial. *Pharmacol Res* 65: 231-238.



44. Palmer C, Bik EM, DiGiulio DB, Relman DA, Brown PO (2007) Development of the human infant intestinal microbiota. *PLoS Biol* 5: e177.
45. Koenig JE, Spor A, Scalfone N, Fricker AD, Stombaugh J, et al. (2011) Succession of microbial consortia in the developing infant gut microbiome. *Proc Natl Acad Sci U S A* 108 Suppl 1: 4578-4585.
46. Matsumiya Y, Kato N, Watanabe K, Kato H, et al. (2002) Molecular epidemiological study of vertical transmission of vaginal *Lactobacillus* species from mothers to newborn infants in Japanese, by arbitrarily primed polymerase chain reaction. *J Infect Chemother* 8: 43-49.
47. Tannock GW, Fuller R, Smith SL, Hall MA (1990) Plasmid profiling of members of the family Enterobacteriaceae, lactobacilli, and bifidobacteria to study the transmission of bacteria from mother to infant. *J Clin Microbiol* 28: 1225-1228.
48. Vaishampayan PA, Kuehl JV, Froula JL, Morgan JL, Ochman H, et al. (2010) Comparative metagenomics and population dynamics of the gut microbiota in mother and infant. *Genome Biol Evol* 2: 53-66.
49. Chris K (2012) Natural Childbirth series, Natural Childbirth VII: Caesarean Risks and Complications.
50. De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, et al. (2010) Impact of diet in shaping gut microbiota revealed by a comparative study in children from Europe and rural Africa. *Proc Natl Acad Sci U S A* 107: 14691-14696.
51. Wu GD, Chen J, Hoffmann C, Bittinger K, Chen YY, et al. (2011) Linking long-term dietary patterns with gut microbial enterotypes. *Science* 334: 105-108.
52. Karlsson CL, Onnerfält J, Xu J, Molin G, Åhrné S, et al. (2012) The microbiota of the gut in preschool children with normal and excessive body weight. *Obesity (Silver Spring)* 20: 2257-2261.
53. Kalliomäki M, Collado MC, Salminen S, Isolauri E (2008) Early differences in fecal microbiota composition in children may predict overweight. *Am J Clin Nutr* 87: 534-538.
54. Barros FC, Matijasevich A, Hallal PC, Horta BL, Barros AJ, et al. (2012) Cesarean section and risk of obesity in childhood, adolescence, and early adulthood: evidence from 3 Brazilian birth cohorts. *Am J Clin Nutr* 95: 465-470.
55. Grissom NM, Reyes TM (2012) Gestational overgrowth and undergrowth affect neurodevelopment: similarities and differences from behavior to epigenetics. *Int J Dev Neurosci*.
56. Collado MC, Isolauri E, Laitinen K, Salminen S (2010) Effect of mother's weight on infant's microbiota acquisition, composition, and activity during early infancy: a prospective follow-up study initiated in early pregnancy. *Am J Clin Nutr* 92: 1023-1030.
57. Jacquot A, Neveu D, Aujoulat F, Mercier G, Marchandin H, et al. (2011) Dynamics and clinical evolution of bacterial gut microflora in extremely premature patients. *J Pediatr* 158: 390-396.
58. Rougé C, Goldenberg O, Ferraris L, Berger B, Rochat F, et al. (2010) Investigation of the intestinal microbiota in preterm infants using different methods. *Anaerobe* 16: 362-370.
59. Diaz Heijtz R, Wang S, Anuar F, Qian Y, Björkholm B, et al. (2011) Normal gut microbiota modulates brain development and behavior. *Proc Natl Acad Sci U S A* 108: 3047-3052.
60. Neufeld KM, Kang N, Bienenstock J, Foster JA (2011) Reduced anxiety-like behavior and central neurochemical change in germ-free mice. *Neurogastroenterol Motil* 23: 255-264, e119.
61. Cryan JF, O'Mahony SM (2011) The microbiome-gut-brain axis: from bowel to behavior. *Neurogastroenterol Motil* 23: 187-192.
62. Wakefield AJ (2002) The gut-brain axis in childhood developmental disorders. *J Pediatr Gastroenterol Nutr* 34 Suppl 1: S14-17.
63. Sekirov I, Russell SL, Antunes LC, Finlay BB (2010) Gut microbiota in health and disease. *Physiol Rev* 90: 859-904.
64. Yap IK, Angley M, Veselkov KA, Holmes E, Lindon JC, et al. (2010) Urinary metabolic phenotyping differentiates children with autism from their unaffected siblings and age-matched controls. *J Proteome Res* 9: 2996-3004.
65. de Theije CG, Wu J, da Silva SL, Kamphuis PJ, Garssen J, et al. (2011) Pathways underlying the gut-to-brain connection in autism spectrum disorders as future targets for disease management. *Eur J Pharmacol* 668 Suppl 1: S70-80.
66. Williams BL, Hornig M, Buie T, Bauman ML, Cho Paik M, et al. (2011) Impaired carbohydrate digestion and transport and mucosal dysbiosis in the intestines of children with autism and gastrointestinal disturbances. *PLoS One* 6: e24585.
67. Finegold SM, Molitoris D, Song Y, Liu C, Vaisanen ML, et al. (2002) Gastrointestinal microflora studies in late-onset autism. *Clin Infect Dis* 35: S6-S16.
68. Finegold SM, Dowd SE, Gontcharova V, Liu C, Henley KE, et al. (2010) Pyrosequencing study of fecal microflora of autistic and control children. *Anaerobe* 16: 444-453.
69. Parracho HM, Bingham MO, Gibson GR, McCartney AL (2005) Differences between the gut microflora of children with autistic spectrum disorders and that of healthy children. *J Med Microbiol* 54: 987-991.
70. Adams JB, Johansen LJ, Powell LD, Quig D, Rubin RA (2011) Gastrointestinal flora and gastrointestinal status in children with autism—comparisons to typical children and correlation with autism severity. *BMC Gastroenterol* 11: 22.
71. Wang L, Christophersen CT, Sorich MJ, Gerber JP, Angley MT, et al. (2012) Elevated fecal short chain fatty acid and ammonia concentrations in children with autism spectrum disorder. *Dig Dis Sci* 57: 2096-2102.
72. MacFabe DF, Cain NE, Boon F, Ossenkopp KP, Cain DP (2011) Effects of the enteric bacterial metabolic product propionic acid on object-directed behaviour, social behaviour, cognition, and neuro-inflammation in adolescent rats: relevance to autism spectrum disorder. *Behav Brain Res* 217: 47-54.
73. El-Ansary AK, Ben Bacha A, Kotb M (2012) Etiology of autistic features: the persisting neurotoxic effects of propionic acid. *J Neuroinflammation* 9: 74.
74. El-Ansary AK, Bacha AG, Al-Ayadhi LY (2011) Plasma fatty acids as diagnostic markers in autistic patients from Saudi Arabia. *Lipids Health Dis* 10: 62.
75. Sandler RH, Finegold SM, Bolte ER, Buchanan CP, Maxwell AP, et al. (2000) Short-term benefit from oral vancomycin treatment of regressive-onset autism. *J Child Neurol* 15: 429-435.
76. Haskå L, Andersson R, Nyman M (2011) The effect of dietary fiber from wheat processing streams on the formation of carboxylic acids and microbiota in the hindgut of rats. *J Agric Food Chem* 59: 3406-3413.
77. Finegold SM (2011) *Desulfovibrio* species are potentially important in regressive autism. *Med Hypotheses* 77: 270-274.
78. Stackebrandt E, Rainey FA (1997) Phylogenetic relationships In: Rood JI, McClane BA, Songer JG, Titball RW, eds. *The clostridia, molecular biology and pathogenesis*. New York, NY: Academic Press.
79. Finegold SM (2011) State of the art; microbiology in health and disease. *Intestinal bacterial flora in autism*. *Anaerobe* 17: 367-368.
80. MacFabe DF, Cain DP, Rodriguez-Capote K, Franklin AE, Hoffman JE, et al. (2007) Neurobiological effects of intraventricular propionic acid in rats: possible role of short chain fatty acids on the pathogenesis and characteristics of autism spectrum disorders. *Behav Brain Res* 176: 149-169.
81. White JF (2003) Intestinal pathophysiology in autism. *Exp Biol Med (Maywood)* 228: 639-649.
82. Atladóttir HO, Thorsen P, Østergaard L, Schendel DE, Lemcke S, et al. (2010) Maternal infection requiring hospitalization during pregnancy and autism spectrum disorders. *J Autism Dev Disord* 40: 1423-1430.
83. Atladóttir HO, Thorsen P, Schendel DE, Østergaard L, Lemcke S, et al. (2010) Association of hospitalization for infection in childhood with diagnosis of autism spectrum disorders: a Danish cohort study. *Arch Pediatr Adolesc Med* 164: 470-477.
84. Horvath K, Papadimitriou JC, Rabsztyrn A, Drachenberg C, Tildon JT (1999) Gastrointestinal abnormalities in children with autistic disorder. *J Pediatr* 135: 559-563.
85. Lagoutte E, Mimoun S, Andriamihaja M, Chaumontet C, Blachier F, et al. (2010) Oxidation of hydrogen sulfide remains a priority in mammalian cells and causes reverse electron transfer in colonocytes. *Biochim Biophys Acta* 1797: 1500-1511.
86. Mellon AF, Deshpande SA, Mathers JC, Bartlett K (2000) Effect of oral antibiotics on intestinal production of propionic acid. *Arch Dis Child* 82: 169-172.
87. Conn AR, Fell DI, Steele RD (1983) Characterization of alpha-keto acid transport across blood-brain barrier in rats. *Am J Physiol* 245: E253-260.
88. El-Ansary AK, Al-Daihan S, Ben Bacha A, Shaker GH, Al-Ayadhi LY (2013)

- Comparative study on the protective effect of carnosine and carnitine against pro-inflammatory/pro-oxidant effects of clindamycin and propionic acid administrations to hamsters. *African Journal of Microbiology Research* 7: 103-114.
89. Hickson M, D'Souza AL, Muthu N, Rogers TR, Want S, et al. (2007) Use of probiotic *Lactobacillus* preparation to prevent diarrhoea associated with antibiotics: randomised double blind placebo controlled trial. *BMJ* 335: 80.
  90. Gibson GR, Probert HM, Loo JV, Rastall RA, Roberfroid MB (2004) Dietary modulation of the human colonic microbiota: updating the concept of prebiotics. *Nutr Res Rev* 17: 259-275.
  91. Roberfroid M (2007) Prebiotics: the concept revisited. *J Nutr* 137: 830S-7S.
  92. Vardakou M, Palop CN, Christakopoulos P, Faulds CB, Gasson MA, et al. (2008) Evaluation of the prebiotic properties of wheat arabinoxylan fractions and induction of hydrolase activity in gut microflora. *Int J Food Microbiol* 123: 166-170.
  93. Silk DB, Davis A, Vulevic J, Tzortzis G, Gibson GR (2009) Clinical trial: the effects of a trans-galactooligosaccharide prebiotic on faecal microbiota and symptoms in irritable bowel syndrome. *Alimentary Pharmacology and Therapeutics* 29: 508-518.
  94. Palmer R (2011) Fecal matters. *Nat Med* 17: 150-152.
  95. Bakken JS (2009) Fecal bacteriotherapy for recurrent *Clostridium difficile* infection. *Anaerobe* 15: 285-289.
  96. Landy J, Al-Hassi HO, McLaughlin SD, Walker AW, Ciclitira PJ, et al. (2011) Review article: faecal transplantation therapy for gastrointestinal disease. *Aliment Pharmacol Ther* 34: 409-415.
  97. Silverman MS, Davis I, Pillai DR (2010) Success of self-administered home fecal transplantation for chronic *Clostridium difficile* infection. *Clin Gastroenterol Hepatol* 8: 471-473.
  98. Hamilton MJ, Weingarten AR, Sadowsky MJ, Khoruts A (2012) Standardized frozen preparation for transplantation of fecal microbiota for recurrent *Clostridium difficile* infection. *Am J Gastroenterol* 107: 761-767.
  99. Chang JY, Antonopoulos DA, Kalra A, Tonelli A, Khalife WT, et al. (2008) Decreased diversity of the fecal Microbiome in recurrent *Clostridium difficile*-associated diarrhea. *J Infect Dis* 197: 435-438.
  100. Rea MC, O'Sullivan O, Shanahan F, O'Toole PW, Stanton C, et al. (2012) *Clostridium difficile* carriage in elderly subjects and associated changes in the intestinal microbiota. *J Clin Microbiol* 50: 867-875.
  101. Brandt LJ, Aroniadis OC, Mellow M, Kanatzar A, Kelly C, et al. (2012) Long-term follow-up of colonoscopic fecal microbiota transplant for recurrent *Clostridium difficile* infection. *Am J Gastroenterol* 107: 1079-1087.
  102. Borody TJ, Khoruts A (2011) Fecal microbiota transplantation and emerging applications. *Nat Rev Gastroenterol Hepatol* 9: 88-96.
  103. Lawley TD, Clare S, Walker AW, Stares MD, Connor TR et al. (2012) Targeted Restoration of the Intestinal Microbiota with a Simple, Defined Bacteriotherapy Resolves Relapsing *Clostridium difficile* Disease in Mice *PLoS Pathog* 8: e1002995.
  104. Knighton DR, Halliday B, Hunt TK (1984) Oxygen as an antibiotic. The effect of inspired oxygen on infection. *Arch Surg* 119: 199-204.
  105. Akin ML, Erenoglu C, Dal A, Erdemoglu A, Elbuen E, et al. (2001) Hyperbaric oxygen prevents bacterial translocation in rats with obstructive jaundice. *Dig Dis Sci* 46: 1657-1662.
  106. Bornside GH, Pakman LM, Ordóñez AA Jr (1975) Inhibition of pathogenic enteric bacteria by hyperbaric oxygen: enhanced antibacterial activity in the absence of carbon dioxide. *Antimicrob Agents Chemother* 7: 682-687.
  107. Bornside GH (1967) Enhancement of antibiotic activity against *Staphylococcus aureus* by exposure to hyperbaric oxygen. *Appl Microbiol* 15: 1020-1024.
  108. Gottlieb SF (1971) Effect of hyperbaric oxygen on microorganisms. *Annu Rev Microbiol* 25: 111-152.
  109. Unsworth IP, Sharp PA (1984) Gas gangrene. An 11-year review of 73 cases managed with hyperbaric oxygen. *Med J Aust* 140: 256-260.
  110. Mader JT, Brown GL, Guckian JC, Wells CH, Reinartz JA (1980) A mechanism for the amelioration by hyperbaric oxygen of experimental staphylococcal osteomyelitis in rabbits. *J Infect Dis* 142: 915-922.
  111. Arao T, Hara Y, Suzuki Y, Tamura K (2005) Effect of high-pressure gas on yeast growth. *Biosci Biotechnol Biochem* 69: 1365-1371.
  112. Baugh MA (2000) HIV: reactive oxygen species, enveloped viruses and hyperbaric oxygen. *Med Hypotheses* 55: 232-238.
  113. Rossignol DA, Bradstreet JJ, Van Dyke K, Schneider C, Freedenfeld SH, et al. (2012) Hyperbaric oxygen treatment in autism spectrum disorders. *Med Gas Res* 2: 16.
  114. Chiranjit M, Atanu A, Tarun KP, Bikas RP, Kesha CM (2012) Study of the cultivable microflora of the large intestine of the rat under varied environmental hyperbaric pressures. *J Microbiol Immunol Infect* 45: 281-286.