

SARS-CoV-2 in Saliva May Be Deactivated with Fatty Acids or Emulsifiers

Daisuke Miyazawa*

Miyazawa Clinic, Hyogo, Japan

EDITORIAL

Long-chain unsaturated fatty acids and medium-chain saturated fatty acids are effective in deactivating enveloped viruses [1]. Short-chain (butyric, caproic, and caprylic) and long-chain saturated (palmitic and stearic) fatty acids had no or a very small antiviral effect [1]. Monocaprin, the 1-monoglyceride of capric acid, and lauric acid caused a >1,00,000-fold reduction in the viral titer of herpes simplex virus in 1 min at a concentration of 20 mM [2]. Unsaturated free fatty acids such as oleic acid, arachidonic acid and linoleic acid inactivate enveloped viruses such as herpes, influenza within minutes of contact at micromolar concentrations [3,4]. Electron micrographs of enveloped treated viruses indicate that the inactivation is associated with disintegration of the virus envelope [3,4]. As SARS-CoV-2 is an enveloped virus, it may also be vulnerable to deactivation by these fatty acids. These fatty acids were effective especially in their monoglyceride form [1], which acts as emulsifiers. Emulsifiers belong to the chemical class of surfactants. The Japanese Ministry of Economy, Trade and Industry recently confirmed that SARS-Cov-2 was deactivated by several diluted surfactants [5]. Emulsifiers and fatty acids are widely used as food additives and are harmless, even if swallowed. Persons affected with SARS-CoV-2 may be able to reduce the infectivity of their saliva or saliva-derived droplets by inclusion of those in their mouth. Saliva is used for PCR based tests to detect COVID-19. However, it can be difficult to collect samples from elderly people with low saliva production [6]. Flavored foods and liquids can stimulate salivation. Therefore, if it is confirmed that emulsifiers and/or fatty acids deactivate SARS-CoV-2 in the saliva and do not adversely affect RNA stability; then there may be a benefit in their inclusion in the mouth at the time of specimen collection, both to stimulate salivation and to reduce the risk to the individual collecting samples. Moreover, the presence of emulsifiers and/or fatty acids in the mouth before or during speech, or dental treatment may reduce the infectivity of the saliva and saliva-derived droplets. If we assume that SARS-CoV-2 infects and replicates in the oral mucosa because angiotensin converting enzyme-2, the host receptor for target cell entry, is highly expressed on the epithelial cells of oral mucosa [7], then repeated introduction of emulsifiers and/or fatty acids in the

mouth may prevent aggravation of the disease due to entry of the virus into the respiratory tract. *In vitro* and *in vivo* research to test this hypothesis is desired.

CONTRIBUTOR AND GUARANTOR INFORMATION

DM contributed to the conception/design of the work; the acquisition, analysis, and interpretation of data; the drafting and revision of the manuscript. DM approved the current version of the manuscript and agree to be accountable for all aspects of the work.

DISCLOSURE

The authors declare no conflicts of interest.

REFERENCES

1. Thormar H, Isaacs CE, Brown HR, Barshatzky MR, Pessolano T. Inactivation of enveloped viruses and killing of cells by fatty acids and monoglycerides. *Antimicrob Agents Chemother*. 1987;31(1):27-31.
2. Kristmundsdóttir T, Arnadóttir SG, Bergsson G, Thormar H. Development and evaluation of microbicidal hydrogels containing monoglyceride as the active ingredient [published correction appears in *J Pharm Sci* 1999 Dec;88(12):1366]. *J Pharm Sci*. 1999;88(10):1011-1015.
3. Kohn A, Gitelman J, Inbar M. Unsaturated free fatty acids inactivate animal enveloped viruses. *Arch Virol*. 1980;66(4):301-307.
4. Kohn A, Gitelman J, Inbar M. Interaction of polyunsaturated fatty acids with animal cells and enveloped viruses. *Antimicrob Agents Chemother*. 1980;18(6):962-968.
5. Ministry of Economy, Trade and Industry, Surfactants and Hypochlorous Acid Solution for Removal of Coronavirus from Surfaces (Final Announcement), 26 June 2020.
6. Xu F, Laguna L, Sarkar A. Aging-related changes in quantity and quality of saliva: Where do we stand in our understanding?. *J Texture Stud*. 2019;50(1):27-35.
7. Xu H, Zhong L, Deng J, Peng J, Dan H, Zeng X, et al. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci*. 2020;12(1):8.

Correspondence to: Daisuke Miyazawa, MD, Miyazawa Clinic, 1-6-5 Akuraminamin Takarazuka, Hyogo 665-0823, Japan, E-mail: kusami1@ybb.ne.jp

Received: September 16, 2020, **Accepted:** November 02, 2020, **Published:** November 09, 2020

Citation: Miyazawa D (2020) SARS-CoV-2 in Saliva May Be Deactivated with Fatty Acids or Emulsifiers. *J Nutr Food Sci*.10:782. doi: 10.35248/2155-9600.20.10.782

Copyright: © 2020 Miyazawa D. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.