

## Segregation of Malignant Hematological Disease in Families with Malignant Lymphoma

Haneef Awan<sup>1</sup>, Johanna Furuholmen<sup>2</sup>, Geir E. Tjønnfjord<sup>3</sup>, Bernt Ly<sup>4</sup>, Tom Børge Johannesen<sup>4</sup>, Anne Tierens<sup>5</sup>, Jan Delabie<sup>5</sup> and Viggo Jonsson<sup>6\*</sup>

<sup>1</sup>IT Section, Institute of Clinical Medicine, University of Oslo, Blindern NO 0318 Oslo, Norway

<sup>2</sup>Medical Faculty, University of Oslo, Blindern NO 0318 Oslo, Norway

<sup>3</sup>Department of Hematology, Institute of Clinical Medicine, University of Oslo, Nydalen NO 0424 Oslo, Norway

<sup>4</sup>Norwegian Cancer Registry, Majorstuen NO 0304 Oslo, Norway

<sup>5</sup>Department of Pathology, University of Oslo, Nydalen NO 0424 Oslo, Norway

<sup>6</sup>Department of Hematology, Institute of Clinical Medicine, University of Oslo, Nydalen NO 0424 Oslo, Norway

### Abstract

Familial malignant lymphoma, viz. more than two cases of malignant hematological disease in a family of which one diagnosis is malignant lymphoma, was seen in 43 (37 per cent) of the families in our database of familial malignant hematological disease. Genealogical examination of the 43 pedigrees after multiple ascertainment showed an equal amount of vertical transmissions (affected parent-offspring and grandparent-parent-offspring combinations) and non-vertical transmissions (affected uncle, aunt-nephew, niece, cousin combinations) without evident Mendelian pattern and no significant difference between observed and expected patrilineal and matrilineal lines in spite of a marked predominance of males. A marked pleiotropic diversity of involved diagnoses comprised 57 (93 per cent) lymphoproliferative- and 4 (7%) myeloproliferative diseases. Both Hodgkin's lymphoma, non-Hodgkin's lymphoma apart from the diffuse large B-cell lymphoma and chronic lymphocytic leukemia had a strong mutual association, and a weaker yet significant association to multiple myeloma and to diffuse large B-cell lymphoma. Compared with the number of patients in the population extracted from the crude age-adjusted incidences, the observed number of patients in familial disease was significantly higher interpreted as a stronger expression of congenic susceptibility among family members with reservations related to different environmental factors. Signs of anticipation in all combinations but no birth order effect were observed. It is discussed that an epigenetic parental genomic imprinting as a modifier for the segregation of linked susceptibility loci theoretically would bring about a similar pattern with male predominance and pleiotropic diversity of diagnoses away from any Mendelian expectation.

**Keywords:** Malignant Lymphoma; Leukemia; Genetics; Epidemiology

### Introduction

Malignant Lymphomas (ML) comprising Hodgkin's Lymphoma (HL) and Non-Hodgkin's Lymphoma (NHL) are characterized by malignant transformation of lymphoid cells where a monoclonal expansion of the tumor cell emerge from a mutation in a lymphoid progenitor cell at the lymphoid differential pathway [1,2]. About 40 different ML diagnoses are defined, each with specific histological, immunophenotypic and clinical features [3]. In principle, the etiology of ML consists of both a genetic inborn susceptibility and a possible environmental influence, e.g. from antigenic drive of the tumor growth from lymphotropic microorganisms, and from a possible mutagenic effect of pesticides, solvents and dyes [4-6].

The genetic etiology is supposed to be related to congenic risk alleles [7,8], so far mainly investigated in chronic lymphocytic leukemia (CLL) [9-16]. Presumably, each diagnosis within the entity malignant lymphoproliferative disorders (LPD) has its own risk alleles linked to each other, which eventually explains the observed association between LPD diagnoses with familial, pleiotropic clustering [12,13]. Such congenic risk alleles, also denoted susceptibility loci, or genetic inborn susceptibility for the specific mutation which causes the monoclonal of the diagnosis, have been demonstrated by means of genome-wide association studies. In CLL, for example, up to ten or perhaps a few more susceptibility loci have been demonstrated during the last years at many different positions in the genome, and with a range of putative associations to CLL [15,16]. The same pattern is seen in HL where four or perhaps five susceptibility loci known so far are scattered throughout the genome with a range of putative associations [17,18].

The familial clustering of ML and other types of LPD has been proved in recent computerized estimations of large-scaled data from cancer registries while only few genealogical investigations based on the study of pedigrees from affected families have been published [19-21]. The computerized estimations of data from cancer registries show remarkable inconstancy in frequencies and incidences of clustered LPD diagnoses versus solitary disease, and marked differences in demographic data of the patients [22-25]. Genetic anticipation, which is increased severity in the form of lower age at onset of disease and more and more high-malignant cases down through the generations, and birth order effect with rank order by age of the affected sib in the sib ship, are also reported with marked differences [26-29]. The mode of segregation is so far largely unknown.

The purpose of the present paper is to compare these findings from mainly large-scaled computerized data with data from genealogical assessments of pedigrees from familial ML.

**\*Corresponding author:** Viggo Jonsson, Department of Hematology, Riks Hospital, Institute of Clinical Medicine, University of Oslo, Nydalen, NO 0424 Oslo, Norway, Tel: +47 22 89 46 14; E-mail: [viggo.jonsson@medisin.uio.no](mailto:viggo.jonsson@medisin.uio.no)

**Received** August 25, 2014; **Accepted** October 08, 2014; **Published** October 14, 2014

**Citation:** Awan H, Furuholmen J, Tjønnfjord GE, Ly B, Johannesen TB, et al. (2014) Segregation of Malignant Hematological Disease in Families with Malignant Lymphoma. J Genet Syndr Gene Ther 5: 249. doi:10.4172/2157-7412.1000249

**Copyright:** © 2014 Awan H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Material and Methods

### Families

Our database on familial malignant hematological disease, viz. families with a least two affected family members (116 families), has 43 families in which ML is one of the diagnoses. The database has been collected systematically over the past decade in our clinical work with all types of hematological malignancies in Oslo and Copenhagen. Parts of this material has previously been published [20,21], but never as a comprehensive description of familial ML and never in the form like that presented here.

### Inclusion

Each patient underwent a face-to-face interview about other family members with malignant blood disease and the pedigree was drawn up. To ensure maximal ascertainment, the interview was individually adjusted and modulated to include also old terminology if necessary. HL for example was earlier lympho granulomatosis, and CLL was sometimes “old man’s disease”. Each patient was also asked about the number and position of healthy family members in the family tree, stillborn and extramarital persons. The patients were allowed time for discussion at home and with relatives. All information was crosschecked with the Cancer Registry in Norway and Denmark. Old hospital records with histopathological and laboratory reports were included in the crosscheck if necessary. Each family was included with signed consent of the proband and after information about the purpose

of the study, that data are confidential and unrecognizable outside the study and that the study was approved by the Scientific-Ethical Committees and the Data Protection Agency in Norway and Denmark.

The affiliation of each ML proband in all 43 pedigrees was grouped into the following categories: (A) parent-offspring pairs and (B) grandparent-parent-offspring combinations (A and B represent vertical inheritance), (C) uncle, aunt-nephew, niece cousin combination and (D) sib concordance which denote two or more affected siblings without affected relatives in other generations (C and D represent non-vertical inheritance). The number of patrilineal and matrilineal transmissions was recorded in (A) and (B), in (C) it is mixed and in (D) unknown.

The mean number of healthy family members crosschecked per family was about 50 persons. The Civil Person Registry was used for crosscheck in some few cases with uncertainty about the number or position of healthy relatives in the family tree. About 30 % of the healthy family members were still alive at the end of study. About 150 generations were evaluated and crosschecked, 86 generations (Table 1) with one or more cases of malignant hematological disease.

### Diagnoses

All diagnoses were based on standard criteria according to WHO classification and ICD-10 nomenclature [3]. In case the crosscheck of older family members came up with diagnoses according to former classification systems, mainly Lukes & Collins, Kiel, and Working

DIAGNOSIS	FAMILIAL RELATIONSHIP Number (males, females)			FAMILIES vs. NORMAL			
	Table 2	Table 3	Total	Rate of ML (%)		Ratio males, females	
				families	normal	families	normal
HL	6 (5,1)	5 (3,2)	11 (8,3)	21	10	2.7	1.5
FL	3 (3,0)	6 (3,3)	9 (6,3)	17	20-25	2.0	1.1
DLBCL	8 (5,3)	4 (2,2)	12 (7,5)	23	30-35	1.4	1.2
LPL	3 (2,1)	3 (1,2)	6 (3,3)				
BL	2 (0,2)	2 (2,0)	4 (2,2)				
T	1 (1,0)	1 (1,0)	2 (2,0)				
MCL		1 (1,0)	1 (1,0)				
MZL	1 (1,0)	2 (2,0)	3 (3,0)				
NOS	2 (0,2)	2 (2,0)	4 (2,2)				
NHL	20 (12,8)	21 (14,7)	41 (26,15)	38	35	1.7	1.3
HL & NHL	26 (17,9)	26 (17,9)	52 (34,18)	100	100	1.9	1.2
CLL	22 (14,8)	15 (10,5)	37 (24,13)			1.9	1.9
LGTCCL		2 (1,1)	2 (1,1)				
T PLL		1 (0,1)	1 (0,1)				
MM	3 (3,0)	2 (1,1)	5 (4,1)			4.0	1.4
AML		1 (0,1)	1 (0,1)				
CML		1 (0,1)	1 (0,1)				
MF		1 (0,1)	1 (0,1)				
ET		1 (1,0)	1 (1,0)				
<b>Patients</b>	<b>51 (34,17)</b>	<b>50 (30,20)</b>	<b>101 (64,37)</b>				
<b>Families</b>		<b>22</b>		<b>21</b>		<b>43</b>	
<b>Generations</b>		<b>45</b>		<b>41</b>		<b>86</b>	
<b>Patrilineal</b>		<b>13</b>		<b>3</b>		<b>16</b>	
<b>Matrilineal</b>		<b>7</b>		<b>9</b>		<b>16</b>	
<b>Patri- and matrilineal</b>		<b>2</b>				<b>2</b>	

**Abbreviations:** HL Hodgkin's lymphoma; FL follicular lymphoma grade I-III; DLBCL diffuse large B-cell lymphoma; LPL lymphoplasmocytic lymphoma; BL Burkitt's lymphoma; T Precursor T-lymphoblastic lymphoma; MCL mantle cell lymphoma; MZL marginal zone B-cell lymphoma; NOS not otherwise specified; NHL non-Hodgkin's lymphoma, CLL chronic lymphocytic leukemia; LGTCCL large granular T-cell leukemia; T PLL T-cell prolymphocytic leukemia; MM multiple myeloma; AML acute myeloblastic leukemia; CML chronic myeloid leukemia, MF myelofibrosis; ET essential thrombocytopenia

**Table 1:** Malignant hematological disorders in 43 families with malignant lymphoma.

formulation, the diagnosis was converted to the WHO system and ICD-10 nomenclature to make the material homogenous and feasible for comparison (Table 1). This conversion was generally easy with some few exceptions denoted NOS (not otherwise specified) in the tables.

### Statistics

The diagnoses in familial ML were compared with the occurrence in the population by means of simple regression of the scores observed (Tables 1-3) and crude age-adjusted incidences, viz. number of patients per 100.000 persons in the population with the age of the diagnosis in question. The incidences used are an estimated mean for Norway and Denmark from the observation period 1950-2010, based on official Cancer Registry reports [30-32]. An estimated mean has been used to adjust for an increasing incidence of ML in general and a specific rise in DLBCL during the past decades [30-32]. Age at onset of disease in parent-offspring pairs, and male-female ratios were estimated by means of Wilcoxon test, pair difference. Two-sided P-value < 0.05 was considered significant. Haldane Smith test was used for estimation of birth order effect [33]. In this test the sum (6A) of the birth orders, viz. the rank in the sib ship by age of affected sibs, is compared with the theoretical value, expressed as the 95 % confidence interval (CI 95%) [33].

### Segregation analysis

X<sup>2</sup>- test statistics with one degree of freedom was used to test for difference between observed and expected number of patri- and matrilineal transmissions in vertical inheritance (category A and B). The expected numbers were calculated on the assumption that the material is random distributed with frequencies for patrilineal (father-son: np<sup>2</sup>; father-daughter: npq) and matrilineal transmission (mother-son: nqp; mother daughter: nq<sup>2</sup>) where p and q denote the proportions of males and females, respectively, and n the total number of families included [34].

### Pattern recognition

Trends towards a likely Mendelian pattern in the distribution of affected family members in the families was done from a visual inspection of each pedigree compared with the standard pattern in autosomal dominant-, autosomal recessive- and Y and X-linked hybridization [35].

### Results

#### Diagnoses

ML was found in 43 (37 per cent) of 116 families with two or more

ONE AFFECTED OFFSPRING							
Family number	Diagnosis and age at onset of disease (years)					Birth order	
	Father		Son		Daughter		Rank of affected sib/ total number of sibs
74	HL	63	CLL	60			1/1
86	HL	49			CLL	58	2/3
30	FL	55	CLL	43			2/3
25	DLBCL	67	HL	36			2/3
11	DLBCL	74	CLL	46			2/3
46	DLBCL	54	CLL	43			3/3
68	LPL	65	MM	42			2/4
5-29	MZL	66	CLL	42			2/2
6	CLL	66			DLBCL	64	3/3
Age at onset, father-offspring: P= 0.02 (Wilcoxon test, pair difference, two- sided)							Birth order effect 6A (95%CI): 114 (74-130) ,P > 0.05 (Haldane Smith test)
	Mother		Son		Daughter		
69	DLBCL	70			DLBCL	49	1/1
27	BL	68	CLL	47			1/3
67	LPL	75	HL	26			2/2
75	NOS	70	CLL	61			1/2
107	CLL	82	FL	46			2/2
16	CLL	78	LPL	76			2/3
Age at onset, mother-offspring P = 0.032							54 (40-74) P>0.05
TWO AFFECTED OFFSPRING							
	Father		Son		Daughter		
114	HL	55			CLL ,CLL	60,60	1/2, 2/2
38	FL	75	CLL, MM	61,60			2/3, 3/3
1	CLL	75	T, CLL	57, 70			2/5, 4/5
80	MM	91	DLBCL,CLL	35, 52			1/2, 2/2
	Mother						
97	NOS	88			CLL, CLL	69, 67	3/4, 4/4
Age at onset, parent-offspring P = 0.014							144 (99-153) P> 0.05
BOTH PARENT'S LINES AFFECTED							
82	Father DLBCL 80 yr - Daughter CLL 61 yr Mother's sister HL						
115	Mother BL 76 yr – Son CLL 63 yr Father's father CLL						

**Abbreviations:** 6A of the Haldane Smith test for birth order effect is the sum of birth orders from affected persons, compared with the theoretical value, expressed as the 95 per cent confidence interval. Other abbreviations, see footnote to Table 1.

**Table 2:** Familial malignant lymphoma. Parent-offspring combinations.

GRAND PARENT – PARENT – OFFSPRING			
Family number	Combination	Diagnosis and age at onset (years)	Birth order
10	Grand father-father-son	CLL 72 – CLL 58 – HL 32	4/7, 2/3, 2/2
10	Father's sister	CLL 61	3/3
35	Grand mother – mother - son	CLL 71 – FL 61 – HL 36	US, 1/1, 1/2
95	Grand mother – mother - son	CLL 86 – normal – BL 21	US, 2/4, 1/4
96	Grand mother – mother - son	T PLL 42 – normal – NOS 36	1/6, 1/4, 2/2
37	Grand mother – mother - daughter	MM 79 – LPL 56 – FL 42	US, 1/3, 1/2
Age at onset, grandparent – parent and parent - offspring P = 0.004			Birth order effect 6A (95%CI): 114 (100-176) ,P > 0.05 (Haldane Smith test)
UNCLE, AUNT – NEPHEW, NIECE, COUSIN			
Family number	Combination	Diagnosis and sex (m male, f female)	Evaluable sibships (n) and Birth order
77	Proband – brother's daughter	DLBCLm – CLLf	(2) 3/3, 3/3
89	Proband – brother's son	HLm – CLLm	(2) 2/3, 2/4
89	Proband's brother	MMm	3/3
76	Proband – mothers's sister's son	FLf – CLLm	(2) 1/1, 1/3
78H	Proband – son	DLBCLf – CLLm	(2) 1/2, 1/2
78H	Proband's sister	AMLf	2/2
102	Proband –mother's sister's son	DLBCLf – CLLm	(2) 1/2, 5/7
102	Proband's brother	FLm	2/2
104	Proband – mother's sister	MCLm – CMLf	(1) 2/2, US
104	Proband's brother	CLLm	1/2
108	Proband –mother's brother's son	Tm – CLLm	(2) 4/5, 3/4
			222 (158-244) P> 0.05
SIB CONCORDANCE			
Combination and birth order			
Family number	Brother – Brother	Family number	Sister – Brother
78B	LPL – LGTCL 1/4 , 4/4	79B	LGTCL – FL 2/3, 3/3
85	BL – CLL 1/3, 2/3		
82B	MZL – FL 2/3, 3/3		Sister – Sister
81B	ET – DLBCL 1/2 , 2/2	61	HL - HL 1/3, 3/3
	Brother – Sister		
83 B	MZL –MF 1/3, 2/3		
58	NOS – CLL 1/2 , 2/2		
79A	CLL – LPL 1/4, 4/4		

Abbreviations: US uncertain. Other abbreviation, see footnote to Table 1 and 2.

Table 3: Familial malignant lymphoma. Other combinations.

cases of malignant hematological disease. The diversity of malignant hematological disorder associated with familial ML includes 17 different diagnoses, 13 Lympho proliferative Disorders (LPD), and 4 Myelo proliferative Disorders (MPD) in 101 patients, 97 (96 per cent) with LPD and 4 (4 per cent) with MPD. Per family, we found 2.3 cases (mean) of LPD, 0.1 cases (mean) of MPD and 1.2 cases (mean) of ML (Table 1).

The number of affected family members was significantly higher than expected (P < 0.001) estimated from the crude, age-adjusted Scandinavian incidences, viz. the number of patients per 100.000 persons in the population. About 2.000 persons were screened within the 43 families, 50 persons as a mean per family, giving: HL incidence 2.5/100.000, expected per 2.000 persons 0.05, observed 11; NHL incidence 8.0/100.000, expected 0.16, observed 41; CLL incidence

5.5/100.000, expected 0.1, observed 37; multiple myeloma (MM) incidence 5.0/100.1000, expected 0.1, observed 5 (Table 1). Based on these figures, the 43 ML probands are associated especially with CLL (37 patients), and ML (52-43 = 9 patients). In spite of nearly the same incidences of CLL and MM in the Scandinavian population (5.5 versus 5.0/100.000), only 5 cases of MM were seen in the families compared with 37 CLL patients (Table 1). No cases of ALL were noted (Table 1).

The rate of lymphoma subtypes in the population compared with that of the 43 families shows predominance of HL in familial ML (expected 10%, observed 21%, P< 0.05) , and a significant reduction of DLBCL in familial ML (expected 30-35%, observed 23 %, P<0.01) (Table 1).

A total of four cases of MPD in 43 families with about 2.000 persons

screened is significantly more than expected ( $P < 0.01$ ) and indicate an association between LPD and MPD.

The male/female ratio was generally higher in familial ML (expected 1.3, observed 1.7,  $P < 0.005$ ) and highest in HL (expected 1.5, observed 2.7), sample size 8 males and 3 females (Table 1), counts too small for convincing statistical description.

## Segregation

Vertical transmission was seen in 22 families (Tables 1 and 2), non-vertical transmission in 21 families (Tables 1, 3). Ten LPD diagnoses were seen in the vertical transmission while 13 LPD diagnoses were seen in non-vertical transmissions (Tables 1-3). All 4 cases of malignant myeloproliferative disorders were found in non-vertical transmissions (Tables 1 and 3). Taken together, the distribution of affected family members in all 43 pedigrees do not fit into one, common Mendelian modality of segregation. Neither was such pattern recognition successful with the diagnoses ML, NHL, HL and CLL separately.

A birth order effect was not observed in familial ML (Tables 1-3). Neither did we see a birth order effect from an estimation of patrilineal (6A 282, 95% CI: 211 – 299,  $n = 16$ ,  $P > 0.05$ ) vs. matrilineal (6A 162, 95% CI: 158 – 238,  $n = 16$ ,  $P > 0.05$ ) transmissions.

The age at onset of disease was significantly higher ( $P < 0.01$ ) in parents than in offspring apart from two families with parental HL (nos. 86 and 114, Table 2).

The number of patrilineal and matrilineal transmissions was the same when all types of transmissions were evaluated (Table 1).

Patrilineal transmission was predominant in the direct vertical transmissions, 13 transmissions from father's side and 7 from mother's side (Table 1) due to surplus of affected males in this material. The observed vs. expected numbers of patrilineal and matrilineal transmissions were not different ( $\chi^2$ -test with one degree of freedom), and no mode of transmission such as father-son (F-S), father-daughter (F-D), mother-son (M-S) or mother-daughter (M-D) was superior after correlation for the gender of the included patients in the asserted vertical transmission (Tables 1 and 2):

ML correlated.  $n = 22$ ,  $p = 0.65$ ,  $q = 0.35$ , F-S expected  $np^2 = 9$ , observed 10, F-D expected  $npq = 5$ , observed 5, M-S expected  $nq^2 = 5$ , observed 5, M-D expected  $nq^2 = 3$ , observed 1 ( $P$  ranging between 0.5 and 0.25).

LPD correlated.  $n = 22$ ,  $p = 0.67$ ,  $q = 0.33$ . F-S expected 10, observed 10, F-D expected 5, observed 5, M-S expected 5, observed 6, M-D expected 2, observed 1 ( $P$  ranging between 0.5 and 0.25).

CLL correlated.  $n = 18$ ,  $p = 0.64$ ,  $q = 0.36$ . F-S expected 7, observed 9, F-D expected 4, observed 4, M-S expected 4, observed 5, M-D expected 2, observed 0 ( $P$  ranging between 0.25 and 0.1).

Combined patrilineal- and matrilineal transmission, viz. affected family members in both father's and mother's lines, was seen in only 2 families, both with affected parent-offspring (Tables 1 and 2).

Sex concordance in sibling pairs (male-male  $np^2$ ), (female-female  $nq^2$ ) compared with sex discordance (male-female and female-male  $2npq$ ) showed no significant findings: Male-male plus female-female expected 9, observed 13. Male-female, female-male expected 8, observed 5 when  $n = 18$ ,  $p = 0.64$ ,  $q = 0.36$  ( $P$  ranging between 0.25 and 0.1).

## Discussion

Bias in our material is related to the low-grade disorders which sometimes are overlooked due to no or only minor symptoms.

Typical examples are MGUS (Monoclonal Gammopathy of Uncertain Significance), stage A CLL and polycythemia vera. The impact of this systematic bias is presumably equal in the multiple familial cases, in solitary disease, and in the incidences used for a comparison between occurrence of disease in families and in the population. One main problem with the underestimated low-grade cases is their absence in the family trees when to estimate the number and position of healthy persons vs. patients. We do not have a systematic screening of all family members in affected families for a precise estimation of this bias. Such a screening will imply the recruitment of healthy, uninformed people without formal social contact with the proband, e.g. family members at a distance from the proband in the pedigree and therefore, it will not be approved by our national ethical committees. Furthermore, the unknown number of early abortions, experienced as a late, strong menstruation, which should rightly be included in the family size when to estimate parity, invalidate our data while legal- and late abortions and still born persons are already included. About 30 per cent of the healthy family members are alive at the time of investigation and still at risk for later malignant hematological disease which could change the findings. We saw a significant higher age at onset of disease in parents than in offspring (Tables 1-3), interpreted as a sign of anticipation with the reservation that the age of the mother at the time of delivery has a trend towards a higher age during the course of the observation period, and that age at onset as a parameter in anticipation theoretically should be related to the mean age of onset of the disease in question. In CLL for example, we found the highest age at onset in the parents [20] with some few exceptions related to e.g. parental HL with a generally lower age at onset than in CLL (cf. family nos. 86 and 114 Table 2). Finally, environmental factor such as infections with antigenic drive and toxic reagents with mutagenic effect, which evidently can affect the expression of ML [6] may have changed in the observation period.

With these reservations, 13 LPD diagnoses and four MPD diagnoses were found within the pleiotropic diversity to familial ML (Tables 1-3). A strong linking between NHL and HL and between ML and CLL was seen in contrast to a weaker association between ML and MM. The higher rate of affected males in familial than in solitary cases (Tables 1-3) may well be explained by a stronger expression of genetic LPD-susceptibility and hence a higher rate of affected males in families with multiple cases than in families with solitary cases. These genealogical data convey a description of the so-called first generation familial disorders known from large-scale estimations of malignant hematological disease [22-25] and prove that no simple Mendelian segregation is on question.

The predominance of HL and the reduction in DLBCL in familial ML (Tables 1-3) can be interpreted as an unequal affinity in the pleiotropic clustering ("HL goes together with HL, but DLBCL has its own ways"). Such positive and negative linking between LPD-diagnoses would be the expected outcome of segregation of mosaics of susceptibility alleles with unequal putative associations. In some segregation, one would predict an additive enhancement of the susceptibility alleles in the mosaic; in other segregations no such synergism would be expected. Finally, an antagonistic, suppressive effect between the susceptibility alleles in the mosaic is a theoretical outcome which could explain the occurrence of single cases (without signs of familial aggregation) of disease in the population.

In such a genetic system with linked susceptibility, male predominance, and segregation away from any Mendelian expectation, the presence of a modifier of segregation should be considered [36]. In principle, such a modifier causing transgenerational segregation

distortions is a genetic mechanism which violate Mendel's first law about equal representation of the two alleles at a heterozygous locus in the functional products of meiosis (egg or sperm) [36]. Models with modifiers linked and unlinked to the primary heterozygote locus have been described for instance in parental genomic imprinting [37-43], all with different segregation in males and females explained by an unequal effect of the modifier in spermiogenesis and oogenesis [36,44].

It is intriguing that the malignant hematological disorders fulfill a number of criteria for the presence of a modifier, among other things the lack of a clear Mendelian pattern in the segregation down through the generations of affected families in combination with male predominance, seen in large subsets of disease, for example in ML and CLL. Genomic imprinting with maternal down regulation, viz. "imprinting of paternal susceptibility" has previously been attributed to CLL-susceptibility as a possible mechanism in the transmission of CLL via fetomaternal microchimerism [20], but very few genealogical publications on the segregation of malignant hematological disease are available for comparison [for review see 19-21]. Sex-concordance, viz. surplus of male-male and female-female pairs vs. male-female pairs in siblings with CLL [45], and the lack of "any genetic effect" in lymphoreticular malignancies associated familial, endemic HL [19] may be interpreted as signs of distortion.

We found an equal number of expected versus observed patrilineal and matrilineal vertical transmissions to sons and daughters when data has been  $X^2$ -adjusted for the male predominance. In this way the predominance of father-son transmissions is explained as a consequence of the surplus of males in the material among an expected proportion of father-daughter and mother-son, mother-daughter transmissions. In other words, the father-son predominance is a distortion away from a Mendelian expectation which is not caused by impaired transmissions to daughters. However, a qualitative difference between matrilineal and patrilineal transmission has been reported in CLL, where a birth order effect to affected offspring could be seen only in patrilineal lines [20]. No such birth order effect was seen in the present material on ML (Tables 2-3). A number of parental transgenerational distortions in epigenetic genomic imprinting are based on reciprocal heterozygotes, viz. persons with the same genotype but with parental origin of their two alleles reversed [36]. In such heterozygotes, pre-conceptual parental environment experiences can sometimes be incorporated into the germ line through changes in the male gametes due to nearly life-long production in the spermiogenesis with a standing capacity to catch up phenotypic, epigenetic impulses, in contrast to the oocytogenesis with its limited pool of cells at birth and no further production [44,46,47]. Since no relationship between susceptibility to malignant hematological disease and the Y or X chromosome has been described, for all we know, the effect of such heterozygotes in parental genomic imprinting may perhaps represent a plausible, alternative explanation.

Recommendation: There can be no doubt about an inherited genetic component in the etiology of LPD and most likely, the susceptibility alleles represent the genotype of this genetic component. Further analyze of susceptibility loci to each LPD diagnosis is therefore crucial when to explore the pattern of segregation. Together with genealogical data from pedigrees from affected families, we know today that the segregation is non-Mendelian and, most likely, influenced by parental genomic imprinting, genetic anticipation, birth order effect and a modifier effect directed towards the susceptibility alleles. We are only at the beginning and we need more family studies to complete the picture. Besides the theoretical understanding of a genetic entity,

knowledge on the segregation of LPD will undoubtedly provide a splendid practical tool for the diagnostic and the prognostic estimation of each patient, and when to look for family donors and disregard those who have a load of susceptibility without manifest disease.

#### Acknowledgement

We are grateful to the patients and their families for their participation and to our colleagues who referred patients for the study. Professor David Haig, Department of Organismic and Evolutionary Biology, Harvard University, US, is thanked for valuable comments to the manuscript. The study was undertaken in accordance with the Declaration of Helsinki and approved by the Committee for Medical Research Ethics in Norway (record no. REK S-06353b), updated and reconfirmed, December 10, 2013 (record no. REK 2013/2002/REK south-south east B), the Data Inspectorate in Norway (record no. 07/00254-2) and the Social and Health Directorate of Norway (record no. 07/324). The Royal Danish Archives and the Provincial Archives of Sealand (record no. 2000-441-0023), the Danish Data Protection Agency (record no.2000-41-0184), the Danish Board of Health (record no.123-63-2000), and the Danish Scientific-Ethical Committees (record no 01-224/01). The Norwegian part of the study was done under the patronage of the Norwegian Hematological Society.

#### References

1. Kadin ME (2001) Hodgkin's disease, cell of origin, immunobiology, and pathogenesis. In: Neoplastic Hematopathology, Knowles DM (ed.) Philadelphia: Lippincott Williams & Wilkins 667-690.
2. Bain BJ (2001) Overview, Cytogenetics in Hematology. In: Best Practice & Research Clinical Haematology, Buckle VJ, Kearney L (ed.) London: Baillière Tindall Pub., 14: 463-477.
3. Jaffe E, Harris NL, Stein H et al. (2001) WHO classification of tumours, pathology and genetics of tumours of the haematopoietic and lymphoid tissues. Lyon: IARC Press.
4. Hjalgrim H, Smedby KE, Rostgaard K, Molin D, Hamilton-Dutoit S, et al. (2007) Infectious mononucleosis, childhood social environment, and risk of Hodgkin lymphoma. *Cancer Res* 67: 2382-2388.
5. Skibola CF, Curry JD, Nieters A (2007) Genetic susceptibility to lymphoma. *Haematologica* 92: 960-969.
6. Morton LM, Wang SS, Cozen W, Linet MS, Chatterjee N, et al. (2008) Etiologic heterogeneity among non-Hodgkin lymphoma subtypes. *Blood* 112: 5150-5160.
7. Auer RL, Dighiero G, Goldin LR, Syndercombe-Court D, Jones C, et al. (2007) Trinucleotide repeat dynamic mutation identifying susceptibility in familial and sporadic chronic lymphocytic leukaemia. *Br J Haematol* 136: 73-79.
8. Hastings PJ, Lupski JR, Rosenberg SM, Ira G (2009) Mechanisms of change in gene copy number. *Nat Rev Genet* 10: 551-564.
9. Houlston RS, Sellick G, Yuille M et al. (2003) Causation of chronic lymphocytic leukaemia - insight from familial disease. *Leuk Res* 27: 871-876.
10. Marti GE, Carter P, Abbasi F et al. (2003) B-cell monoclonal lymphocytosis and B-cell abnormalities in the setting of familial B-cell CLL. *Cytometry, Part B, Clinical Cytometry* 52: 1-12.
11. Sellick GS, Webb EL, Allinson R et al. (2005) A high-density SNP genome-wide linkage scan for chronic lymphocytic leukemia-susceptibility loci. *Am J Hum Genet* 77: 420-429.
12. Di Bernardo MC, Crowther-Swanepoel D, Broderick P, Webb E, Sellick G, et al. (2008) A genome-wide association study identifies six susceptibility loci for chronic lymphocytic leukemia. *Nat Genet* 40: 1204-1210.
13. Sherborne AL, Houlston RS (2010) What are genome-wide association studies telling us about B-cell tumor development? *Oncotarget* 1: 367-372.
14. Crowther-Swanepoel D1, Houlston RS (2010) Genetic variation and risk of chronic lymphocytic leukaemia. *Semin Cancer Biol* 20: 363-369.
15. Slager SL, Skibola CF, Di Bernardo MC, Conde L, Broderick P, et al. (2012) Common variation at 6p21.31 (BAK1) influences the risk of chronic lymphocytic leukemia. *Blood* 120: 843-846.
16. Speedy HE, Di Bernardo MC, Sava GP, Dyer MJ, Holroyd A, et al. (2014) A genome-wide association study identifies multiple susceptibility loci for chronic lymphocytic leukemia. *Nat Genet* 46: 56-60.

17. Enciso-Mora V, Broderick P, Ma Y, Jarrett RF, Hjalgrim H, et al. (2010) A genome-wide association study of Hodgkin's lymphoma identifies new susceptibility loci at 2p16.1 (REL), 8q24.21 and 10p14 (GATA3). *Nat Genet* 42: 1126-1130.
18. Cozen W, Timofeeva MN, Li D, Diepstra A, Hazelett D, et al. (2014) A meta-analysis of Hodgkin lymphoma reveals 19p13.3 TCF3 as a novel susceptibility locus. *Nat Commun* 5: 3856.
19. Thompson EA (1981) Pedigree analysis of Hodgkin's disease in a Newfoundland genealogy. *Ann Hum Genet* 45: 279-292.
20. Jønsson V, Tjønnfjord GE, Johannesen TB, Ly B, Olsen JH, et al. (2010) Familial chronic lymphocytic leukemia in Norway and Denmark. Comments on pleiotropy and birth order. *In Vivo* 24: 85-95.
21. Jønsson V, Awan H, Nyquist E, Maisenhølder M, Johannesen TB, et al. (2011) Familial Hodgkin's lymphoma in Scandinavia. *In Vivo* 25: 431-437.
22. Goldin LR, Pfeiffer RM, Li X, Hemminki K (2004) Familial risk of lymphoproliferative tumors in families of patients with chronic lymphocytic leukemia: results from the Swedish Family-Cancer Database. *Blood* 104: 1850-1854.
23. Goldin LR, Pfeiffer RM, Gridley G, Gail MH, Li X, et al. (2004) Familial aggregation of Hodgkin lymphoma and related tumors. *Cancer* 100: 1902-1908.
24. Segel GB, Lichtman MA (2004) Familial (inherited) leukemia, lymphoma, and myeloma: an overview. *Blood Cells Mol Dis* 32: 246-261.
25. Wang SS, Slanger SL, Brennan P et al. (2007) Family history of hematopoietic malignancies and risk of non-Hodgkin lymphoma (NHL), a pooled analysis of 10211 cases and 11905 controls from the International Lymphoma Epidemiology Consortium (InterLymph). *Blood* 109: 3479-388.
26. Awan H, Jønsson V, Johannesen TB, Ly B, Tjønnfjord GE (2010) Anticipation in families with chronic lymphocytic leukemia and other lymphoproliferative disorders. *TranslOncogenomics* 4: 1-9.
27. Chatenoud L, Gallus S, Altieri A, Negri E, Talamini R, et al. (2005) Number of siblings and risk of Hodgkin's and other lymphoid neoplasms. *Cancer Epidemiol Biomarkers Prev* 14: 552.
28. Jønsson V, Tjønnfjord G, Samuelson SO, Johannesen T, Olsen J, et al. (2007) Birth order pattern in the inheritance of chronic lymphocytic leukaemia and related lymphoproliferative disease. *Leuk Lymphoma* 48: 2387-2396.
29. Von Behren J, Spector LG, Mueller BA, Carozza SE, Chow EJ, et al. (2011) Birth order and risk of childhood cancer: a pooled analysis from five US States. *Int J Cancer* 128: 2709-2716.
30. Report from the Norwegian Cancer Registry
31. Danish Cancer Registry
32. LYFO, Danish Lymphoma Group
33. Emery AEH (1986) Parental age and birth order. In: *Methodology in medical genetics*, Emery AEH (ed) 2nd Edition, Edinburgh: Churchill Livingstone, 140-153.
34. Emery AEH (1986) Segregation analysis. In: *Methodology in medical genetics*, Emery AEH (ed) 2nd Edition. Edinburgh: Churchill Livingstone, 37-54.
35. Hartwell LH, Hood L, Goldberg ML, Reynolds AE, Silver LM, et al. (2000) *Genetics, from genes to genome, genetic analysis according to Mendel*. Boston: McGraw Hill Companies 16-32.
36. Ubeda F, Haig D (2005) On the evolutionary stability of Mendelian segregation. *Genetics* 170: 1345-1357.
37. Hartl DL (1975) Modifier theory and meiotic drive. *TheorPopulBiol* 7: 168-174.
38. Liberman U (1976) Modifier theory of meiotic drive: is Mendelian segregation stable? *TheorPopulBiol* 10: 127-132.
39. Eshel I (1985) Evolutionary genetic stability of Mendelian segregation and the role of free recombination in the chromosomal system. *Am Nat* 125: 412-420.
40. Haig D, Grafen A (1991) Genetic scrambling as a defence against meiotic drive. *J TheorBiol* 153: 531-558.
41. Haig D (2004) Genomic imprinting and kinship: how good is the evidence? *Annu Rev Genet* 38: 553-585.
42. Smallwood SA, Kelsey G (2012) De novo DNA methylation: a germ cell perspective. *Trends Genet* 28: 33-42.
43. Ferguson-Smith AC (2011) Genomic imprinting: the emergence of an epigenetic paradigm. *Nat Rev Genet* 12: 565-575.
44. Morelli MA, Cohen PE (2005) Not all germ cells are created equal: aspects of sexual dimorphism in mammalian meiosis. *Reproduction* 130: 761-781.
45. Sellick GS, Allinson R, Matutes E, Catovsky D, Houlston RS (2004) Increased sex concordance of sibling pairs with chronic lymphocytic leukemia. *Leukemia* 18: 1162-1163.
46. Anway MD, Cupp AS, Uzumcu M, Skinner MK. (2005) Epigenetic transgenerational actions of endocrine disruptors and male fertility. *Science* 308: 1466-1469.
47. Pembrey ME, Bygren LO, Kaati G, Edvinsson S, Northstone K, et al. (2006) Sex-specific, male-line transgenerational responses in humans. *Eur J Hum Genet* 14: 159-166.