

Some Physiological and Biological Studies on Reuterin Production from *Lactobacillus reuteri*

Mohamed MI Helal^{1*}, Amal M Hashem¹, Madeha OI Ghobashy^{2,3} and AI Shima G Shalaby¹

¹Chemistry of Natural and Microbial Products Department, Division of Pharmaceutical and Drug Industries, National Research Center, Dokki, Cairo, Egypt

²Microbiology Department, Faculty of Sciences, Ain Shams University, Cairo, Egypt

³Biology Department, Faculty of Sciences, Tabuk University, Tabuk, Saudi Arabia

Abstract

Lactic Acid Bacteria (LAB) can be used as food preservatives to improve food stability and safety. This is due to its ability to produce antimicrobial substances which can inhibit the growth of the food poisoning organisms. LAB produce antimicrobial compounds named bacteriocins. This study focused on bacteriocin named reuterin which produced from *Lactobacillus reuteri* strain and its optimal production condition. The metabolite *L. reuteri* bacteriocin (reuterin) was extracted and the antimicrobial activity was evaluated against some hospitalized bacterial and fungal pathogens. The reuterin producing *L. reuteri* exhibited the highest inhibition zone (22.2, 22.5 and 22.7 mm) against *E. coli*, *Staphylococcus aureus* and *Candida albicans*, respectively, when grown on optimized condition, i.e., growth on 2% glucose, soy bean (sb) or yeast extract as nitrogen source, all MRS salts medium and inoculated by 21×10^8 cfu/ml, pH 6.5 at 37°C for 24 hr anaerobically. This study gave us the possibility to use reuterin as food preservative to control pathogenic microorganisms and food spoilage.

Keywords: *Lactobacillus reuteri*; Reuterin; Inhibition zone; Amikacin; Fluconazole

Introduction

It is clearly shown that human gut bacteria consists of approximately 10^4 colony forming unit (cfu)/ml and about 500 to 1000 different species and live in their host in symbiosis. This bacterial population is controlled by diets [1]. From few years ago, scientists discovered beneficial bacteria in yogurt, pills and other food and called it probiotics. The Latin and a Greek word “probiotic” means “favorable to life”. According to the Food and Agricultural Organization (FAO) [2]; “they are any living microorganisms administered in adequate amounts which have a beneficial health effect on their host. According to Fuller [3], the probiotics can be defined as “A live microbial feed supplement which beneficially affects the host animal by improving its intestinal microbial balance.” Recently, The word “probiotic” means a large numbers of microorganisms, mainly bacteria (as Lactic acid bacteria (LAB) and non lactic acid bacteria) and few yeasts whom can stay a live until reach to the colon and then improve the host health by its beneficial actions on its host. For example, LAB (e.g., *Lactobacillus reuteri*, *Lactococcus acidophilus*, *Enterococcus faecium*, etc.) and non-lactic acid bacteria (e.g., *Escherichia coli*) also some yeasts (e.g., *Saccharomyces cerevisiae*.) are also considered as probiotics.

These LAB are Gram-positive rods and usually live an-aerobically but in few cases, they can grow also on aerobic conditions [4,5]. Lindgren and Dobrogosz [6] stated that LAB can preserve dairy foods by producing some organic compounds that are antagonistic to other microorganisms. Among these compounds, the bacteriocins which are short chain proteins and have strong ability to inhibit the pathogens growth [7,8]. Many researches had focused on using bacteriocins as food preservatives and control of human bacterial pathogens and disease appearance. The probiotic activity of these bacteria is studied and recorded by Gilliland [9-12]. Bacteriocins production can be influenced by many factors, including medium composition [13] and environmental factors [14,15]. De Man Rogosa and Sharpe (MRS) medium is the standard, specific culture medium for (LAB), but it is not suitable for industrial-scale production due to its high cost. MRS had used for fermentation, while the purpose was to produce

bacteriocins, lactic acid, some enzymes, or other metabolites [16,17]. The lowering of the culture media cost has been published [18-20], not referring to industrial fermentation. The aim of this study was to increase the production and efficiency of the studied reuterin and reduce the fermentation cost to increase the industrial production rate.

Materials and Methods

Bacterial and fungal strains

Lactobacillus reuteri (*L. reuteri*) which is Gram-positive bacillus bacterium kindly provided from Chr. Hansen’s Lab. Inc., Denmark. On other hand, Pathogenic strains, *Escherichia coli*, and *Staphylococcus aureus* in addition to *Candida albicans* as yeast fungi were obtained from the clinical lab. of the El Demerdash Hospital in Cairo, Egypt and they are multi drug resistant (MDR) isolates.

Positive control of antibiotics: It’s include Amikacin (32 µg) as antibacterial antibiotic and Fluconazole (150 µg) as antifungal antibiotic derived from pfizer pharmaceutical company, Cairo, Egypt.

Sugar cane molasses: by product obtained from “The Sugar and Integrated Industries Co.”, El Hawamdia, Giza, Egypt.

Corn steep liquor (CSL) by product was obtained from Egyptian Starch and Glucose Company, Moustoroured, Cairo, Egypt.

Soybean meal-extract (SBE): prepared by aqueous extraction (at 121°C for 20 min.) of a commercial sample of soybean seeds (solid/

***Corresponding author:** Dr. Mohamed MI Helal, Chemistry of Natural and Microbial Products Department, Division of Pharmaceutical and Drug Industries, National Research Center, Dokki, Cairo, Egypt, Tel: +20233371211; E-mail: dmohamedhelal@yahoo.com

Received July 06, 2016; **Accepted** October 07, 2016; **Published** November 07, 2016

Citation: Helal MMI, Hashem AM, Ghobashy MOI, Shalaby AISG (2016) Some Physiological and Biological Studies on Reuterin Production from *Lactobacillus reuteri*. J Prob Health 4: 156. doi: 10.4172/2329-8901.1000156

Copyright: © 2016 Helal MMI, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

liquid ratio 1/20, w/v) followed by filtration and freeze drying of the resulted extract.

Gas generating kits: used for generation of hydrogen and carbon dioxide in metal or plastic anaerobic jars. It was purchased from Oxoid Ltd, Basingstoke, Hampshire, England.

Culture Media

Nutrient agar medium was used for the growth and maintenance of the pathogenic strains *Escherichia coli* and *Staphylococcus aureus* [21].

MRS medium (De Man- Rogosa- Sharp- medium): According to De Man et al. This medium was used for the growth and maintenances of the probiotic *L. reuteri*, it was prepared broth (without addition of agar) and can be used for determination of the growth density of the investigated probiotic and its bacteriocin production [22].

Potato dextrose agar (PDA): Was used for the growth and maintenances of the yeast strain *Candida albicans* [23].

Isolation and purification of bacteriocin

Variations in the level of reuterin production were evaluated by the well diffusion assay [24] and using cork pore diameter 5 mm. Experiments were carried out in duplicate. 0.1% (v/v) of *L. reuteri* was inoculated at a final concentration of about 10^8 CFU/ml according to Emanuel Goldman, Lorrence H Green [25]. The antimicrobial activity of the supernatants was evaluated by the critical dilution assay of Barefoot and Klaenhammer [26] with using of Amikacin (32 µg) as antibacterial antibiotic and Fluconazole (150 µg) as antifungal antibiotic. Reuterin activity was defined as the reciprocal of the highest dilution showing definite inhibition of the indicator strains and was expressed as activity units per milliliter (AU/ml).

Determination of the minimum inhibitory concentration (MIC) of *L. reuteri* supernatant on indicator pathogens: According to Andrews et al. Minimum inhibitory concentration (MIC) is the lowest concentration of an antimicrobial metabolite that inhibits the visible growth of a microorganism after overnight incubation [27]. Supernatants obtained from *L. reuteri* bacterium was used to determine the MIC of each by using the original supernatant of each as follow: 0.025, 0.05, 0.10, 0.15, 0.20 and 0.25 ml. Well-cut diffusion technique

was used to determine the minimal inhibitory effect of the previous dilutions against the indicator pathogens.

Preparation of cell-free culture supernatants: Cell-free culture supernatants (CFCS) were obtained by centrifugation at 5000 rpm for 20 min of *L. reuteri* culture grown under specific cultivation conditions under study.

Optimization studies: It include incubation periods, inoculum size, incubation temperature ranged from 25-45°C and initial pH ranged from 5.0 to 8.5. Effect of various carbon sources, nitrogen sources and medium salt elimination were evaluated in relation to reuterin production.

Data analysis: The recorded data and results were summarized and analyzed in the tables according to Coulombier et al. [28].

Results

Effect of using different incubation times

Table 1 clearly shown that the reuterin production by *L. reuteri* being affected by incubation time, it reached a maximum at 24 and 48 hr. which was 19.0 mm inhibition zone for *E. coli* and 22.0 mm for both *Staph. aureus* or *Candida albicans* comparing to positive control (Amikacin 32 µg/ml as antibacterial agent and Fluconazole 150 µg/ml as antifungal agent) in addition to sterile distilled water as negative control. So 24 hr. is the best incubation time for reuterin production, there for we select 24 hr. incubation period for reuterin production.

Effect of using different initial pH values

According to the nature of the studied organism which is acidophilic, the reuterin production by *L. reuteri* was highly affected by the initial pH values of the media. Eight pH values were selected to study their effects on reuterin production ranged from 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0 and 8.5. All environmental and nutritional factors were fixed and incubated for 24 hr. Table 2 clearly illustrate that the pH 6.5 and 7.0 were the best pH values for maximum reuterin production than that of others under the same experimental conditions.

The maximum production of reuterin is after 24 hr. and pH 6.5 gave 19.0 mm inhibition zone for *E. coli* and 22.0 mm for both *Staph.*

Incubation period (hr)	Cell biomass g/l	Final pH	Inhibition zone (mm)		
			<i>E. coli</i>	<i>Staph. aureus</i>	<i>Candida albicans</i>
24	2.1	5.6	19.0	22.0	22.0
48	2.3	5.6	19.0	22.0	22.0
72	3.5	5.5	18.0	20.0	20.0
96	3.5	5.5	16.0	17.0	17.0
Amikacin 32 µg/ml			20.0	20.0	-
Fluconazole 150 µg/ml			-	-	18.0

Table 1: Effect of using different incubation periods on reuterin production.

pH value	Cell biomass g/l	Final pH	Inhibition zone (mm)		
			<i>E. coli</i>	<i>Staph. aureus</i>	<i>Candida albicans</i>
5.0	1.8	5.0	17.0	16.0	16.0
5.5	1.8	5.0	18.0	18.0	18.0
6.0	2.0	5.5	18.0	18.0	18.0
6.5	2.1	5.5	19.0	22.0	22.0
7.0	2.7	6.0	19.0	21.0	21.0
7.5	2.0	6.3	17.0	18.0	16.0
8.0	1.4	7.0	11.0	11.0	10.0
8.5	1.0	7.1	7.0	8.0	6.0

Table 2: The effect of using different pH values on reuterin production.

aureus or *Candida albicans* comparing to positive control (Amikacin 32 µg/ml as antibacterial agent and Fluconazole 150 µg/ml as antifungal agent).

Effect of using different incubation temperatures

At this experiment, the inoculated test tubes will incubated at 25, 30, 37 and 45°C for 24 hr. and pH 6.5 with fixation of other environmental and nutritional factors and the cell biomass, final pH and reuterin productivity were recorded and listed in Table 3.

The recorded data show that the reuterin productivity was clearly affected by the temperature of incubation. The reuterin productivity was maximized at 30 - 37°C while at elevated temperature (45°C) or low temperature (25°C) the activity decreased. At 25°C the activity decreased to 19 mm inhibition zone diameter while at high temperature (45°C) there was no activity.

Effect of using different inoculum sizes

The Table 4 clearly illustrate the effect of using different inoculum sizes (3, 6, 9, 12, 15, 18, 21, 24 or 27 × 10⁸ CFU/ml) on reuterin production with the fixation of all other environmental or nutritional factors according to the best final condition. According to the Table 4, the best inoculum size gives the highest reuterin production was 21 × 10⁸.

Effect of using different carbon sources

Carbon source is the most important nutritional factor for both growth and reuterin production of *L. reuteri* strain. This study discusses the effect of using of different carbon sources instead of 2% glucose of the reuterin production medium. Carbon sources include; fructose, lactose, sucrose or cane sugar molasses in a concentrations ranged from 1-4%. Table 5 show that the best carbon sources were glucose (2%) and lactose (2%) followed by sucrose while fructose gave weak results on other hand, cane sugar molasses was not suitable for reuterin production (negative results).

Effect of using different nitrogen sources

six nitrogenous compounds were used instead of nitrogenous component of the reuterin production medium, at fixed optimum

conditions (i.e., at pH 6.5, temperature 37°C, inoculum size 21 × 10⁸ with 2% glucose). These nitrogen sources were; peptone, yeast extract, beef extract, casein hydrolysate, SBE and CSL (according to the equivalent N-basis). Table 6 showed that the most favourable nitrogenous compound was yeast extract which gave the highest activity expressed by inhibition zone diameter (The maximum production of reuterin is 22.2 mm inhibition zone for *E. coli* and 22.5 mm for *Staph. aureus* and 22.8 mm for *Candida albicans* comparing to positive controls) followed by corn steep liquor which gave almost the near same reuterin productivity. Results of Table 6 revealed that there is inversely proportionality between reuterin productivity and the cell biomass i.e., the reuterin production will decrease when the cell biomass increased (and vice versa).

Effect of salt elimination

The results illustrated in Table 7 showed that elimination of one salt resulted in decrease of reuterin productivity, so the reuterin productivity resulted from original MRS medium is highly affected by the culture medium component and elimination of one salt will decrease the reuterin productivity. So, all the medium salts are important for reuterin production with concentrations as in MRS medium.

The minimum inhibitory concentration (MIC) test

The results of Table 8 show that, the best volume for culture filtrate (free of cells) containing reuterin is 0.05 ml, below which negative effect of reuterin against studied bacterial and yeast strains was recorded and then it can be considered as the MIC. On the other hand, the volume 0.15 ml as well as 0.2 ml and 0.25 ml of culture filtrate were the best concentrations gives the highest reuterin production and antimicrobial activity against all studied pathogenic strains.

Discussion

This study focused on the effect of two growing factors; environmental and nutritional factors. The effect of environmental factors on reuterin production, including incubation periods, pH values, incubation temperatures and inoculum size were studied by many researchers [29]. On other hand, the nutritional factors (medium composition) play an important role in bacteriocin production [30].

Incubation temperature (°C)	Cell biomass g/l	Final pH	Inhibition zone (mm)		
			<i>E. coli</i>	<i>Staph. aureus</i>	<i>Candida albicans</i>
25	1.8	5.0	19.0	19.0	19.0
30	1.8	5.0	19.0	22.0	22.0
37	2.1	5.5	19.0	22.0	22.0
45	0.4	6.3	-ve	-ve	-ve

-ve: Negative (no reuterin productivity).

Table 3: the effect of using different incubation temperatures on reuterin production.

Inoculum size × 10 ⁸ cfu/ml	Cell biomass g/l	Final pH	Inhibition zone (mm)		
			<i>E. coli</i>	<i>Staph. aureus</i>	<i>Candida albicans</i>
3	1.8	5.8	17.0	18.0	17.0
6	1.9	5.8	18.0	18.0	18.0
9	1.9	5.7	18.0	20.0	19.0
12	2.1	5.5	19.0	20.0	20.0
15	2.1	5.5	19.0	20.0	20.0
18	2.3	5.5	19.0	22.1	22.0
21	2.5	5.5	22.2	22.5	22.8
24	2.5	5.5	22.2	22.5	22.7
27	2.3	5.0	22.2	22.4	22.7

Table 4: The effect of using different inoculum sizes on reuterin production.

Furthermore, reuterin production is strongly affected by medium composition [31]. Our study showed that, the pH values ranged from 6.5 to 7.0 were the optimum pH values gave the highest reuterin productivity with moderate cell biomass of the studied *L. reuteri*. Above or below this pH value will lead to decrease in reuterin productivity. These results agree with Leroy and De Vuyst [32], they proved that the bacteriocin activity of *B. bifidum* was lower at pH ranging from 7.5 to 8.0 and from 5.5 to 6.5.

Generally, reuterin production by *L. reuteri* was considered as a temperature sensitive process. Furthermore, the optimal growth temperature definitely not as like as the optimal bacteriocin production temperature [33] and it consistent with our results which exhibited a temperatures ranged from (30-37°C) were optimum for reuterin production while temperature 37°C only is optimum for cell growth. A higher production of bacteriocin in the range of temperature for optimal strain development has been observed in many LAB strains, such as *E. faecium* RZS C5 [32], *Lactobacillus sakei* Lb 706 [24]. However, no significant differences were reported for bacteriocin production at 30 and 37°C, which may prove that growth temperature in that range does not play a defining role in bacteriocin activity and it coincide with Todorov and Dicks [34] with regards to *L. plantarum* bacteriocins.

The studying of using different nutritional factors on reuterin production aimed to attain higher reuterin yields with very low cost media as food industry wastes (e.g., sugar cane molasses and CSL) or low cost proteins as SBE. With regards to studied carbohydrates, low reuterin production was produced when using sucrose and fructose while sugar cane molasses does not initiate reuterin production for the studied strain, it may be due to that the studied organism did not have the enzymatic system responsible for the digestion of sugar

cane molasses. Since fructose and molasses reduced cell biomass and reuterin productivity, we deduced that the studied probiotic could not utilize these saccharides as a sole carbon source, on the other hand, the sole carbon sources which were readily to be utilized and gave high reuterin productivity were glucose, lactose followed by sucrose. Bing et al. observed that the glucose concentration over 2.0% could reduce the reuterin yield [35]. On the other hand, the nitrogenous compound SBE reduces markedly the reuterin production but the cell growth was too high. It explain that the studied probiotic used the SBE to produce cells only not for reuterin production while CSL gave high reuterin production (more than the control medium) with moderately high cell growth, this agree with Helal et al. [36], although several researchers observed that higher bacteriocin production were recorded when increasing the concentrations of nitrogen content of the medium [37,38]. Salt elimination test explain clearly that all medium salts needed for high reuterin production in a different degrees and the less effective one was $MnSO_4 \cdot 4H_2O$ which gave slight effect.

Finally, our study tested the effect of reuterin concentrations in the culture filtrate on the studied pathogens, it was found that the MIC for activity is 0.05 ml of the reuterin below which there is no antimicrobial activity and concentration 0.15 ml of the culture filtrate is the optimal concentration gave the best antimicrobial activity, this results agree with Helal et al. [36].

Conclusion

This study revealed that the reuterin productivity and cell biomass of the studied *Lactobacillus reuteri* is highly sensitive to environmental and nutritional factors which affecting on the studied probiotic. The relation between the cell biomass and reuterin productivity is not in linear relationship i.e., reuterin productivity is optimized with

Carbon Source	Cell biomass g/l	Final pH	Inhibition zone (mm)		
			<i>E. coli</i>	<i>Staph. aureus</i>	<i>Candida albicans</i>
glucose					
1 %	1.5	6.0	18.0	18.0	18.0
2 %	2.5	6.0	22.2	22.5	22.8
3 %	3.1	5.5	18.0	20.0	20.0
4 %	3.4	5.5	17.0	18.0	16.0
fructose					
1 %	1.2	6.0	10.0	10.0	10.0
2 %	1.3	5.5	11.0	10.0	9.0
3 %	1.3	5.5	10.0	10.0	9.0
4 %	1.3	5.5	10.0	10.0	9.0
Sucrose					
1 %	1.6	6.0	16.0	16.0	16.0
2 %	1.9	6.0	17.0	20.0	20.0
3 %	2.4	5.5	17.0	18.0	17.0
4 %	2.5	5.5	17.0	16.0	16.0
lactose					
1 %	1.5	6.0	18.0	19.0	20.0
2 %	2.5	6.0	19.2	22.0	22.1
3 %	3.1	5.5	18.1	20.0	20.0
4 %	3.4	5.5	17.1	18.0	16.2
Cane sugar molasses					
1 %	0.25	5.0	-ve	-ve	-ve
2 %	0.25	5.0	-ve	-ve	-ve
3 %	0.30	5.0	-ve	-ve	-ve
4 %	0.30	5.0	-ve	-ve	-ve

-ve : negative (no reuterin productivity)

Table 5: The effect of using different carbon sources with different concentrations on reuterin production.

Nitrogen Source	Cell biomass g/l	Final pH	Inhibition zone (mm)		
			<i>E. coli</i>	<i>Staph. aureus</i>	<i>Candida albicans</i>
peptone	1.5	6.0	18.0	18.0	18.0
Yeast extract	2.5	5.5	22.2	22.5	22.8
Beef extract	3.1	5.5	17.0	18.0	18.0
Casein hydrolysate	3.4	5.5	17.0	19.0	19.0
Soy bean	5.8	6.7	11.0	10.2	10.1
Corn steep liquor	3.0	6.5	22.2	22.5	22.7
Control [†]	2.1	5.6	19.0	22.0	22.0

Control[†]: standard MRS broth medium.

Table 6: Effect of using different nitrogen sources on reuterin production (expressed as inhibition zone diameter).

Eliminated salt	Cell biomass g/l	Final pH	Inhibition zone (mm)		
			<i>E. coli</i>	<i>Staph. aureus</i>	<i>Candida albicans</i>
K ₂ HPO ₄	1.5	5.5	10.0	12.0	12.0
Sod. Acetate trihydrate	3.1	5.5	10.0	12.0	11.0
Triammonium citrate	3.4	5.5	10.0	14.0	11.0
MgSO ₄ 7 H ₂ O	5.8	6.7	17.0	17.2	18.1
MnSO ₄ 4 H ₂ O	3.6	6.5	18.0	19.0	19.1
Control [†]	2.1	5.6	19.0	22.0	22.0

Control[†]: standard MRS broth medium.

Table 7: Effect of salt elimination on reuterin productivity.

Culture filtrate conc.(ml)	<i>Escherichia coli</i>	<i>Staphylococcus aureus</i>	<i>Candida albicans</i>
	24 h	24 h	24 h
0.025	-ve	-ve	-ve
0.05	10.0	10.0	12.0
0.10	19.0	22.0	22.0
0.15	22.2	22.5	22.7
0.20	22.2	22.5	22.7
0.25	22.2	22.5	22.7

-ve: negative (no reuterin productivity)

Table 8: The effect of using different culture filtrate concentrations on reuterin production as inhibition zone diameter (mm).

moderate cell growth while increasing the cell biomass did not affecting on the reuterin productivity.

Acknowledgements

All thankful was sent to the National Research Centre, Cairo, Egypt and Ain Shams University, Cairo, Egypt and Tabuk University, Tabuk region, Saudi Arabia, for their support which facilitate making the research practical work.

References

- Lee YK, Salminen S (2009) Handbook of probiotics and prebiotics. 2nd edn. Hoboken, NJ: John Wiley and Sons Inc., USA.
- FAO/WHO (2001) Evaluation of health and nutritional properties of powder milk and live lactic acid bacteria. Food and Agriculture Organization of the United Nations and World Health Organization Expert Consultation Report.
- Fuller R (1989) Probiotics in man and animals. J Appl Bact 66: 365-378.
- Holzappel WH, Haberer P, Geisen R, Björkroth J, Schillinger U (2001) Taxonomy and important features of probiotic microorganisms in food and nutrition. American Journal of Clinical Nutrition 73: 365S-373S.
- Anal AK, Singh H (2007) Recent advances in microencapsulation of probiotics for industrial application and targeted delivery. Trends Food Sci Technol 18: 240-251.
- Lindgren SE, Dobrogosz WJ (1990) Antagonistic activities of lactic acid bacteria in food and feed fermentations. FEMS Microbiol Rev 87: 149-164.
- Ruiz-Barbara JL, Cathcart DP, Warner PJ, Jimenez-Diaz R (1994) Use of *L. plantarum* LPCO10, a bacteriocin producer as a starter culture in Spanish style green olive fermentations. Appl Environ Microbiol, pp: 2059-2064.
- Benkerroum N, Ghouati Y, Ghalfi H (2007) Screening for bacteriocin producing LAB from various Moroccan food products and partial characterization of putative bacteriocins. Biotechnol 4: 481-488.
- Gilliand SE (1990) Health and nutritional benefits from lactic acid bacteria. FEMS Microbiol Rev 87: 175-178.
- Cleveland J, Montville TJ, Nes IF, Chikindas ML (2001) Bacteriocins: safe, natural antimicrobial for food preservation. Int J Food Microbiol 71: 1-20.
- Mojgani N, Ashtiani MP (2006) In vitro inhibition of mastitis pathogens by bacteriocin RN 86 produced by an indigenous strain of *L. casei* RN 86. J Appl Sciences 6: 2629-2634.
- Diez-Gonzalez F (2007) Applications of bacteriocins in livestock. Curr Issues Intestinal Microbiol 8: 15-24.
- Zhou XX, Pan YJ, Wang YB, Li WF (2008) Optimization of medium composition for nisin fermentation with response surface methodology. J Food Sci 73: 245-249.
- Leal-Sánchez MV, Jiménez-Díaz R, Maldonado-Barragán A, Garrido-Fernández A, Ruiz-Barba JL (2002) Optimization of bacteriocin production by batch fermentation of *Lactobacillus plantarum* LPCO10. Appl Environ Microbiol 68: 4465-4471.
- Motta AS, Brandelli A (2008) Evaluation of environmental conditions for production of bacteriocin-like substance by *Bacillus* sp. Strain P34. World J Microbiol Biotechnol 24: 641-646.
- Hummel W, Schütte H, Kula MR (1983) Large scale production of D lactate dehydrogenase for the stereospecific reduction of pyruvate and phenylpyruvate. Eur J Appl Microbiol Biotechnol 18: 75-85.
- Lu Z, Breidt F, Fleming HP, Altermann E, Klaenhammer TR (2003) Isolation and characterization of a *Lactobacillus plantarum* bacteriophage, JL-1, from a cucumber fermentation. Int J Food Microbiol, 84: 225-235.
- Dominguez APM, Bizani D, Cladera-Olivera F, Brandelli A (2007) Cerein 8A production in soybean protein using response surface methodology. Biochem Eng J 35: 238-243.

19. Trinetta V, Rollini M, Manzoni M (2008) Development of a low cost culture medium for sakacin. A production by *L. sakei*. Process Biochem 43: 1275-1280.
20. Wiese B, Bru E, Juárez Tomás MS, Nader-Macías MEF (2010) Optimization of low-cost culture media for the production of biomass and bacteriocin by a Urogenital *Lactobacillus salivarius* strain. Probiotics and Antimicro Prot 2: 2-11.
21. American Public Health Association, American Chemical Society, Association of Official Agricultural Chemists (1920) Standard Methods for the Examination of Water and Sewage. American public health association. pp: 95.
22. De Man JD, Rogosa M, Sharpe ME (1960) A medium for the cultivation of lactobacilli. J appl Bact 23: 130-135.
23. Harold E (1998) Making Bacteria Media from Potato. Indiana Biolab. disknet.com. Retrieved 2011: 03-04.
24. Schillinger V, Luke KK (1989) Antibacterial activity of *Lactobacillus sake* isolated from meat. Appl Environ Microbiol 55: 1091-1096.
25. Emanuel G, Lorrence HG (2008) USA. CRC Press, Taylor and Francis Group, p: 864.
26. Barefoot SF, Klaenhammer TR (1983) Detection and activity of lactacin B, a bacteriocin produced by *Lactobacillus acidophilus*. Appl Environ Microbiol 45: 1808-1815.
27. Andrews JM (2001) Determination of minimal inhibitory concentrations. J Antimicrob Chemoth 48: 5-16.
28. Coulombier D, Fagan R, Hathcock L, Smith C (2001) Epi Info 6 Version 6.04. A Word Processing, Database and Statistical Program for Public Health. CDC, Atlanta, Delaware, USA.
29. Delgado A, López NAL, Brito D, Peres C, Feveireiro P, et al. (2007) Optimum bacteriocin production by *Lactobacillus plantarum* 17.2b requires absence of NaCl and apparently follows a mixed metabolite kinetics. J Biotechnol 130: 193-201.
30. Li C, Bai, JH, Cai ZL, Ouyang F (2002) Optimization of a cultural medium for bacteriocin production by *Lactococcus lactis* using response surface methodology. J Biotechnol 93: 27-34.
31. Ganzle MG, Weber S, Hammes WP (1999) Effect of ecological factors on the inhibitory spectrum and activity of bacteriocins. International Journal of Food Microbiology 46: 207-217.
32. Leroy F, De Vuyst L (2002) Bacteriocin production by *Enterococcus faecium* RZS C5 is cell density limited and occurs in the very early growth phase. International Journal of Food Microbiology 72: 155-164.
33. Leroy F, De Vuyst L (1999b) The presence of salt and a curing agent reduces bacteriocin production by *Lactobacillus sakei* CTC 494, a potential starter culture for sausage fermentation. Applied and Environmental Microbiology 65: 5350-5356.
34. Todorov SD, Dicks LMT (2005) Growth parameters influencing the production of *Lactobacillus rhamnosus* bacteriocins ST461BZ and ST462BZ. Ann Microbiol 55: 283-289.
35. Han B, Yu Z, Liu B, Ma Q, Zhang R (2011) Optimization of bacteriocin production by *Lactobacillus plantarum* YJG, isolated from the mucosa of the gut of healthy chickens. African Journal of Microbiology Research 5: 1147-1155.
36. Mohamed MI, Helal NAM, Abdel-Fattah A, Madeha OI, Ghobashy, et al. (2013) Studies on the production of antimicrobial substances produced from *Bifidobacterium bifidum*. J of applied science research 9: 1554-1563.
37. Aasen IM, Moretro T, Katla T, Axelsson L, Storro I (2000) Influence of complex nutrient, temperature and pH on bacteriocin production by *Lactobacillus sakei* CCUG42687. Appl Microbiol Biotechnol 53: 159-166.
38. Guerra NP, Pastrana L (2001) Enhanced nisin and pediocin production on whey supplemented with different nitrogen sources. Biotechnol Lett 23: 609-612.