

# Spectrum of Congenital Heart Diseases in Children with Down Syndrome in Ajdaybia, Libya

#### Rugea Mahmud Mahmmed\*

Department of Pediatric, Faculty of Medicine, Ajdabiya University, Ajdabiya, Libya

#### ABSTRACT

Down syndrome: Is by far the most common and best known chromosomal disorder in humans.

**Aim of study:** To define the frequency and patterns of congenital heart disease among children with down syndrome in Ejdabyia North eastern of Libya.

**Patients and methods:** This is prospective descriptive study include all phenotypic down syndrome children referred to Echo department at AL-Mugref Teaching Hospital, North eastern of Libya. Verbal consent was taken from guardiants of all participants using GE Vivid S5 Echo Machine.

**Results:** Out of 813 cases 50 (6.1%) case were down syndrome included in our study. 29 (58%) them were female and 21 (42%) were male with Female : Male ratio 1:3:1. 46 (92%) of cases have congenital heart disease. where 40 (87%) have single cardiac defect and the most common defect was AV Canal 27 (54%) case and other form include ASD 6 (12%), VSD 3 (6%), PDA 3(6%) and TOF 1 (2%). the combined lesions were present in 6 (13%) cases include 3(6%) (TOF+AV Canal) and 3 (6%) (VSD+ASD+PDA).

**Conculsion:** Highly prevalence of CHDs among Down syndrome patients with Female dominance. AV Canal was the most common Congenital Heart lesions. So, important to early referral and screening for CHDs in DS. **Keywords:** Down syndrome; CHDs

#### ABBREVIATIONS

DS: Down Syndrome; CHDs: Congenital Heart Diseases; AVC: Atrioventricular Canal; ASD: Atrioseptal Defect; VSD: Ventricular Septal Defect; PDA: Patent Ductal Arteriosus; TOF: Tetralogy Of Fallot.

### INTRODUCTION

**Down syndrome:** Is by far the most common and best known chromosomal disorder in humans and the most common cause of intellectual disability. It is primarily caused by trisomy of chromosomal 21 which gives rise to multiple systematic complications as part of the syndrome [1]. However, not all defects occurs in all patient and there is a wide range of phenotypic variation [2]. It occurs in around 1 in ever 700 pregnancies, the research suggests that there is a higher chance if the mother is older than 35 yrs of age. It is characterized by

intellectual disability, dysmorphic facial features and other distinctive phenotypictraits [3].

The association between Down Syndrome (DS) and Congenital Heart Disease (CHD) was first recognised by Garrod in 1894 and it is the most common cause of death in people with Down syndrome in the first 2 year of life. Therefore, echocardiography is recommended for early detection of CHDs which may help to prevent many complications [4].

**Aim of study:** To define the frequency and patterns of there congenital heart disease among children with down syndrome in Ejdabyia North eastern of Libya.

### MATERIAL AND METHODS

**Study design and sitting:** This is prospective descriptive study include all down syndrome children referred to Echo

**Correspondence to:** Department of Pediatric, Faculty of Medicine, Ajdabiya University, Ajdabiya, Libya, Tel: +218944815824, E mail: sunsetmah@gmail.com

Received: April 20, 2021; Accepted: May 05, 2021; Published: May 12, 2021

Citation: Mahmmed RM (2021) Spectrum of Congenital Heart Diseases in Children with Down Syndrome in Ajdybia, Libya. Clin Pediatr. 6:2319.

**Copyright:** © 2021 Mahmmed RM. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

#### OPEN O ACCESS Freely available online

#### Mahmmed RM

department at AL-Mugref Teaching Hospital, Ejdybia North eastern of Libya. [5]

**Study population:** All potential cases underwent full clinical assessment including phenotypic features that suggested Down Syndrome are included in our study (n=50 cases aged between

2 days to 8 years) with verbal consent were taken from the guardiants of all the participants. Echocardiography examination was performed using a GE Vivid S5 ultrasound Machine [6].

**Data analysis:** Data were exported from collected database to Excel spread sheets in preparation for data analyses [7].

**Results:** Out of 813 cases 50 (6.1%) case were down syndrome included in our study. where 29 (58%) of them were female and 21 (42%) were male. 46 (92%) of cases have congenital heart disease [8].



Data were 40 (87%) have single cardiac defect, the most common defect was AV Canal 27 (54%) case and other form include ASD 6 (12%), VSD 3 (6%), PDA 3(6%) and TOF 1 (2%). the combined lesions were present in 6 (13%) cases include (TOF +AV Canal) 3 (6%) and (VSD+ASD+PDA) 3 (6%).



Figure 2: Distribution of patient according to pattern of CHD.

Where the female predominance seen in cases of AV Canal, PDA, VSD, and (AVCanal+TOF) and TOF. Male dominance seen in (VSD+ASD+ PDA).

The association between Down Syndrome (DS) and Congenital Heart Disease (CHD) was first recognised by Garrod in 1894 and it is the most common cause of death in people with Down syndrome in the first 2 year of life.





#### DISSCUSSION

## Distribution of patient according to presence of congenital heart disease

Congenital heart disease are common in children with Down syndrome where our study showed that: among 813 cases, 50 (6.1%) of them were down syndrome with prevalence of congenital Heart disease among them was 92% and this is going with .The female preponderance with F:M ratio was 1.3: 1. Where 29 (58%) of them were female and 21 (42%) were male and this is going with [9].

### Distribution of patient according to pattern of congenital heart disease

It appears that the distribution and frequency of CHD in DS varies in different geographical regions.

In this study, 40 (87%) of patient have single cardiac lesion and the most common a cyanotic cardiac defect were AV Canal (54%) followed by ASD (12%) and these are going with .Where, the commonest cyanotic cardiac defect was TOF (2%) and this is going with [10]. There are wide difference in combined lesion Types among Down Syndrome patient's studies, in our study 6 (13%) of cases have (AVC+TOF and VSD+ASD+PDA) and no other study going with our study result in this point [11].

**Female preponderance:** AV Canal and this is going with VSD and this is going with. PDA and this is going with. In (AV Canal +TOF) and TOF, no study correlate with my result [12].

Male preponderance: (VSD+ASD+PDA) and this is going with [13].

We certainly are strong proponents of providing high-quality health care for people with Down syndrome [14]. We are certainly not in favor of short-changing a person with Down syndrome. However, the first rule of medicine is "Do no harm." We want to provide optimal care for people with Down syndrome. It is important that we continue to study what that means [15]. Sometimes that means providing the same care as people without DS and sometimes that treatment should be modified based on disease patterns in people with DS [16].

#### LIMITATIONS

- 1. The figures reported herein are not population based and of only one center.
- 2. The cytogenetic studies were not performed and the diagnosis was mainly based on clinical grounds. As a result we could not comment on the frequency of CHD in different chromosomal alterations of DS.
- 3. The frequency was not compared with international rates.

#### CONCLUSION

Prevalence of Down Syndrome in my study was 6.1%. Prevalence of congenital heart disease among Down Syndrome was 92% with Male to Female ratio was 1:1.3.

AV Canal was the most common CHDs among Down Syndrome patient's. Female preponderance among patient with AV Canal, VSD, PDA, TOF and (AVC+TOF) and Male dominance in (VSD+ASD+PDA).

#### RECOMMENDATIONS

- 1. Congenital heart disease is highly prevalence in Down syndrome patients so,
- 2. All children with Down syndrome should have a cardiac evaluation at birth. We stress on the importance of early referral and screening for CHDs in DS.
- 3. The Effort has to be made to do Karyotyping in all suspected cases.
- 4. National registry for Down syndrome and Congenital Heart Disease would provide more accurate data to study contributing factors for CHD in DS.

#### REFERENCES

- 1. Khaira B, Zoubida Z, Mounira A, Naima H, Malika M, Amalou SA, et al. Study of congenital heart disease in patients with down syndrome in Algeria. East Mediterr Health J. 2017;23(9):632-636.
- Levenson D. Talking about down syndrome. Am J Med Gene A. 2009; 149(4).
- 3. Muntha A, Moges T. Congenital cardiovascular anomalies among cases of down syndrome: A hospital based review of cases in tikur an bessa

specialized hospital, Ethiopia. Ethiopia J Health Sci. 2019;29(2):165-174.

- Beatriz E, Bermudez, Sandra L, Medeiros. Down syndrome: Prevalence and distribution of congenital heart disease in Brazil. Sao Paulo Med J 2015;133(6):521-524.
- Mourato FA, Villachan LRR. Prevalence and profile congenital heart disease and pulmonary hypertension in down syndrome in a pediatric cardiology service. Rev Paul Pediatr. 2014;32(2):159-163.
- 6. Elmagrpy Z, Rayani A, Habas E, Aburawi EH. Down syndrome and congenital heart diseas: Why the regional difference as observed the Libyan experience?. Cardiovasc J Afr. 2011;22(6):306-309.
- 7. De Rubens Figueroa J, Del Pozzo Magaña B. Heart malformations children with down syndrome. Rev Esp Cardiol. 2003;56(9):894-899.
- 8. Jaiyesimi O, Baichoo V. Cardiovascular malformations omani Arab children with down's syndrome. Cardio Young. 2007;17(2):166-171.
- 9. Morsy MM, Algrigri O, Salem S, Abosedera MM, Alnajjar AA. The spectrum of congenital heart disease in down syndrome. Saudi Med J. 2016;37(7):767-772.
- Asim A, Agarwal S and Panigrahi I. Frequency of an congenital heart defects in Indian children with down syndrome. Austin J Genet Genomic Res. 2016;3(1):1016.
- Mokhtar M. M, Abdel-Fattah. M. Major birth defects among infants with down syndrome in Alexandria, Egypt: Trends and risk factors. J Trop Pediatr. 2002;48(4):447-249.
- Tereza Cristina Pinheiro Diogenes, Mourato FA, Lima Filho JL, Mattos SS. Gender difference in the prevalence of congenital heart disease in Down syndrome: a brief meta-analysis. BMC Medical Genet. 2017;18(1):111-129.
- Yaqoob M, Manzoor J, Hyder SN, Sadiq M. Congenital heart disease and thyroid dysfunction in Down syndrome reported at children's Hospital, Lahore, Pakistan. Turk J Pediatr. 2019;61(6):915-924.
- Almawazini AM, Sharkawy AA, Eldadah OM, Sumaily YA. Congential heart diseases in down syndrome children at AlbalaArea, Suadi Arabia Suadi Arabia. Neon Pediatr Med. 2017;3(134).
- 15. Shazali O, Ahmed H, El-Shazali H. The spectrum of congenital heart defects in infants with down's syndrome, Khartoum, Sudan. J Pediatr Neonatal Care. 2015;2(5):S00091.
- Chicoine B, Fitzpatrick V, Rivelli A, Bria A. Down syndrome and heart disease between 1996 and 2016. J Am Board Fam Med. 2020; 33(6):923-931.