

Stress and Anxiety A Scientific View of Mechanisms Effects and Management Approaches

Noah Bennett*

Department of Psychiatry, Monash University, School of Clinical Sciences Melbourne, Australia

DESCRIPTION

Stress and anxiety are common psychological experiences that influence human behavior emotions and overall health. The interaction between biological processes psychological patterns and environmental conditions determines how individuals experience and manage stress and anxiety. Stress and anxiety are increasingly recognized as major concerns in modern society [1]. Daily responsibilities social pressures academic demands workplace challenges and personal uncertainties can create emotional strain that affects individuals across different age groups. Although stress and anxiety are often used together they represent different psychological processes. Stress generally develops as a response to external demands while anxiety involves feelings of fear worry or anticipation about possible threats [2]. A certain level of stress and anxiety can be beneficial because they improve alertness motivation and problem solving abilities. However when these responses become excessive persistent or difficult to control they may contribute to emotional exhaustion and reduced quality of life. Understanding the relationship between stress and anxiety is essential for developing effective strategies to support mental health [3]. The relationship between stress and anxiety is strongly influenced by biological mechanisms involved in threat detection and emotional regulation. This process involves the nervous system and hormonal pathways that regulate physiological reactions.

The hypothalamic pituitary adrenal axis plays an important role in stress regulation by controlling the release of stress related hormones. Short term activation of this system supports adaptation but prolonged activation may contribute to emotional imbalance and increased anxiety symptoms [4]. Brain regions involved in emotional processing including the amygdala and prefrontal cortex influence how individuals perceive and respond to stressful experiences. The amygdala helps detect potential threats while the prefrontal cortex contributes to emotional control and decision making. Imbalance between these systems may increase sensitivity to stress and anxiety [5]. Research approaches such as in vitro studies help examine cellular responses related to stress while in vivo investigations

provide information about biological reactions within living systems. Computational methods using in silico approaches allow researchers to explore complex interactions involved in stress and anxiety regulation. Psychological factors strongly influence how individuals experience stress and anxiety. Two people exposed to similar circumstances may experience different levels of stress depending on their psychological resources [6].

Persistent stress can influence thought patterns and emotional stability. Individuals experiencing chronic stress may develop excessive worry difficulty concentrating and feelings of being overwhelmed. Developing effective coping strategies is important for maintaining emotional balance [7,8]. Techniques that improve self awareness relaxation problem solving and adaptive thinking can help individuals manage stressful situations more effectively. Psychological support can also provide valuable tools for understanding and controlling anxiety responses. Long term stress and anxiety can affect multiple aspects of health. Continuous emotional pressure may influence sleep quality energy levels immune function and cardiovascular health. Stress and anxiety may also influence behavior by reducing motivation affecting social interactions and decreasing productivity [9]. Individuals experiencing prolonged anxiety may avoid certain situations which can further limit personal growth and daily functioning. The effects of stress and anxiety highlight the importance of viewing mental health as a complete system involving emotional biological and social factors. Supporting mental well being requires attention to both psychological experiences and physical health. Environmental conditions strongly shape stress and anxiety levels. Supportive relationships and positive social connections often reduce the negative effects of stress [10].

CONCLUSION

Modern lifestyles have introduced additional sources of psychological pressure including constant digital communication information overload and reduced opportunities for relaxation. However technology also provides new possibilities for mental

Correspondence to: Noah Bennett, Department of Psychiatry, Monash University, School of Clinical Sciences Melbourne, Australia, E-mail: noahbennett@gmail.com

Received: 01-Apr-2025, Manuscript No. JDA-25-42927; **Editor assigned:** 03-Apr-2025, PreQC No. JDA-25-42927 (PQ); **Reviewed:** 17-Apr-2025, QC No. JDA-25-42927; **Revised:** 25-Apr-2025, Manuscript No. JDA-25-42927 (R); **Published:** 01-May-2025, DOI: 10.35258/2167-1044.25.14.571

Citation: Bennett N (2025). Stress and Anxiety A Scientific View of Mechanisms Effects and Management Approaches. J Dep Anxiety. 14:571.

Copyright: © 2025 Bennett N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

health education online support and access to resources. Creating healthier environments requires collective efforts from individuals communities and healthcare systems. Promoting awareness and reducing stigma can encourage people to seek support before stress and anxiety become overwhelming.

REFERENCES

1. Algate DL, Beck C, Kolanowski A, Whall A, Berent S, Richards K, et al. Need-driven dementia-compromised behavior: An alternative view of disruptive behavior. *Am J Alzheimer's Dis.* 1996;11(6):10-19.
2. Hsiao HC, Chao HC, Wang JJ. Features of problematic eating behaviors among community-dwelling older adults with dementia: Family caregivers' experience. *Geriatr Nurs.* 2013;34(5):361-365.
3. Kales HC, Gitlin LN, Lyketsos CG. Assessment and management of behavioral and psychological symptoms of dementia. *BMJ.* 2015;350.
4. Kovach CR, Noonan PE, Schlidt AM, Wells T. Need-driven dementia-compromised behavior: An alternative view of disruptive behavior. *J Nurs Scholar.* 2005;37(2):134-140.
5. Reis RC, Dalpai D, Camozzato A. Staff training to reduce behavioral and psychiatric symptoms of dementia in nursing home residents: A systematic review of intervention reproducibility. *Dement Neuropsychol.* 2013;7:292-297.
6. Sampson EL, White N, Leurent B, Scott S, Lord K, Round J, et al. Behavioural and psychiatric symptoms in people with dementia admitted to the acute hospital: Prospective cohort study. *Br J Psychiatr.* 2014;205(3):189-196.
7. Shih YH, Pai MC, Huang YC, Wang JJ. Sundown syndrome, sleep quality, and walking among community-dwelling people with Alzheimer disease. *J Am Med Director Assoc.* 2017;18(5):396-401.
8. Wang CJ, Pai MC, Hsiao HS, Wang JJ. The investigation and comparison of the underlying needs of common disruptive behaviours in patients with Alzheimer's disease. *Scand J Caring Sci.* 2015;29(4):769-775.
9. Wang JJ, Cheng WY, Lai PR, Pai MC. Delusions and underlying needs in older adults with Alzheimer's disease: Influence of earlier life experiences and the current environment. *J Gerontol Nurs.* 2014;40(12):38-47.
10. Wang JJ, Feldt K, Cheng WY. Characteristics and underlying meaning of hoarding behavior in elders with Alzheimer's dementia: Caregivers' perspective. *J Nurs Res.* 2012;20(3):189-196.