

Suicidal Behavior and Neurological Illnesses

Coughlin Steven S^{1,2} and @c Sher^{3,4}

¹Post-Deployment Health Epidemiology Program, Office of Public Health, Department of Veterans Affairs, Washington, DC, USA

²Current Affiliation: Department of Epidemiology, Rollins School of Public Health, Emory University, Atlanta, GA, USA

³James J. Peters Veterans' Affairs Medical Center, New York, USA

⁴Mount Sinai School of Medicine, New York, USA

Abstract

Objective: Suicidal ideation and behavior have been associated with a variety of neurological illnesses. Studies are ongoing in combat veterans and other groups to examine possible mechanisms and pathways that account for such associations.

Method: This article provides a review of the literature on suicide ideation and suicidal behavior in patients with neurological illnesses including publications on veteran's health and military medicine. Studies of suicide attempts and deaths in people with neurological illnesses are also reviewed.

Results: The studies summarized in this review indicate that there are important linkages between suicidal ideation and behavior and neurological conditions, including epilepsy, multiple sclerosis, and amyotrophic lateral sclerosis.

Conclusion: Additional studies are needed to further clarify why suicide ideation and suicidal behavior are associated with neurological diseases, in order to improve quality of life, alleviate patient distress, and prevent nonfatal and fatal suicide attempts in veteran and non-veteran populations.

Keywords: Amyotrophic lateral sclerosis; Epilepsy; Multiple sclerosis; Suicide; Veterans

Introduction

Over 36,000 people in the U.S. die each year from suicide. A much larger number of people attempt suicide, but survive the experience. An estimated 25 suicide attempts occur for every fatal suicide [1]. Suicide is more frequent among men whereas non-fatal suicidal behaviors are more prevalent among women and in persons who are unmarried or have a psychiatric disorder [2]. Psychological autopsy studies have shown that more than 90% of people who commit suicide suffer from psychiatric disorders [3,4]. Demographic risk factors for suicidal ideation, developing a suicide plan, and non-fatal suicidal behaviors include being female, being younger, having less educational attainment, and being unemployed [5]. The study of suicidality and suicide prevention programs are important aspects of post-deployment health and military medicine [6]. About 20% of all known suicides in the U.S. occur among current or former military personnel [7].

Although it is widely recognized that suicide ideation and suicidal behaviors occur more frequently among persons with certain psychiatric illnesses (e.g., major depression, bipolar disorder, personality disorders, schizophrenia, substance abuse) [3,8-10], additional factors are likely to account for the transition from suicidal ideation to suicide plans or attempts [4]. For example, lack of psychosocial support from family and friends, social disadvantage, psychosocial stress due to financial problems or a disrupted relationship, and the absence of a therapeutic alliance with a clinical provider can be detrimental [1].

Previous authors have noted that suicide ideation and suicidal behaviors can be related to an individual's physical illness and to unhealthy behaviors such as heavy alcohol consumption [11-13]. Suicidal ideation and behavior have been associated with a variety of neurological illnesses, although relatively few attempts have been made to examine possible mechanisms and pathways that account for such associations, while comparing and contrasting key pathways across various neurological conditions.

The objective of this article is to review the recent literature on suicidal ideation and behavior occurring among adults suffering from neurological illnesses including the literature on veteran's health, including epilepsy, multiple sclerosis, and amyotrophic lateral sclerosis. The general approach used to identify references cited in this article was to identify relevant medical subject headings (MeSH) used by the National Library of Medicine and to search PubMed bibliographic databases for appropriate articles. Of particular interest were clinical and epidemiologic studies of suicidal ideation and suicidality published since 2000. Clinical descriptions of individual patients and case series involving very small numbers of patients were omitted. The bibliographic searches were limited to studies published in English. However, because suicidality and neurological illnesses are global health concerns, the bibliographic searches were not limited to studies carried out in the U.S. Additional citations were identified by reviewing systematic reviews and review articles identified through PubMed searches (for example, recent review articles on the epidemiology of suicidality). The equally important topic of suicidality and Traumatic Brain Injury (TBI), spinal cord injury, and polytrauma will be examined in a future article. In order to narrow the focus of the current review, suicide ideation and behavior occurring in Parkinson's disease and Alzheimer disease are not considered.

Overview of Suicidal Ideation and Suicidal Behaviors

Suicidal ideation and suicidal behaviors can be conceptualized as occurring over a continuum from suicidal ideation, development

***Corresponding author:** Steven S Coughlin, 1648 Beverly Drive, Carson City, NV 89706, USA, Tel: (404) 983-2524; E-mail: stevecatlanta@aol.com

Received March 02, 2013; **Accepted** March 18, 2013; **Published** April 18, 2013

Citation: Coughlin Steven S, Sher L (2013) Suicidal Behavior and Neurological Illnesses. J Depress Anxiety S9: 001. doi:[10.4172/2167-1044.S9-001](https://doi.org/10.4172/2167-1044.S9-001)

Copyright: © 2013 Coughlin Steven S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

of a suicide plan, non-fatal suicide attempt(s), or death from suicide [5]. Suicides reflect only a small proportion of the total impact of suicidal behavior; substantially more persons are hospitalized as a result of nonfatal suicidal behavior than are fatally injured. An even greater number are either treated in ambulatory settings or not treated at all. Nock et al. [2] defined nonfatal suicidal thoughts and behaviors (i.e., suicidal behaviors) as suicide ideation (which refers to thoughts of engaging in behavior intended to end one's life), suicide plan (the formulation of a specific method through which one intends to die), and suicide attempt (engagement in potentially self-injurious behavior in which there is at least some intent to die). Nonsuicidal self-injury such as self-cutting can also occur in which a person has no intent to die [2]. While suicidal behavior and nonsuicidal self-injury are distinct behaviors, nonsuicidal self-injury is a risk factor for suicide attempts and suicide. Roughly 55-85% of self-injurers have made at least one suicide attempt [14,15]. The suicide rate for patients with borderline personality disorder and nonsuicidal self-injury is about 8-10% [16,17]. Community surveys of suicide behaviors have sometimes categorized suicidal thought or behavior into 4 groups of increasing severity: suicide ideation only, suicide plan, unplanned suicide attempt and planned suicide attempt [18]. The Department of Veterans Affairs recently adopted the U.S. Centers for Disease Control and Prevention's Self-Directed Violence Classification System nomenclature to facilitate research and epidemiologic surveillance efforts [19].

Retrospective studies suggest that suicidal ideation fluctuates and that the process that people may go through before they attempt suicide is complex [20]. Suicidal thoughts and behaviours tend to be transient in nature, coming and going repeatedly over time [18]. Mathias et al. [21] noted that suicidal ideation should be thought of as a state that would be expected to fluctuate in intensity over time rather than as a psychological trait. Suicide attempters are distinguished from nonattempters by specific traits and by a family history of suicidal behavior. Suicidal behavior is usually associated with mood and anxiety disorders, borderline and antisocial personality disorders, alcohol and drug dependence, and greater lifetime aggression and impulsivity in attempters than in nonattempters [22-24]. A further issue is that some differences have been noted between the characteristics of persons who attempt suicide and those who complete suicide [25].

The risk of nonfatal self-injury (both suicidal and nonsuicidal in nature) rises significantly in the U.S. during adolescence and young adulthood and then decreases monotonically throughout adulthood [2]. Rates of non-fatal self-injury are higher among females than in males, in contrast to a pattern of higher suicide rates among males. Only a small percentage of people who attempt suicide eventually complete suicide, although suicide attempts are an indicator of extreme psychological distress and an important predictor of eventual completed suicide [5]. In fact, the best predictor of suicide is a history of a previous suicide attempt. It has also been observed that more severe methods of attempted suicide confer greater risk of completed suicide [26].

Risk factors for attempted suicide include suicide ideation, developing a suicide plan, prior suicide attempts, hopelessness, having a positive family history of suicide or psychiatric illness, and recent loss of an important relationship [1]. Suicide ideation and suicidality have been related to an individual's psychiatric or physical illness [2,11]. In over 90% of suicides, this fatal outcome is a complication of a psychiatric disorder. Sixty percent of all suicide victims have a mood disorder at the time of death [22,27-29]. Personally knowing somebody who completed or attempted suicide is associated with an increased risk of experiencing suicidal ideation or attempting suicide, especially

among young people [20,30]. Other risk factors for suicide attempts and death from suicide include lack of social support, unemployment, and low income [31]. Social disadvantage, financial distress, and legal difficulties have been identified as risk factors for suicidal ideation and precipitants of suicidal acts [20,32,33].

The lifetime prevalence of suicide ideation has been estimated to be 5.6-14.3 percent [2]. The 12-month prevalence of suicide ideation has been estimated to be 2.1-10.0 percent. The variation in rates is likely due, in part, to variability in the methods used to assess suicidal behaviors [2]. Bossarte et al. [34] recently examined the prevalence of suicide ideation among active military and veteran participants in a national health survey in the U.S. The prevalence of suicide ideation and attempts identified in their study were similar to published estimates of those outcomes in the U.S. general population, 3.7% and 0.5%, respectively. In a study of Operation Enduring Freedom/Operation Iraqi Freedom (OEF/OIF) veterans from Connecticut, Pietrzak et al. [35] found that 12.5% (34 of 272) of the respondents reported contemplating suicide in the 2 weeks prior to completing the survey. The veterans who reported suicidal ideation were more likely to screen positive for post-traumatic stress disorder (PTSD), depression, or problem drinking [35]. They also scored higher on measures of psychosocial difficulties, stigma, and barriers to care, and lower on measures of resilience and social support. Other studies have shown that suicidal ideation is associated with combat exposure [36]. Among the highest published rates of suicidal ideation (63%) are those from a study of homeless men and women [37].

Ideation as a risk factor for suicide attempt or death: Persons who experience suicidal ideation, especially those who have comorbid conditions such as major depression, bipolar disorder, alcohol abuse or dependence, or chronic pain, have a higher risk of nonfatal or fatal suicide attempt [38-40]. Suicidal ideation has been associated with increased risk of suicidal behavior in community-based studies [12,20,40] and in follow-up studies of patients seen for psychiatric care [31,41]. Combat veterans are more likely to have suicidal ideation (often associated with PTSD and depression) and they may be more likely to act on a suicidal plan [42]. A recent study by Britton et al. [38] identified suicidal ideation as a warning sign for suicide among U.S. veterans who died within 1 week of receiving healthcare from the Veterans Health Administration (VHA).

There is likely to be variability in the extent to which suicidal ideation predicts suicide in clinical and community samples. In a 12-year follow-up study of a community sample in Baltimore, Maryland, Kuo et al. [40] found that persons who reported suicidal ideation at baseline were more likely to report having attempted suicide at follow-up (relative risk=6.09, 95% confidence interval 2.58-14.36). In a longitudinal study in a community sample, adolescents who had suicidal ideation at age 15 were more likely to have attempted suicide by age 30 [33]. An estimated 34 percent of persons who experience suicidal ideation go on to make a suicide plan and about 72 percent of persons with a suicide plan go on to make a suicide attempt [2,43]. An estimated 26 percent of persons who experience suicidal ideation make an unplanned suicide attempt.

A variety of preventive factors have been identified that can help protect people with suicidal ideation from attempting suicide. These include family responsibilities and child-related concerns, fear of suicide, fear of social disapproval, and life-affirming cultural values [1]. Concern shown by others has been inversely associated with suicidal ideation [44]. Psychosocial support from family, friends, and community, and having a sense of purpose, have been reported to protect against suicidal behavior among OEF/OIF veterans who may

have PTSD [35]. Skills in problem-solving and having a therapeutic alliance with a clinical provider can also be important [1]. Resilience to suicidal behavior is influenced by genetic factors, parenting, psychiatric and medical illnesses, especially affecting the brain, e.g., epilepsy, migraine, Huntington's disease, alcoholism and substance abuse [22,27,45-49]. The relationships may be in different directions; genetic factors may both increase or decrease resilience to suicidality; illnesses affecting the brain can increase suicide risk. Important suicide protectors include effective clinical care for physical and psychiatric disorders (including substance abuse); easy access to a variety of clinical interventions and support for help seeking; restricted access to highly lethal means of suicide; strong connections to family and community support, and cultural and religious beliefs that discourage suicide and support self preservation [50].

Assessment: The identification of people who are having thoughts of suicide or of harming themselves in some way, and ensuring their access to mental health care services and appropriate follow-up, has been highlighted in the psychiatric, medical, and public health literature [7]. Suicidal ideation can be a highly distressing experience and is a valid target for clinical intervention [51]. Such ideation can be passive, fleeting, intermittent, active, and intense, with or without the intent to die [1]. As part of clinical evaluation and suicide risk assessment, suicidal ideation can be distinguished from suicidal intent. The latter is the subjective expectation and desire to die by a self-destructive act [1].

Suicide assessments consider such factors as previous suicide attempts, mood and anxiety symptoms, coexisting psychiatric disorders including alcohol and drug abuse, the role of any medications (stimulants, antidepressants, other), and the lack of family and social support. The presence of a suicide plan or the possibility of implementing a suicide plan is also considered.

Studies of suicide assessment tools such as the Scale for Suicidal Ideation and the Beck Hopelessness Scale have shown associations between higher scores and death by suicide [7]. The utility of such tools in routine screening in primary care practice is uncertain because of the potential for false positive and false negative identification of persons at risk of suicide and because busy primary care practitioners do not have time to administer these or other scales [7]. Suicide risk assessments performed as part of psychiatric care are determinations made for the purpose of treatment and patient management and are not long-term predictions of suicide risk [1]. Suicide risk assessment has a quasi-quantitative dimension in that risk of suicide increases with various combinations of risk factors [1].

Epilepsy

Epilepsy affects about 3 million Americans. About 200,000 new cases of epilepsy and seizures occur each year in the U.S. A complex relationship exists between epilepsy and suicidality [52,53]. Epilepsy both increases the risk of the development of psychiatric conditions (for example, depression) and the risk of suicide among people with psychiatric disorders [52]. In addition to psychiatric illness, risk factors for suicide among people with epilepsy include temporal-lobe epilepsy, female sex, and the onset of epilepsy at an earlier age [52]. People with epilepsy have a higher risk of suicide ideation and suicidal behaviors than the general population [54]. Suicide risk increases soon after the onset of epilepsy and then gradually decreases [52]. Onset of epilepsy in the first 17 years of life is associated with a significant increase in risk for suicide compared with those whose epilepsy started at an age of more than 29 years [53]. The relation is one of mutual causation since injuries sustained during a suicide attempt can injure the brain and cause

epileptic seizures. A further issue is that epilepsy and suicidal ideation may share common neurobiological mechanisms such as disturbances of neurotransmitters [54]. For example, serotonergic disturbances are present in both epilepsy and suicidal behavior. Antiepileptic drugs have been associated with increased risk of suicidality by some regulatory agencies, although the finding has not been supported by some studies and meta-analyses [52,55-57]. The U.S. Food and Drug Administration requires that labeling of antiepileptic drugs include a warning and that a medication guide be provided to patients, informing them of the risks of suicidal thoughts or actions. The existence of a causal association between antiepileptic medications and suicidality has been a topic of continuing scientific discussion and debate. One reason for the elevated risk of suicide in people with epilepsy might be social disadvantages and stigma associated with epilepsy.

Post-traumatic forms of epilepsy are especially important in combat veterans who are more likely to experience mild, moderate, or severe traumatic brain injuries, PTSD, and major depression than other veterans [58-60]. Epileptic seizures, PTSD, depression and suicide ideation and attempts are more frequent among persons with traumatic brain injury [58,59,61]. People with epilepsy should be regularly evaluated for depression and suicidality by their physician, and when depression or suicidality is present, therapeutic interventions are necessary.

Multiple Sclerosis

Multiple sclerosis (MS) is a chronic neurological disease with a variable clinical course [62]. The pathophysiological mechanisms include neuronal/axonal damage and loss, demyelination, inflammation, remyelination and repair, and gliosis; the latter refers to a proliferation of astrocytes in damaged areas of the central nervous system [62]. The axonal loss and gliosis that occurs in MS can result in a progression of disability. The disability experienced by MS patients varies widely. The overall life expectancy for MS patients has been estimated to be 6-7 years less than that of persons without the disease [63]. Estimates of the annual incidence of MS vary by country, sex, race, and history of military service [64-66]. Women have a higher risk of MS than men. Among U.S. Armed Forces personnel from 2000-2009, the annual incidence rate was 12.9 per 100,000 person-years [67]. Black non-Hispanics had a higher incidence rate (18.3 per 100,000 person-years) than white non-Hispanics (12.5 per 100,000 person-years) [67]. Because many MS patients do not die from their chronic illness, estimates of the prevalence of MS are much higher. For example, Noonan et al. [68] found that the prevalence of MS in 3 US communities ranged from 47.2-109.5 per 100,000 persons. In a recent study from France the prevalence of MS was estimated to be between 170.9-188.2 cases per 100,000 persons [69]. Epidemiologic and genetic studies have shown that MS likely results from a complex interaction between genetic susceptibility and environmental factors [70]. Genetic studies have confirmed that the immune system plays a key role in the development of MS [71].

Depression is common in patients with MS and other neurodegenerative diseases [72-75]. Paparrigopoulos et al. [76] postulated that depression occurring in MS may be due to a complex interplay of biological, disease-related, behavioral, and psychosocial factors. The neuroinflammatory reaction is involved in the pathophysiology of MS. Immune sensitization to neural antigens could develop within the systemic compartment consequent to exposure to cross-reacting peptides and play an important role in the neurobiology of MS [73,74]. The etiology of depression in MS patients

may be due to a psychological reaction to the diagnosis or associated disability, to neuroanatomical or neurochemical changes associated with neurodegeneration, or to psychiatric side effects of corticosteroids or interferons used to treat MS [77-79]. Factors that may contribute to depression among MS patients include maladaptive coping strategies, loss of recreational opportunities, having poor quality relationships, and high levels of stress and fatigue [80]. In evaluating depression occurring in patients with a neurodegenerative disease such as MS, providers should draw a distinction between the syndrome of major depression (as defined by the Diagnostic and Statistical Manual of Mental Disorders, 4th revision) and symptoms of MS itself [72]. The lifetime prevalence of major depression in MS patients may approach 50% [72].

Population-based studies have generally shown that people with MS have about a two-fold increased risk of suicide, after adjustment for age [81-83], although no significant association was observed between suicide and MS in a study conducted in Finland between 1964 and 1993 (standardized mortality ratio=1.7, 95% confidence interval 0.9-2.7) [84]. Suicide risk may be particularly high in the first year following the diagnosis of MS, and among young men [81]. A recent study found that 3.7% (15 of 404) of MS patients in Denmark had attempted suicide; the rate was not significantly different than that of the general population of the County of Funen, Denmark [85].

Over a quarter of MS patients may contemplate suicide [72,86]. In a clinic-based study of 140 MS patients in Toronto, Canada, Feinstein [86] found that 28.6% (40 of 140) of patients had thought about committing suicide and 6.4% (9 of 140) had attempted suicide. Those with a history of suicidal intent were more likely to have a history of major depression than the non-suicidal group (83% versus 18%). Factors associated with suicidal intent among MS patients included living alone, having a positive family history of mental illness, reporting more social stress, and having a lifetime history of major depression, anxiety disorder, or alcohol abuse [86]. Turner et al. [39] studied suicidal ideation in 445 VHA patients with MS by linking computerized medical records with information obtained via a mail survey. Suicidal ideation was assessed using the PHQ suicide item. About 29.4% (131 of 445) of respondents had suicidal ideation and 7.9% (35 of 445) had persistent suicidal ideation (more than half the days) over the past 2 weeks. In bivariate analyses, suicidal ideation was associated with younger age, earlier disease course, progressive disease subtype, lower income, not being married, having less social support, not driving, having higher levels of physical disability, and depression [39].

Previous authors have noted that providers who care for MS patients should closely monitor their patients for depression and suicide ideation, that MS patients with suicidal ideation should be referred for immediate psychiatric consultation, and that depression in MS patients can be effectively treated with antidepressant medication or psychotherapy [72,76,78,87]. Current and emerging MS therapies that may show promise across the spectrum of disease have recently been reviewed by Castro-Borrero et al. [88] and Bompreszi et al. [89]. It is important to make sure that new therapies will not be associated with suicidality.

Amyotrophic Lateral Sclerosis

Amyotrophic lateral sclerosis (ALS) is a devastating disease characterized by progressive loss of motor neurons [90]. In the U.S., the prevalence of ALS is about 5 cases per 100,000 persons, with somewhat higher rates among men. Most people with ALS are between the ages of 40 and 70. About 5% of ALS cases are familial [91]. Considerable

evidence from both patients and animal models indicates impairment of all neurovascular unit components including the blood-brain and blood-spinal cord barriers [92]. The neurodegeneration seen in ALS can occur within the corticospinal tracts, brain stem nuclei, and spinal cord anterior horns [93]. The neurodegeneration can extend beyond motor cerebral neurons into the frontal and temporal lobes although frank dementia is rare [93]. Patients often experience weakness of limb, abdominal, thoracic, or bulbar muscles. The latter are muscles of the mouth and throat that are responsible for speech and swallowing. The one relatively effective drug for ALS, Riluzole, a medication that preferentially blocks tetrodotoxin-sensitive sodium channels which reduces influx of calcium ions and probably prevents stimulation of glutamate receptors [94], prolongs survival for up to 6 months [95,96]. Currently, most therapeutic efforts for ALS patients are centered on symptom management, including such important areas as respiration, nutrition, secretions, communication, physical therapy and exercise, muscle cramps and spasticity, pain, and depression [97]. Although ALS patients often die within 3 to 5 years from diagnosis about 10% of patients survive for more than 10 years [98].

Palmieri et al. [99] examined suicidal ideation in 42 patients with ALS. The Rorschach test was used in the assessments. The investigators found that suicidal ideation was more frequent in recently diagnosed ALS patients [99]. A population-based study in Sweden identified 6,642 patients with incident ALS between 1965 and 2004 using the Swedish Inpatient Register [90]. The investigators found an almost 6-fold increased risk for suicide among ALS patients (standardized mortality ratio 5.8, 95% confidence interval 3.6-8.8). Patients who committed suicide were on average about 7 years younger at the time of their first hospitalization as compared with those who did not commit suicide.

It is important to note that ALS is associated with cognitive impairments [100]. Cognitive abnormalities are associated with suicidal behavior [101,102]. Therefore, cognitive impairments may mediate the effect of ALS on suicidality.

In a study by Rabkin et al. [103] of 80 patients with late-stage ALS, about 8-9% of patients had major depression at the time of any visit. The frequency of depression did not generally increase as death approached [103]. More severe depressive symptoms and greater hopelessness were correlated with higher scores on measures of suffering, anger, perceived caregiver burden, and weariness [103]. During the study, 14 patients received antidepressant medication and 6 saw a counselor or therapist [103]. The authors noted that depression affects only a small subset of patients with late-stage ALS and that the overall results from their study provide a "picture of resiliency rather than despair" [103]. Lule et al. [104] conducted two studies of depression and quality of life among ALS patients. One study was longitudinal in design and the other compared ALS patients with healthy controls. About 28% of the patients had signs of depression based upon the ALS Depression Inventory; the severity of depression was inversely related to educational status [104]. Depressive symptoms were not found to be correlated with extent of physical impairment or time since diagnosis [104]. A measure of quality of life was similar among ALS patients and healthy controls, indicating that ALS patients can experience a satisfactory quality of life without manifestations of depression even if they are severely physically impaired [104]. Additional studies of the psychological health of patients with ALS have also found that clinically significant depression is not as frequent or severe as might be expected (based upon studies of other life-threatening medical illnesses), although the psychological impact of ALS on patients, family, and caregivers is profound [105,106]. Several studies have shown that ALS patients often report surprisingly

high subjective quality of life despite their marked physical impairment [104,107].

Summary and Conclusions

The results of this literature review highlight the importance of an awareness of the psychological distress experienced by patients with neurological conditions such as, epilepsy, MS, and ALS. Scientific understanding of the psychoneurobiological mechanisms and pathways that account for suicidal ideation and suicidal behaviors in patients with neurological disorders is still in its infancy. Measures of executive dysfunction have been associated with suicidal behavior in some neuropsychological studies of research participants with a history of suicidal behavior and healthy controls [108]. Twin-studies and other family studies have provided evidence of a heritable risk of suicidal behavior [2,109-115]. Much of the family history of suicidal behavior may be explained by the risk associated with psychiatric conditions such as major depression, bipolar illness, and schizophrenia.

Advances in proteomics, genomics, and metabolomics, and ongoing and planned clinical trials for neurorestoration within the human central nervous system, offer hope for patients suffering from MS, ALS, and for their family members [98,116-118].

Researchers have examined specific psychological constructs that may explain why neuropsychiatric disorders are associated with suicidal behaviors. These constructs include the presence of hopelessness, anhedonia, impulsiveness, and high emotional reactivity [2]. A variety of psychological and psychoanalytic models and theories of suicidal behaviors have been proposed. For example, in the Interpersonal Theory of Suicide, interpersonal connectedness is posited to be a protective factor for suicidal behavior [119]. The presence of a psychiatric disorder is a consistently reported risk factor for suicidal ideation and behavior [2].

Physical illnesses and chronic pain are important risk factors for suicidal ideation and behaviors. Epilepsy can occur during childhood, adolescence, and young adulthood, or later in life, whereas neurodegenerative diseases such as Parkinson's disease, stroke, Alzheimer disease, and dementia—which were outside the scope of the current review—are primarily illnesses that afflict the elderly. Multiple sclerosis and ALS may also occur in the elderly and in middle-aged persons. In people of all ages, physical illnesses, depression, and loss of important relationships are important risk factors for suicidal behaviors. Primary care providers can play an important role in suicide prevention among higher-risk patients including the elderly [2]. In a study by Isometsa et al. [29], 41% of older adults had seen their primary care physician within 28 days of committing suicide. A meta-analysis of 40 studies found that almost 45% of people of various ages who committed suicide had contact with primary care providers within 1 month of the suicide [120]. The availability of effective clinical care for physical, mental, and substance abuse disorders is important for the prevention of suicide behaviors. Bruffaerts et al. [18] noted that the help seeking process of suicidal people is complex. People experiencing suicidal ideation often feel pessimistic and hopeless and they may not have positive expectations that treatment will be helpful for them [18]. Once someone has decided to seek treatment, they may wait and see whether the problem subsides by itself and whether they can “ride out the storm.” Adherence with recommended treatments is also important. A variety of strategies have been proposed to encourage people experiencing suicidal ideation or intent to seek help, to provide referrals, and to improve referral follow-through and attendance [18,121,122]. In the U.S., these strategies include a national network

of suicide prevention crisis lines, including the VA national crisis line (veteranscrisisline.net), the National Suicide Prevention Lifeline (<http://www.suicidepreventionlifeline.org>), and comparable resources for U.S. military personnel. A congressionally mandated comprehensive VA suicide prevention program began in 2007 [123]. National strategies for suicide prevention are periodically updated as new scientific information becomes available.

Acknowledgement

The authors are grateful to Dr. Lisa Brenner for helpful comments on this manuscript.

References

1. Simon RI (2008) Suicide. In: Hales RE, Yudofsky SC, Gabbard GO, editors. *The American Psychiatric Publishing Textbook of Psychiatry* (5th edn) USA.
2. Nock MK, Borges G, Bromet EJ, Cha CB, Kessler RC, et al. (2008) Suicide and suicidal behavior. *Epidemiol Rev* 30: 133-154.
3. Mann JJ (2002) A current perspective of suicide and attempted suicide. *Ann Intern Med* 136: 302-311.
4. Nock MK, Borges G, Bromet EJ, Alonso J, Angermeyer M, et al. (2008) Cross-national prevalence and risk factors for suicidal ideation, plans and attempts. *Br J Psychiatry* 192: 98-105.
5. Kessler RC, Berglund P, Borges G, Nock M, Wang PS (2005) Trends in suicide ideation, plans, gestures, and attempts in the United States, 1990-1992 to 2001-2003. *JAMA* 293: 2487-2495.
6. Mills PD, Huber SJ, Vince Watts B, Bagian JP (2011) Systemic vulnerabilities to suicide among veterans from the Iraq and Afghanistan Conflicts: review of case reports from a National Veterans Affairs Database. *Suicide Life Threat Behav* 41: 21-32.
7. Haney EM, O'Neil ME, Carson S, et al. (2012) Suicide Risk Factors and Risk Assessment Tools: A Systematic Review. VA-ESP Project 05-225.
8. Hawton K, Houston K, Haw C, Townsend E, Harriss L (2003) Comorbidity of axis I and axis II disorders in patients who attempted suicide. *Am J Psychiatry* 160: 1494-1500.
9. Cottler LB, Campbell W, Krishna VA, Cunningham-Williams RM, Abdallah AB (2005) Predictors of high rates of suicidal ideation among drug users. *J Nerv Ment Dis* 193: 431-437.
10. Maris RW (2002) Suicide. *Lancet* 360: 319-326.
11. Hawgood J, Spathonis K, De Leo D (2004) Physical illness and suicide behavior. In: De Leo D, Bille-Brahe U, Kerkhof A, Schmidtke A, editors. *Suicidal behavior: theories and research findings*. Gottingen, Germany: Hogrefe & Huber, 139-163.
12. Khang YH, Kim HR, Cho SJ (2010) Relationships of suicide ideation with cause-specific mortality in a longitudinal study of South Koreans. *Suicide Life Threat Behav* 40: 465-475.
13. Pfaff JJ, Almeida OP, Witte TK, Waesche MC, Joiner TE Jr (2007) Relationship between quantity and frequency of alcohol use and indices of suicidal behavior in an elderly Australian sample. *Suicide Life Threat Behav* 37: 616-626.
14. Fyer MR, Frances AJ, Sullivan T, Hurt SW, Clarkin J (1988) Suicide attempts in patients with borderline personality disorder. *Am J Psychiatry* 145: 737-739.
15. Glenn CR, Klonsky ED (2011) Prospective prediction of nonsuicidal self-injury: a 1-year longitudinal study in young adults. *Behav Ther* 42: 751-762.
16. Frances A, Fyer M, Clarkin J (1986) Personality and suicide. *Ann N Y Acad Sci* 487: 281-293.
17. Stone MH, Stone DK, Hurt SW (1987) Natural history of borderline patients treated by intensive hospitalization. *Psychiatr Clin North Am* 10: 185-206.
18. Bruffaerts R, Demyttenaere K, Hwang I, Chiu WT, Sampson N, et al. (2011) Treatment of suicidal people around the world. *Br J Psychiatry* 199: 64-70.
19. Brenner LA, Breshears RE, Betthausen LM, Bellon KK, Holman E, et al. (2011) Implementation of a suicide nomenclature within two VA healthcare settings. *J Clin Psychol Med Settings* 18: 116-128.
20. De Leo D, Cerin E, Spathonis K, Burgis S (2005) Lifetime risk of suicide ideation

- and attempts in an Australian community: prevalence, suicidal process, and help-seeking behavior. *J Affect Disord* 86:215-224.
21. Mathias CW, Furr RM, Sheftall AH, Nathalie Hill-Kapturczak, Paige Crum, et al. (2012) What's the harm in asking about suicidal ideation? *Suicide and Life-Threatening Behavior* 42: 341-351.
 22. Sher L, Oquendo MA, Mann JJ (2001) Risk of suicide in mood disorders. *Clin Neurosci Res* 1:337-344.
 23. Oquendo MA, Galfalvy H, Russo S, Ellis SP, Grunebaum MF, et al. (2004) Prospective study of clinical predictors of suicidal acts after a major depressive episode in patients with major depressive disorder or bipolar disorder. *Am J Psychiatry* 161: 1433-1441.
 24. Mann JJ, Apter A, Bertolote J, Beautrais A, Currier D, et al. (2005) Suicide prevention strategies: a systematic review. *JAMA* 294: 2064-2074.
 25. Beautrais A (2001) Suicides and serious suicide attempts: two populations or one? *Psychological Med* 31:837-845.
 26. Runeson B, Tidemalm D, Dahlin M, Lichtenstein P, Långström N (2010) Method of attempted suicide as predictor of subsequent successful suicide: national long term cohort study. *BMJ* 341: c3222.
 27. Mann JJ (1998) The neurobiology of suicide. *Nat Med* 4: 25-30.
 28. Rich CL, Fowler RC, Fogarty LA, Young D (1988) San Diego Suicide Study. III. Relationships between diagnoses and stressors. *Arch Gen Psychiatry* 45: 589-592.
 29. Isometsä E, Henriksson M, Marttunen M, Heikkinen M, Aro H, et al. (1995) Mental disorders in young and middle aged men who commit suicide. *BMJ* 310: 1366-1367.
 30. O'Carroll PW, Potter LB (1994) Suicide contagion and the reporting of suicide: recommendations from a national workshop. United States Department of Health and Human Services. *MMWR Recomm Rep* 43: 9-17.
 31. Large M, Sharma S, Cannon E, Ryan C, Nielssen O (2011) Risk factors for suicide within a year of discharge from psychiatric hospital: a systematic meta-analysis. *Aust N Z J Psychiatry* 45: 619-628.
 32. Kuo WH, Gallo JJ, Eaton WW (2004) Hopelessness, depression, substance disorder, and suicidality—a 13-year community-based study. *Soc Psychiatry Psychiatr Epidemiol* 39: 497-501.
 33. Reinherz HZ, Tanner JL, Berger SR, Beardslee WR, Fitzmaurice GM (2006) Adolescent suicidal ideation as predictive of psychopathology, suicidal behavior, and compromised functioning at age 30. *Am J Psychiatry* 163: 1226-1232.
 34. Bossarte RM, Knox KL, Piegari R, Altieri J, Kemp J, et al. (2012) Prevalence and characteristics of suicide ideation and attempts among active military and veteran participants in a national health survey. *Am J Public Health* 102: S38-40.
 35. Pietrzak RH, Goldstein MB, Malley JC, Rivers AJ, Johnson DC, et al. (2010) Risk and protective factors associated with suicidal ideation in veterans of Operations Enduring Freedom and Iraqi Freedom. *J Affect Disord* 123: 102-107.
 36. Thoresen S, Mehlum L (2008) Traumatic stress and suicidal ideation in Norwegian male peacekeepers. *J Nerv Ment Dis* 196: 814-821.
 37. Desai RA, Liu-Mares W, Dausey DJ, Rosenheck RA (2003) Suicidal ideation and suicide attempts in a sample of homeless people with mental illness. *J Nerv Ment Dis* 191: 365-371.
 38. Britton PC, Ilgen MA, Rudd MD, Conner KR (2012) Warning signs for suicide within a week of healthcare contact in Veteran decedents. *Psychiatry Res* 200: 395-399.
 39. Turner AP, Williams RM, Bowen JD, Kivlahan DR, Haselkorn JK (2006) Suicidal ideation in multiple sclerosis. *Arch Phys Med Rehabil* 87: 1073-1078.
 40. Kuo WH, Gallo JJ, Tien AY (2001) Incidence of suicide ideation and attempts in adults: the 13-year follow-up of a community sample in Baltimore, Maryland. *Psychol Med* 31: 1181-1191.
 41. Beck AT, Brown GK, Steer RA, Dahlsgaard KK, Grisham JR (1999) Suicide ideation at its worst point: a predictor of eventual suicide in psychiatric outpatients. *Suicide Life Threat Behav* 29: 1-9.
 42. Sher L, Braquehais MD, Casas M (2012) Posttraumatic stress disorder, depression, and suicide in veterans. *Cleve Clin J Med* 79: 92-97.
 43. Kessler RC, Borges G, Walters EE (1999) Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Arch Gen Psychiatry* 56: 617-626.
 44. Casey PR, Dunn G, Kelly BD, Birkbeck G, Dalgard OS, et al. (2006) Factors associated with suicidal ideation in the general population: five-centre analysis from the ODIN study. *Br J Psychiatry* 189: 410-415.
 45. Schoenfeld M, Myers RH, Cupples LA, Berkman B, Sax DS, et al. (1984) Increased rate of suicide among patients with Huntington's disease. *J Neurol Neurosurg Psychiatry* 47: 1283-1287.
 46. Breslau N, Davis GC, Schultz LR, Peterson EL (1994) Joint 1994 Wolff Award Presentation. Migraine and major depression: a longitudinal study. *Headache* 34: 387-393.
 47. Brent DA, Bridge J, Johnson BA, Connolly J (1996) Suicidal behavior runs in families. A controlled family study of adolescent suicide victims. *Arch Gen Psychiatry* 53: 1145-1152.
 48. Conner KR, Duberstein PR (2004) Predisposing and precipitating factors for suicide among alcoholics: empirical review and conceptual integration. *Alcohol Clin Exp Res* 28: 6S-17S.
 49. Sher L (2006) Alcohol consumption and suicide. *QJM* 99: 57-61.
 50. Sher L (2012) Protective factors. In: Shrivastava A, Kimbrell M, Lester D, editors. *Suicide from a global perspective: psychosocial approaches*. Hauppauge, New York: Nova Science Publishers, 57-65.
 51. Tarrier N, Taylor K, Gooding P (2008) Cognitive-behavioral interventions to reduce suicide behavior: a systematic review and meta-analysis. *Behav Modif* 32: 77-108.
 52. Arana A, Wentworth CE, Ayuso-Mateos JL, Arellano FM (2010) Suicide-related events in patients treated with antiepileptic drugs. *N Engl J Med* 363: 542-551.
 53. Nilsson L, Ahlbom A, Farahmand BY, Asberg M, Tomson T (2002) Risk factors for suicide in epilepsy: a case control study. *Epilepsia* 43: 644-651.
 54. Hecimovic H, Salpekar J, Kanner AM, Barry JJ (2011) Suicidality and epilepsy: a neuropsychobiological perspective. *Epilepsy Behav* 22: 77-84.
 55. Wen X, Meador KJ, Loring DW, Eisenschenk S, Segal R, et al. (2010) Is antiepileptic drug use related to depression and suicidal ideation among patients with epilepsy? *Epilepsy Behav* 19: 494-500.
 56. Machado RA, Espinosa AG, Melendrez D, González YR, García VF, et al. (2011) Suicidal risk and suicide attempts in people treated with antiepileptic drugs for epilepsy. *Seizure* 20: 280-284.
 57. Leon AC, Demirtas H, Li C, Hedeker D (2012) Two propensity score-based strategies for a three-decade observational study: investigating psychotropic medications and suicide risk. *Stat Med* 31: 3255-3260.
 58. Breshears RE, Brenner LA, Harwood JE, Gutierrez PM (2010) Predicting suicidal behavior in veterans with traumatic brain injury: the utility of the personality assessment inventory. *J Pers Assess* 92: 349-355.
 59. Reeves RR, Panguluri RL (2011) Neuropsychiatric complications of traumatic brain injury. *J Psychosoc Nurs Ment Health Serv* 49: 42-50.
 60. Coughlin SS, Sullivan PM, Chapman JC (2012) Post-traumatic stress disorder and traumatic brain injury. In: Coughlin SS (ed.) *Post-traumatic stress disorder and chronic health conditions*. Washington, DC: American Public Health Association.187-229.
 61. Simpson G, Tate R (2007) Suicidality in people surviving a traumatic brain injury: prevalence, risk factors and implications for clinical management. *Brain Inj* 21: 1335-1351.
 62. Miller E (2012) Multiple sclerosis. *Adv Exp Med Biol* 724: 222-238.
 63. Sadovnick AD, Ebers GC, Wilson RW, Paty DW (1992) Life expectancy in patients attending multiple sclerosis clinics. *Neurology* 42: 991-994.
 64. Wallin MT, Culpepper WJ, Coffman P, Pulaski S, Maloni H, et al. (2012) The Gulf War era multiple sclerosis cohort: age and incidence rates by race, sex and service. *Brain* 135: 1778-1785.
 65. Fromont A, Binquet C, Sauleau E, Fournel I, Despalins R, et al. (2012) National estimate of multiple sclerosis incidence in France (2001-2007) *Mult Scler* 18: 1108-1115.
 66. Alonso A, Jick SS, Olek MJ, Hernán MA (2007) Incidence of multiple sclerosis

- in the United Kingdom : findings from a population-based cohort. *J Neurol* 254: 1736-1741.
67. Deussing EC, Jankosky CJ, Clark LL, Otto JL (2012) Estimated incidence of multiple sclerosis among United States Armed Forces personnel using the Defense Medical Surveillance System. *Mil Med* 177: 594-600.
 68. Noonan CW, Williamson DM, Henry JP, Indian R, Lynch SG, et al. (2010) The prevalence of multiple sclerosis in 3 US communities. *Prev Chronic Dis* 7: A12.
 69. El Adssi H, Debouverie M, Guillemin F; LORSEP Group (2012) Estimating the prevalence and incidence of multiple sclerosis in the Lorraine region, France, by the capture-recapture method. *Mult Scler* 18: 1244-1250.
 70. Baranzini SE, Nickles D (2012) Genetics of multiple sclerosis: swimming in an ocean of data. *Curr Opin Neurol* 25: 239-245.
 71. Gourraud PA, Harbo HF, Hauser SL, Baranzini SE (2012) The genetics of multiple sclerosis: an up-to-date review. *Immunol Rev* 248: 87-103.
 72. Feinstein A (2011) Multiple sclerosis and depression. *Mult Scler* 17: 1276-1281.
 73. Prat A, Antel J (2005) Pathogenesis of multiple sclerosis. *Curr Opin Neurol* 18: 225-230.
 74. Hemmer B, Cepok S, Zhou D, Sommer N (2004) Multiple sclerosis -- a coordinated immune attack across the blood brain barrier. *Curr Neurovasc Res* 1: 141-150.
 75. Goldman Consensus Group (2005) The Goldman Consensus statement on depression in multiple sclerosis. *Mult Scler* 11: 328-337.
 76. Paparrigopoulos T, Ferentinos P, Kouzoupis A, Koutsis G, Papadimitriou GN (2010) The neuropsychiatry of multiple sclerosis: focus on disorders of mood, affect and behaviour. *Int Rev Psychiatry* 22: 14-21.
 77. Raskind MA (2008) Diagnosis and treatment of depression comorbid with neurologic disorders. *Am J Med* 121: S28-37.
 78. Goeb JL, Even C, Nicolas G, Gohier B, Dubas F, et al. (2006) Psychiatric side effects of interferon-beta in multiple sclerosis. *Eur Psychiatry* 21: 186-193.
 79. Patten SB, Francis G, Metz LM, Lopez-Bresnahan M, Chang P, et al. (2005) The relationship between depression and interferon beta-1a therapy in patients with multiple sclerosis. *Mult Scler* 11: 175-181.
 80. Feinstein A (2007) The clinical neuropsychiatry of multiple sclerosis. New York: Cambridge University Press(2ndedn) UK. 51-52
 81. Fredrikson S, Cheng Q, Jiang GX, Wasserman D (2003) Elevated suicide risk among patients with multiple sclerosis in Sweden. *Neuroepidemiology* 22: 146-152.
 82. Stenager EN, Stenager E, Koch-Henriksen N, Brønnum-Hansen H, Hyllested K, et al. (1992) Suicide and multiple sclerosis: an epidemiological investigation. *J Neurol Neurosurg Psychiatry* 55: 542-545.
 83. Brønnum-Hansen H, Stenager E, Nylev Stenager E, Koch-Henriksen N (2005) Suicide among Danes with multiple sclerosis. *J Neurol Neurosurg Psychiatry* 76: 1457-1459.
 84. Sumelahti ML, Hakama M, Elovaara I, Pukkala E (2010) Causes of death among patients with multiple sclerosis. *Mult Scler* 16: 1437-1442.
 85. Stenager EN, Jensen B, Stenager M, Stenager K, Stenager E (2011) Suicide attempts in multiple sclerosis. *Mult Scler* 17: 1265-1268.
 86. Feinstein A (2002) An examination of suicidal intent in patients with multiple sclerosis. *Neurology* 59: 674-678.
 87. Berman AL, Pompili M (2011) Suicide risk associated with medical conditions. Washington, DC: American Association of Suicidology,
 88. Castro-Borrero W, Graves D, Frohman TC, Flores AB, Hardeman P, et al. (2012) Current and emerging therapies in multiple sclerosis: a systematic review. *Ther Adv Neurol Disord* 5: 205-220.
 89. Bomprezzi R, Okuda DT, Alderazi YJ, Stüve O, Frohman EM (2012) From injection therapies to natalizumab: views on the treatment of multiple sclerosis. *Ther Adv Neurol Disord* 5: 97-104.
 90. Fang F, Valdimarsdóttir U, Fürst CJ, Hultman C, Fall K, et al. (2008) Suicide among patients with amyotrophic lateral sclerosis. *Brain* 131: 2729-2733.
 91. Venkova-Hristova K, Christov A, Kamaluddin Z, Kobalka P, Hensley K (2012) Progress in therapy development for amyotrophic lateral sclerosis. *Neurol Res Int* 2012: 187234.
 92. Garbuzova-Davis S, Rodrigues MC, Hernandez-Ontiveros DG, Louis MK, Willing AE, et al. (2011) Amyotrophic lateral sclerosis: a neurovascular disease. *Brain Res* 1398: 113-125.
 93. Turner MR, Agosta F, Bede P, Govind V, Lulé D, et al. (2012) Neuroimaging in amyotrophic lateral sclerosis. *Biomark Med* 6: 319-337.
 94. Cheah BC, Vucic S, Krishnan AV, Kiernan MC (2010) Riluzole, neuroprotection and amyotrophic lateral sclerosis. *Curr Med Chem* 17: 1942-1199.
 95. Bensimon G, Lacomblez L, Meininger V (1994) A controlled trial of riluzole in amyotrophic lateral sclerosis. ALS/Riluzole Study Group. *N Engl J Med* 330: 585-591.
 96. Miller RG, Mitchell JD, Moore DH (2012) Riluzole for amyotrophic lateral sclerosis (ALS)/motor neuron disease (MND) *Cochrane Database Syst Rev* 3: CD001447.
 97. Simmons Z (2005) Management strategies for patients with amyotrophic lateral sclerosis from diagnosis through death. *Neurologist* 11: 257-270.
 98. Morren JA, Galvez-Jimenez N (2012) Current and prospective disease-modifying therapies for amyotrophic lateral sclerosis. *Expert Opin Investig Drugs* 21: 297-320.
 99. Palmieri A, Sorarù G, Albertini E, Semenza C, Vottero-Ris F, et al. (2010) Psychopathological features and suicidal ideation in amyotrophic lateral sclerosis patients. *Neurol Sci* 31: 735-740.
 100. Floris G, Borghero G, Chiò A, Secchi L, Cannas A, et al. (2012) Cognitive screening in patients with amyotrophic lateral sclerosis in early stages. *Amyotroph Lateral Scler* 13: 95-101.
 101. Sher L (2007) Possible link between cognitive dysfunction and suicidal behavior in individuals with alcohol use disorders. *Int J Disabil Hum Dev* 6:149-52.
 102. Jollant F, Lawrence NL, Olié E, Guillaume S, Courtet P (2011) The suicidal mind and brain: a review of neuropsychological and neuroimaging studies. *World J Biol Psychiatry* 12: 319-339.
 103. Rabkin JG, Albert SM, Del Bene ML, O'Sullivan I, Tider T, et al. (2005) Prevalence of depressive disorders and change over time in late-stage ALS. *Neurology* 65: 62-67.
 104. Lulé D, Häcker S, Ludolph A, Birbaumer N, Kübler A (2008) Depression and quality of life in patients with amyotrophic lateral sclerosis. *Dtsch Arztebl Int* 105: 397-403.
 105. Averill AJ, Kasarskis EJ, Segerstrom SC (2007) Psychological health in patients with amyotrophic lateral sclerosis. *Amyotroph Lateral Scler* 8: 243-254.
 106. Pagnini F (2012) Psychological wellbeing and quality of life in amyotrophic lateral sclerosis: A review. *Int J Psychol* .
 107. Hogg KE, Goldstein LH, Leigh PN (1994) The psychological impact of motor neurone disease. *Psychol Med* 24: 625-632.
 108. Keilp JG, Gorlyn M, Russell M, Oquendo MA, Burke AK, et al. (2013) Neuropsychological function and suicidal behavior: attention control, memory and executive dysfunction in suicide attempt. *Psychol Med* 43: 539-551.
 109. Baldessarini RJ, Hennen J (2004) Genetics of suicide: an overview. *Harv Rev Psychiatry* 12: 1-13.
 110. Roy A, Segal NL (2001) Suicidal behavior in twins: a replication. *J Affect Disord* 66: 71-74.
 111. Fu Q, Heath AC, Bucholz KK, Nelson EC, Glowinski AL, et al. (2002) A twin study of genetic and environmental influences on suicidality in men. *Psychol Med* 32: 11-24.
 112. Smith AR, Ribeiro JD, Mikolajewski A, Taylor J, Joiner TE, et al. (2012) An examination of environmental and genetic contributions to the determinants of suicidal behavior among male twins. *Psychiatry Res* 197: 60-65.
 113. Voracek M, Loibl LM (2007) Genetics of suicide: a systematic review of twin studies. *Wien Klin Wochenschr* 119: 463-475.

114. Althoff RR, Hudziak JJ, Willemsen G, Hudziak V, Bartels M, et al. (2012) Genetic and environmental contributions to self-reported thoughts of self-harm and suicide. *Am J Med Genet B Neuropsychiatr Genet* 159B: 120-127.
115. Segal NL (2009) Suicidal behaviors in surviving monozygotic and dizygotic co-twins: is the nature of the co-twin's cause of death a factor? *Suicide Life Threat Behav* 39: 569-575.
116. Bednar MM, Perry A (2012) Neurorestoration therapeutics for neurodegenerative and psychiatric disease. *Neurol Res* 34: 129-142.
117. Van Langenhove T, van der Zee J, Van Broeckhoven C (2012) The molecular basis of the frontotemporal lobar degeneration-amyotrophic lateral sclerosis spectrum. *Ann Med* 44: 817-828.
118. Pandya RS, Mao LL, Zhou EW, Bowser R, Zhu Z, et al. (2012) Neuroprotection for amyotrophic lateral sclerosis: role of stem cells, growth factors, and gene therapy. *Cent Nerv Syst Agents Med Chem* 12: 15-27.
119. Van Orden KA, Witte TK, Cukrowicz KC, Braithwaite SR, Selby EA, et al. (2010) The interpersonal theory of suicide. *Psychol Rev* 117: 575-600.
120. Luoma JB, Martin CE, Pearson JL (2002) Contact with mental health and primary care providers before suicide: a review of the evidence. *Am J Psychiatry* 159: 909-916.
121. Luxton DD, June JD, Comtois KA (2013) Can postdischarge follow-up contacts prevent suicide and suicidal behavior? A review of the evidence. *Crisis* 34: 32-41.
122. Snarr JD, Heyman RE, Slep AM (2010) Recent suicidal ideation and suicide attempts in a large-scale survey of the U.S. Air Force: prevalences and demographic risk factors. *Suicide Life Threat Behav* 40: 544-552.
123. Blow FC, Bohnert AS, Ilgen MA, Ignacio R, McCarthy JF, et al. (2012) Suicide mortality among patients treated by the Veterans Health Administration from 2000 to 2007. *Am J Public Health* 102: S98-104.

This article was originally published in a special issue, **Treatment Resistant Mood Disorders** handled by Editor(s). Dr. Jan Fawcett, University of New Mexico School of Medicine, USA