

Takotsubo Cardiomyopathy Shortly Emerged after Falling through the Ice - A Case Report

Shestakova M, Kalashnikov V, Terekhin S, Melkozerov K, Pesegova E and Shvangiradze T*

Endocrinology Research Centre, Moscow, Russia

Abstract

This case is dedicated to a 71-year-old woman with stress induced cardiomyopathy emerged shortly after falling through the ice. She suffered from acute heart failure and fever. Electrocardiogram showed a small-scale ST elevation and echocardiography revealed typical dyskinesia of the apical segments, while coronary angiography demonstrated minimal changes in coronary arteries. The diagnosis of Takotsubo cardiomyopathy was suggested. The treatment of the patient included beta-blockers, angiotensin-converting enzyme inhibitors, aspirin, diuretics and antibiotics. In a while, the left ventricular function had fully recovered.

Keywords: Takotsubo cardiomyopathy; Ventricular dysfunction; Apical ballooning syndrome

Introduction

Takotsubo cardiomyopathy (TCM), also known as stress-induced cardiomyopathy, apical ballooning syndrome, and broken heart syndrome, is condition characterized by left ventricular apical ballooning, electrocardiographic changes with ST elevation or depression, formation of abnormal Q-wave, and elevation of cardiac enzymes mimics acute myocardial infarction, but without evidence of coronary artery disease [1,2].

Initially TCM was described by Satoh et al. [3]. In Japanese, there is a fisherman's octopus pot, the takotsubo, with rounded body and narrow neck [4]. The similar appearance we can see on echocardiogram, cardiac MRI, or ventriculography, at the end of systole the heart looks like a bulb with short neck and round bottom. The syndrome was renamed stress cardiomyopathy in 2006 [5].

There are a lot of theories but the etiology of the syndrome isn't clear yet. These theories include multivessel epicardial spasm, microvascular coronary dysfunction, catecholamine-induced injury and neurohumoral-related myocardial stunning [6].

The true incidence of stress cardiomyopathy is higher than estimated, as it remains unrecognized in many cases. TCM predominantly occurs in elderly postmenopausal women (over 90%) with the median age from 63 to 76 years of age. Mortality is only 1.1% [7,8].

Case Report

71-year-old female with history of hypertension, hyperlipidemia was skiing and suddenly fell through the ice. It took her about 10-20 min to get out from the cold water, and 1 hour to reach home. She was hospitalized in 4 hours after an accident with weakness, shortness of breath, tachycardia. The initial examination revealed a light drop of blood pressure (105/70 mm Hg), sinus tachycardia (130 beats/min), bubbling wheezes in medium and basal parts of the lungs, mostly on the right side.

A twelve-lead standard electrocardiogram (ECG) showed sinus tachycardia (129 beats/min) with small-scale ST-segment elevation in V_4 - V_6 leads. Gradually negative T waves appeared in I, aVL, V_4 - V_6 leads. Previous ECG was not presented.

Routine laboratory tests showed increased levels of white blood cells $18.2 \times 10^9/L$ (NR- $9 \times 10^9/L$), neutrophils 81.3% (NR-75%), troponin T level was 1.10 ng/ml (NR<0.03 ng/ml), SaO₂ 80%.

X-ray examination and the chest computer-aided tomography showed dimming of the all lung fields and pleural effusion about 150 ml. Echocardiography revealed left ventricular systolic dysfunction (hyperkinesis of basal segments and akinesis of apical segments of the left ventricular). Left ventricular ejection fraction (LVEF) was reduced to 42%. Obstructive coronary heart disease was rejected during the coronarography (Figure 1).

The rest myocardial perfusion scintigraphy by gated single photon emission computed tomography was performed. It revealed a characteristic shape of the left ventricle, perfusion defects in apical and average part of anterior wall (Figure 2).

The patient was treated symptomatically (with β -blockers, angiotensin-converting enzyme inhibitors, aspirin, antibiotics and diuretics). The condition of the patient improved steadily, weakness and shortness of breath disappeared. The treatment in hospital took 7 days.

At the follow-up visit, echocardiography showed significant improvement in the left ventricular systolic function with a normal ejection fraction (about 63%) and normal wall motion. In two months myocardial perfusion scintigraphy revealed no relevant perfusion defects (Figure 3).

Discussion

The described case of such a rapid progress of the symptoms is not solitary. In the literature, there are many situations with acute appearance of cardiomyopathy just after information about somebody's death or defeat of the favorite soccer team [3,9].

Takotsubo cardiomyopathy is a rare, but an increasingly recognized type of cardiomyopathy occurring after a recent emotional or physical

*Corresponding author: Teona Shvangiradze, Research Center of Endocrinology, Dmitriya Ulyanova street 11, 117036, Russia, Tel: +7(926)583-44-64; E-mail: teona_endo@mail.ru

Received November 02, 2013; Accepted December 05, 2013; Published December 08, 2013

Citation: Shestakova M, Kalashnikov V, Terekhin S, Melkozerov K, Pesegova E, et al. (2013) Takotsubo Cardiomyopathy Shortly Emerged after Falling through the Ice - A Case Report. J Clin Exp Cardiol 4: 277. doi:10.4172/2155-9880.1000277

Copyright: © 2013 Shestakova M, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1: Coronarography. (A) Left main coronary artery. (B) Right coronary artery.




Figure 2: The rest myocardial perfusion scintigraphy made on the second day after incident, revealing perfusion defects in apical and anterior part of anterior wall.




Figure 3: The rest myocardial perfusion scintigraphy made in two months after incident, without any significant perfusion defects.

stress or neurohumoral disorder. The prevalence of Takotsubo cardiomyopathy in the general population is estimated to be 1.2% to 2.2% [10]. Most patients are postmenopausal women.

Since the first description in Japanese studies, several case series

have been published. Despite this the etiology and pathophysiology is not fully understood. There are a lot of mechanisms underlying the development of this transient ventricular dysfunction.

Researchers have implicated gender-related differences in this syndrome. Akashi et al. have hypothesized that the reduced estrogen levels after menopause explain the predisposition of elderly women to this cardiomyopathy. Estrogen is thought to promote vasodilatation through endothelial nitric oxide synthase. Postmenopausal women lose this protective effect, which may predispose them to coronary spasm and myocardial stunning in the setting of elevated catecholamine levels. But this does not explain the occurrence of the disease in men [11].

So a central role in the pathogenesis of Takotsubo cardiomyopathy has been identified for the level of catecholamines. Kume et al. have described elevated levels of catecholamines as the cause of cardiomyopathy. There are a lot of studies confirming the increased level of catecholamine in Takotsubo cardiomyopathy in compared with myocardial infarction [12,13].

Under usual catecholamine release, epinephrine and norepinephrine contact to myocardial β_1 and β_2 adrenoreceptors and increase heart contractility and rate (the positive inotropic effect). It is worth noting, that there is an increased density of beta adrenoreceptors in the apex. During the emotional stress excess release of catecholamine initiates a cascade of intracellular events that leads to a decrease inotropy. In connection with the pathophysiological features of the location of the β -receptors we can see characteristic apical ballooning and hyperkinesis of basal segments [14].

Another way of transient dysfunction of the left ventricular is microvascular dysfunction or a multivessel acute epicardial coronary spasm that caused by increased level norepinephrine. It is interesting to suggest, that that cold water also contributed to the development of TCM. It is known that cooling causes reflexive vasoconstriction, due to increased affinity of alpha-adrenergic receptors for norepinephrine in the vascular walls [15].

Some studies indicate that ultrastructural damage of cardiomyocytes can be connected with direct action of high level catecholamine, that to cyclic adenosine mono-phosphate (AMP)-mediated calcium overload of the cell, with the resultant decrease in synthetic activity and viability, or oxygen-derived free radicals [10].

Among the evidence suggesting an important role of catecholamines in the pathophysiology of cardiomyopathy is the observation that cardiomyopathy is common in patients with pheochromocytoma [16].

To assist in the diagnosis of stress-induced cardiomyopathy, the proposed Mayo criteria suggested that all of the following 4 conditions be met [16,17]:

1. Transient akinesis or dyskinesis of the left ventricular apical and mid-ventricular segments with regional wall-motion abnormalities extending beyond a single epicardial vascular distribution.
2. Absence of obstructive coronary disease or angiographic evidence of acute plaque rupture.
3. New ECG abnormalities (either ST-segment elevation or T-wave inversion).
4. Absence of: pheochromocytoma, myocarditis, intracranial bleeding.

There are no specific treatments for Takotsubo cardiomyopathy, but supportive measures are needed in many cases.

The prognosis is usually good. The average time to recovery is about 2-3 weeks. The risk of recurrence ranges from 2-10%, and the long-term prognosis is usually excellent.

Conclusion

There is the description of many reasons connected with the development of Takotsubo cardiomyopathy including emotional stress, pheochromocytoma, physical distress, surgery, but catecholamines are believed to play a central role. Such a rapid increase of symptoms, similar with an acute coronary syndrome (chest pain, dyspnea, ischemic changes on electrocardiogram (ST changes, or T-wave inversion) and cardiac enzyme elevation, which leads to the need of supplementary methods (coronary angiography) to establish the diagnosis. It may sparing patients the risks of unnecessary thrombolytic therapy.

Takotsubo cardiomyopathy should be considered in all patients presenting with acute onset chest pain and elevated cardiac biomarkers.

References

1. Murase K, Takagi K (2012) Takotsubo cardiomyopathy in a snake bite victim: a case report. *Pan Afr Med J* 13: 51.
2. Doesch C, Burgstahler C, Seeger A, Miller S, May AE (2009) Chest pain and reversible midventricular ballooning in a woman after witnessing sudden cardiac death: a possible variant of takotsubo cardiomyopathy. *Can J Cardiol* 25: e22.
3. Fijalkowski M, Fijalkowska M, Nowak R, Rynkiewicz A (2013) Takotsubo cardiomyopathy in a male during a Euro 2012 football match. *Clin Res Cardiol* 102: 319-321.
4. Velankar P, Buegler J (2012) A mid-ventricular variant of Takotsubo cardiomyopathy. *Methodist Debaquey Cardiovasc J* 8: 37-39.
5. Maron BJ, Towbin JA, Thiene G, Antzelevitch C, Corrado D, et al. (2006) Contemporary definitions and classification of the cardiomyopathies: an American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary Working Groups; and Council on Epidemiology and Prevention. *Circulation* 113: 1807-1816.
6. Kepez A, Yesildag O, Erdogan O, Aktas B (2012) Takotsubo cardiomyopathy in a patient with lung adenocarcinoma. *Heart Views* 13: 107-110.
7. Jenab Y, Taher M, Shirzad S (2012) Broken heart syndrome: a case report. *J Tehran Heart Cent* 7: 136-139.
8. Yamashita K, Ishii H, Hirota K, Sato M, Tanabe H, et al. (2012) Takotsubo cardiomyopathy during ambulatory anesthesia for bladder hydrodistension therapy -A case report-. *Korean J Anesthesiol* 62: 484-487.
9. Potter B, Gobeil F, Oiknine A, Laramée P (2010) The first case of takotsubo cardiomyopathy associated with sodium tetradecyl sulphate sclerotherapy. *Can J Cardiol* 26: 146-148.
10. Manzanal A, Ruiz L, Madrazo J, Makan M, Perez J (2013) Inverted Takotsubo cardiomyopathy and the fundamental diagnostic role of echocardiography. *Tex Heart Inst J* 40: 56-59.
11. Bielecka-Dabrowa A, Mikhailidis DP, Hannam S, Rysz J, Michalska M, et al. (2010) Takotsubo cardiomyopathy--the current state of knowledge. *Int J Cardiol* 142: 120-125.
12. Bounhoure JP (2012) Takotsubo or stress cardiomyopathy. *Cardiovasc Psychiatry Neurol* 2012: 637672.
13. Turley A, Graham R, Hall J (2008) Takotsubo cardiomyopathy in two female patients: two case reports. *Cases J* 1: 325.
14. Singh A, Southwick F, Gumm E, Makar JS (2012) Exploring takotsubo cardiomyopathy in an elderly patient with acute anxiety attack. *W V Med J* 108: 14-17.
15. White GE, Wells GD (2013) Cold-water immersion and other forms of cryotherapy: physiological changes potentially affecting recovery from high-intensity exercise. *Extrem Physiol Med* 2: 26.
16. Redfors B, Shao Y, Omerovic E (2013) Stress-induced cardiomyopathy (Takotsubo)--broken heart and mind? *Vasc Health Risk Manag* 9: 149-154.
17. Daly MJ, Dixon LJ (2009) Tako-tsubo cardiomyopathy presenting with acute pulmonary edema. *Congest Heart Fail* 15: 46-48.

Citation: Shestakova M, Kalashnikov V, Terekhin S, Melkozerov K, Pesegova E, et al. (2013) Takotsubo Cardiomyopathy Shortly Emerged after Falling through the Ice - A Case Report. *J Clin Exp Cardiol* 4: 277. doi:[10.4172/2155-9880.1000277](https://doi.org/10.4172/2155-9880.1000277)