

# The Biological Screening of Extracts/Fractions of Various Parts of *Pistacia integerrima* Stewart against Pathogenic Fungi

Ghias Uddin<sup>1\*</sup>, Abdur Rauf<sup>1</sup>, Bina Shaheen Siddiqui<sup>2</sup> and Haroon Khan<sup>3</sup>

<sup>1</sup>Institute of Chemical Sciences, University of Peshawar, Pakistan

<sup>2</sup>H.E.J. Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Pakistan

<sup>3</sup>Department of Pharmacy, Abdul Wali Khan University Mardan, Pakistan

## Abstract

The goal of the current study was to scrutinize the antifungal activity of the crude extracts and subsequent solvent fractions of different parts of *Pistacia integerrima* against selected pathogenic fungi using agar tube dilution method. The results illustrated susceptibility of extracts/fractions of different parts of the plant against *Microsporum canis*, *Aspergillus flavus* and *Fusarium solani*. However, the rest of fungi were not sensitive to extracts/fractions. In conclusions, our study provided a strong evidence for the isolation of secondary metabolites from different parts of the plant as antifungal.

**Keywords:** *Pistacia integerrima*; Extracts/fractions of different parts; Antifungal activity

## Introduction

*Pistacia integerrima* (J. L. Stewart ex Brandis) belongs to family anacardiaceae. It mostly grows at a height of 12000 to 8000 feet in Eastern Himalayan regions [1]. It is a variable sized tree that can grow up to forty feet. *P. integerrima* has been used in the treatment of inflammation, diabetes, blood purification, gastrointestinal problems, and as an expectorant. Indian are using this plant as antiasthmatic, antipyretic, antiemetic and antidiarrheal [2,3]. In Pakistan, galls of *P. integerrima* are used for the treatment of hepatitis and other liver disorders [4,5], infections, diabetes, pain, inflammatory conditions, and fever [6].

Phytochemically, monoterpenes, triterpenoids sterols, dihydromalvalic acid and flavonoids have been isolated from the different parts of *Pistacia* species [7,8]. In this research article, we present the experimental findings of various fractions of different parts of *P. integerrima* against selected pathogenic fungi.

## Material and Methods

### Collection of plant materials

*P. integerrima* were collected from Toormang, Razagram (District Dir), Khyber Pakhtunkhwa, Pakistan in the month of February, 2010. The identification of plant material was done by Prof. Dr. Abdur Rashid plant taxonomist Department of Botany, University of Peshawar. A voucher specimen (Bot.20037(PUP)) was deposited in the herbarium of the same institution.

### Extraction and fractionation

Shade dried and crushed galls, leaves, bark and roots of *P. integerrima* (Stewart) were extraction with ethanol. The resulting methanolic extract of each part was suspended in water and sequentially fractionated with n-hexane, chloroform and ethyl acetate in order to get respective major fraction.

### Fungal strains

Fungal strains include *Trichyton logifusus* (clinical isolate), *Candida albicans* ATCC 2091, *Candida glabrata* ATCC 90030 *Aspergillus flavus* ATCC 32611, *Microsporum canis* ATCC 11622, and *Fusarium solani* 11712. They were maintained on agar slant at 4°C. Before any test, the activation of strains was occurred at 37°C for 24 h on Sabouraud Glucose Agar (SGA).

## Antifungal assay

Agar tube dilution method was employed for the assessment of antifungal activity of extracts and fractions of various parts *P. integerrima* [9]. The extracts as well as fractions (24 mg/ ml) were dissolved in the sterile Dimethyl sulfoxide (DMSO) for the preparation of stock solution. Sabouraud Dextrose Agar (SDA) 4 ml was dispensed into screw cap tubes, which were autoclaved at 120°C for 15 min and then cooled to 15°C. The non-solidified SDA media was poisoned with stock solution (66.6 µl) giving the final concentration of 400 µg of the extract per ml of SDA. Tubes were allowed to solidify in the slanted position at room temperature. Each tube was inoculated with a piece (4 mm diameter) of inoculum removed from a seven days old culture of fungi for non-mycelial growth; an agar surface streak was employed. DMSO was used as control while miconazol and amphotericin-B as standard drugs. Inhibition of fungal growth was observed after 7 days of incubation at 28 ± 1°C and relative humidity (40-50%). The test tubes were analyzed for the visible growth of the microorganisms and % antifungal activity was determined.

## Statistical analysis

The experimental findings are mean ± SEM of three different readings. Analysis was performed by using GraphPad Prism version 6.

## Results

### Antifungal effect of the galls *P. integerrima*

The antifungal activity of the methanolic extract and subsequent solvent fractions of the galls of *P. integerrima* are presented in Table 1. Of the test fungi, *M. canis* was the only susceptible fungus to various extracts and fractions of the galls of the plant. However, the rest of fungi were not receptive to extract and fractions.

**\*Corresponding author:** Ghias Uddin, Institute of Chemical Sciences, University of Peshawar, K.P.K Peshawar-25120, Pakistan, Tel: +92-345 325 6686; E-mail: [drghiasuddin@hotmail.com](mailto:drghiasuddin@hotmail.com), [Ghiasuddin@upesh.edu.pk](mailto:Ghiasuddin@upesh.edu.pk)

**Received** November 01, 2013; **Accepted** December 10, 2013; **Published** December 13, 2013

**Citation:** Uddin G, Rauf A, Siddiqui BS, Khan H (2013) The Biological Screening of Extracts/Fractions of Various Parts of *Pistacia integerrima* Stewart against Pathogenic Fungi. Transl Med 4: 120. doi:10.4172/2161-1025.1000120

**Copyright:** © 2013 Uddin G et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

### Antifungal effect of the leaves of *P. integerrima*

The results of antifungal methanolic extract and subsequent solvent fractions of the leaves of *P. integerrima* are presented in Table 2. *M. canis*, *A. flavus* and *F. solani* evoked sensitive to extract/fractions among the test fungi.

### Antifungal effect of the bark of *P. integerrima*

The effects of test extract/fractions of the bark of *P. integerrima* are presented in Table 3. The results showed susceptibility of extract/fractions only against *M. canis* and *A. flavus*.

### Antifungal effect of the roots of *P. integerrima*

The antifungal effects of the roots of *P. integerrima* against selected pathogenic fungi are illustrated in Table 4. The extract/fractions exhibited activity only against *M. canis* and *A. flavus*.

### Discussion

The current article was designed to evaluate the antifungal activity of extracts/fractions of various parts (galls, leaves, bark and roots) of *P. integerrima* against various pathogenic fungi.

In the course of time, dramatic changes have been observed in human mycoses. Along with a significant increased in the number of infected patients, there have been noted changes in the fungal pathogens. The best possible explanation for these alterations is both the emergence of new fungal pathogens and the expanding spectrum with more serious infections due to classic fungal pathogens. Probably, these effects are paralleled and directly related to increases in patient populations that are highly susceptible to fungal infections. These effects mostly including patients with surgery, cancers, bone marrow and solid organ transplant recipients, and other immunosuppressed patients, most notably those infected with the Human-Immune-

Name of Fungus	% Zone of inhibition (mm)					
	n-Hexane	Chloroform	Ethyl acetate	Methanol	STD	MIC (µg/ml)
<i>Trichophyton longifusus</i>	-	-	-	-	Miconazole	
<i>Candida albicans</i>	-	-	-	-	Miconazole	110.8
<i>Aspergillus flavus</i>	-	20 ± 0.50	-	-	Amphotericin B	20.20
<i>Microsporium canis</i>	20 ± 0.42	35 ± 0.46	25 ± 0.57	15 ± 0.21	Miconazole	98.4
<i>Fusarium solani</i>	-	-	-	-	Miconazole	73.25
<i>Candida glabrata</i>	-	-	-	-	Miconazole	110.8

Results represent the mean of three different experimental readings

**Table 1:** Antifungal (%) extract and fractions of the galls of *P. integerrima* against test fungi

Name of Fungus	% Zone of inhibition (mm)					
	n-Hexane	Chloroform	Ethyl acetate	Methanol	STD	MIC (µg/ml)
<i>Trichophyton longifusus</i>	-	-	-	-	Miconazole	
<i>Candida albicans</i>	-	-	-	-	Miconazole	110.8
<i>Aspergillus flavus</i>	20 ± 0.22	25 ± 0.53	-	20 ± 0.23	Amphotericin B	20.20
<i>Microsporium canis</i>	-	30 ± 0.68	30 ± 0.63	20 ± 0.12	Miconazole	98.4
<i>Fusarium solani</i>	20 ± 0.34	-	20 ± 0.37	-	Miconazole	73.25
<i>Candida glabrata</i>	-	-	-	-	Miconazole	110.8

Results represent the mean of three different experimental readings

**Table 2:** Antifungal (%) extract and fractions of the leaves of *P. integerrima* against test fungi

Name of Fungus	% Zone of inhibition (mm)					
	n-Hexane	Chloroform	Ethyl acetate	Methanol	STD	MIC (µg/ml)
<i>Trichophyton longifusus</i>	-	-	-	-	Miconazole	
<i>Candida albicans</i>	-	-	-	-	Miconazole	110.8
<i>Aspergillus flavus</i>	-	20 ± 0.02	-	-	Amphotericin B	20.20
<i>Microsporium canis</i>	20 ± 0.21	35 ± 0.49	25 ± 0.72	15 ± 0.03	Miconazole	98.4
<i>Fusarium solani</i>	-	-	-	-	Miconazole	73.25
<i>Candida glabrata</i>	-	-	-	-	Miconazole	110.8

Results represent the mean of three different experimental readings

**Table 3:** Antifungal (%) extract and fractions of the barks of *P. integerrima* against test fungi

Name of Fungus	% Zone of inhibition (mm)					
	n-Hexane	Chloroform	Ethyl acetate	Methanol	STD	MIC (µg/ml)
<i>Trichophyton longifusus</i>	-	-	-	-	Miconazole	
<i>Candida albicans</i>	-	-	-	-	Miconazole	110.8
<i>Aspergillus flavus</i>	-	20 ± 0.32	-	-	Amphotericin B	20.20
<i>Microsporium canis</i>	20 ± 0.08	30 ± 0.63	25 ± 0.47	20 ± 0.19	Miconazole	98.4
<i>Fusarium solani</i>	-	-	-	-	Miconazole	73.25
<i>Candida glabrata</i>	-	-	-	-	Miconazole	110.8

Results represent the mean ± SEM of three different experimental readings

**Table 4:** Antifungal (%) extract and fractions of the roots of *P. integerrima* against test fungi

Deficiency Virus (HIV) [10].

*M. canis* is involved in the pathogenesis of tinea capitis in humans, and simple ringworm in animals (pets). The organism's major reservoir in companion animals is within domestic cats and dogs. Under Wood's lamp examination, it gives bright green fluorescence. It is closely related to other dermatophytes [11]. The extracts/fractions of different parts of the plant showed antifungal activity against *M. canis*. It is therefore, suggested that the extracts/fractions of different parts of the plant possessed active constituents that were sensitive to *M. canis*.

*A. flavus* is a common filamentous fungus that produces aflatoxins and presents a major threat to agriculture and human health [12]. The extracts of different parts of the test plant illustrated some activity against *A. flavus*. The results suggested the presence of pharmacologically active secondary metabolites that exhibited antifungal activity.

Members of the *F. solani* species are increasingly implicated as the causative agents of human mycoses, particularly in the expanding immunocompromised and immunosuppressed patient populations [13]. Based on our finding, extracts possessed active molecules that had some antifungal potential.

In short, the present study demonstrated wide antifungal sensitive of extracts/fractions of *P. integerrima* against different pathogenic fungi. Isolation of secondary metabolites from different parts of the plant will explain the chemical background of current study.

## References

1. Uddin G, Rauf A, Rehman TU, Qaisar M (2011) Phytochemical Screening of *Pistacia chinensis* var. *integerrima*. *Middle-East J Sci Res* 7(5): 707-711.
2. Pant S, Samant SS (2010) Ethanobotanical observation in the Momaula Reserve Forest of Koumoun., West Himalaya. *India Ethanobotanical Leaflets* 1493.
3. Uddin G, Rauf A, Al-Othman AM, Collina S, Arfan M, et al. (2012) Pistagremic acid, a glucosidase inhibitor from *Pistacia integerrima*. *Fitoterapia* 83: 1648-1652.
4. Uddin G, Rauf A, Arfan M, Waliullah, Khan I, et al. (2012) Pistagremic acid a new leishmanicidal triterpene isolated from *Pistacia integerrima* Stewart. *J Enzyme Inhib Med Chem* 27: 646-648.
5. Uddin G, Rauf A, Siddiqui BS, Khan H (2013) Cytotoxic Activity of Extracts/Fractions of Various Parts of *Pistacia integerrima* Stewart. *Transl Med* 3: 2161-1025.
6. Ahmad NS, Farman M, Najmi MH, Mian KB, Hasan A (2008) Pharmacological basis for use of *Pistacia integerrima* leaves in hyperuricemia and gout. *J Ethnopharmacol* 117: 478-482.
7. Ahmad NS, Waheed A, Farman M, Qayyum A (2010) Analgesic and anti-inflammatory effects of *Pistacia integerrima* extracts in mice. *J Ethnopharmacol* 129: 250-253.
8. Arfan M, Rauf A, Tahir MN, Ali M, Uddin G (2011) 2-Methyl-6-(4,4,10,13,14-penta-methyl-3-oxo-2,3,4,5,6,7,10,11,12,13,14,15,16,17-tetra-deca-hydro-1H-cyclo-penta-[a]phenanthren-17-yl)hept-2-enoic acid. *Acta Crystallogr Sect E Struct Rep Online* 67: o711.
9. Khan H, Khan MA, Mahmood T, Choudhary MI (2008) Antimicrobial activities of *Gloriosa superba* Linn (Colchicaceae) extracts. *J Enz Inhib Med Chem* 23: 855-859.
10. King D, Cheever LW, Hood A, Horn TD, Rinaldi MG, et al. (1996) Primary invasive cutaneous *Microsporium canis* infections in immunocompromised patients. *J Clin Microbiol* 34: 460-462.
11. Ginter-Hanselmayer G, Smolle J, Gupta A (2004) Itraconazole in the treatment of tinea capitis caused by *Microsporium canis*: experience in a large cohort. *Pediatr Dermatol* 21: 499-502.
12. Geiser DM, Dörner JW, Horn BW, Taylor JW (2000) The phylogenetics of mycotoxin and sclerotium production in *Aspergillus flavus* and *Aspergillus oryzae*. *Fungal Genet Biol* 31: 169-179.
13. Zhang N, O'Donnell K, Sutton DA, Nalim FA, Summerbell RC, et al. (2006) Members of the *Fusarium solani* species complex that cause infections in both humans and plants are common in the environment. *J Clin Microbiol* 44: 2186-2190.