

The Case for Early Psychotherapy in Aging Combat Veterans Experiencing Late Onset Stress Symptomatology

Rajdip Barman¹ and Mark B. Detweiler²⁻⁵

¹Virginia Tech-Carilion Psychiatry Residency Program, Roanoke, Virginia

²Staff Psychiatrist, Salem Veterans Affairs Medical Center, Salem, Virginia

³Geriatric Research Group, Salem Veterans Affairs Medical Center, Salem, Virginia

⁴Virginia Tech-Carilion School of Medicine, Department of Psychiatry and Behavioral Medicine, Roanoke, Virginia

⁵Edward via College of Osteopathic Medicine Department of Psychiatry, Blacksburg, Virginia

Delayed onset post-traumatic stress disorder (DPTSD) is defined as onset of PTSD symptoms 6 months or more after a trauma. There is a wide range of variation in prevalence of DPTSD in combat veterans, ranging from 8.3-34% [1-3], but many studies cite a narrower prevalence of 20-25% [4-6]. Symptoms of combat related DPTSD may develop even after a prolonged silent period of 20-40 years [7]. The presence of life stressors such as the deterioration of physical health, death of family members and friends, retirement and lack of social support can precipitate trauma related symptoms in later life. Brain aging may also be a variable as one case report found an association with cognitive deficits with delayed onset of trauma and stress related symptoms [8].

DPTSD may be a part of the waxing and waning nature of PTSD [9]. It may present clinically following a symptom free period [10,11] as an exacerbation of sub threshold PTSD symptoms, or following many of the stressful events accompanying aging such as loss of employment. Loss of partner or loss of independence may also precipitate DPTSD [2,5,10]. Individual coping and resilience play important roles in development of these symptoms [12-14].

While PTSD is recognized as a clinical disorder, stress related symptoms may emerge in a non-pathological form called late onset stress symptomatology (LOSS). It is considered to be part of the spectrum of the normal ageing process. However, if not resolved, LOSS may progress to sub threshold or full blown PTSD with more severe clinical symptoms and life disruption [7,15,16]. Davison et al. [7] characterized the LOSS phenomenon in ageing combat veterans who were: a) exposed to highly stressful combat events in their early adulthood; b) asymptomatic for years after trauma exposure; c) functionally successful over the course of their lives; and d) begin to register increased combat related thoughts, feelings and reminiscences as they encounter the changes and challenges of ageing. LOSS symptoms may be precipitated by significant life stressors, and/or emerging cognitive deficits [8]. Rather than having the core symptoms of PTSD such as flashbacks, avoidance and negative emotions, LOSS symptoms comprise repeated thoughts about combat related trauma, irritability and/or nightmares that do not impair daily functioning [15,16].

There are biological factors associated with trauma and stress related disorders such as hippocampus damage following prolonged exposure to stress with decreased volume [17-19], altered hypothalamic-pituitary-adrenal axis function, increased pro-inflammatory cytokines [20-22] and decreased N-Acetyl Aspartate [23]. In one study of middle aged and elderly veterans, there was no difference in hippocampal volume between veterans with and without PTSD. However, veterans that developed PTSD following their first traumatic event had smaller left hippocampal volumes compared to veterans developed PTSD after exposure to multiple traumatic events. Also, veterans with PTSD had significantly lower urinary cortisol levels and diminished memory performance when compared to veterans without PTSD [24].

As Vietnam era combat veterans are entering, or have entered the years when LOSS is most commonly encountered, it is important to understand the differences among LOSS, subthreshold PTSD and full PTSD, as treatment and psychotherapy are not the same for each presentation [8]. LOSS patients do not need psychiatric medications, but may benefit from psych educational or life review therapeutic approaches [25,26]. The life review approach, which is also called reminiscence therapy, is a form of vocal or silent recall of life events either in individual or group settings. During therapy veterans attempt to accept the negative events, resolve past conflicts, recollect positive memories and reconcile the discrepancy between the ideal and reality. Participants are assisted in finding meaning and worth in life as it was lived, which is critical for successful ageing [26]. Cappeliez et al. [27] suggests placing these functions within a narrative that provides an uplifting life frame work. A number of studies show significant benefit in using 'props' such as photographs, videos, scents, music in older adults with depression and cognitive impairment [28-31]. With these psychotherapeutic approaches, ageing combat veterans are able to recall positive events in their premilitary, military and post military life that were the foundation of their positive quality of life prior to the onset of LOSS. Hence, rekindling their faltering or lost positive self-identity prior to the stressors accompanying aging, in addition to building problem solving skills using CBT, may reduce their negative perceptions and undergird the positive cognitive processes that had enabled them in the past to have a healthy life with normal ageing for decades following their combat tours.


If LOSS can be conceptualized as a normal aging event, which is a reversible phase of combat stress, early therapeutic intervention using established non pharmacologic augmented psychotherapeutic methods, may result in significant clinical benefits. As LOSS is considered a healthy, but transitional, state along the continuum of combat stress sequelae, early intervention is recommended to prevent the progression of LOSS to subthreshold PTSD and full PTSD. Such strategies should improve the veterans' quality of life in their elder years and reduce health care system burden (Figure 1).

*Corresponding author: Mark B. Detweiler, Mental Health Service Line (116A7), Veterans Affairs Medical Center, 1970 Roanoke Boulevard, Salem, Virginia, 24153, USA, Tel: 540-982-2463; Fax: 540-983-1080; E-mail: mark.detweiler1@va.gov

Received June 06, 2015; Accepted August 13, 2015; Published August 20, 2015

Citation: Barman R, Detweiler MB (2015) The Case for Early Psychotherapy in Aging Combat Veterans Experiencing Late Onset Stress Symptomatology. J Psychol Psychother 5: 200. doi: [10.4172/2161-0487.1000200](https://doi.org/10.4172/2161-0487.1000200)

Copyright: © 2015 Barman R, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



Conflict of Interest

The authors report no conflict of interest. The report findings do not represent the views of the Department of Veterans Affairs or the United States Government.

Acknowledgment

This material is the result of work supported with resources and the use of facilities at the Salem, Virginia Veterans Affairs Medical Center.

References

- Frueh BC, Grubaugh AL, Yeager DE, Magruder KM (2009) Delayed-onset post-traumatic stress disorder among war veterans in primary care clinics. *Br J Psychiatry* 194: 515-520.
- Op den Velde W, Falger PR, Hovens JE, De Groen JH, Lasschuit LJ, et al. (1993) Posttraumatic stress disorder in Dutch resistance veterans from World War II. In: Wilson JP, Raphael B (Edr), *International Handbook of Traumatic Stress Syndromes*. Plenum Press, New York.
- Solomon Z, Horesch D, Ein-Dor T, Ohry A (2012) Predictors of PTSD trajectories following captivity: a 35-year longitudinal study. *Psychiatry Res* 199: 188-194.
- Prigerson HG, Maciejewski PK, Rosenheck RA (2001) Combat trauma: trauma with highest risk of delayed onset and unresolved posttraumatic stress disorder symptoms, unemployment and abuse among men. *Journal of Nervous and Mental Disease* 189: 99-108.
- Horesch D, Solomon Z, Keinan G, Ein-Dor T (2013) The clinical picture of late-onset PTSD: a 20-year longitudinal study of Israeli war veterans. *Psychiatry Res* 208: 265-273.
- Smid GE, Mooren TT, van der Mast RC, Gersons BP, Kleber RJ (2009) Delayed posttraumatic stress disorder: systematic review, meta-analysis, and meta-regression analysis of prospective studies. *J Clin Psychiatry* 70: 1572-1582.
- King LA, King DW, Vickers K, Davison EH, Spiro A 3rd (2007) Assessing late-onset stress symptomatology among aging male combat veterans. *Aging Ment Health* 11: 175-191.
- Barman R, Detweiler MB (2014) Late onset stress symptomatology, subclinical PTSD or mixed etiologies in previously symptom free aging combat veterans. *Journal of Traumatic Stress disorders & Treatment. J Trauma Stress Disor Treat* 3: 4.
- Gray MJ, Bolton EE, Litz BT (2004) A longitudinal analysis of PTSD symptom course: delayed-onset PTSD in Somalia peacekeepers. *J Consult Clin Psychol* 72: 909-913.
- Herrmann N, Eryavec G (1994) Delayed onset post-traumatic stress disorder in World War II veterans. *Can J Psychiatry* 39: 439-441.
- Andrews B, Brewin CR, Philpott R, Stewart L (2007) Delayed-onset posttraumatic stress disorder: a systematic review of the evidence. *Am J Psychiatry* 164: 1319-1326.
- Aldwin CM, Levenson MR, Spiro A 3rd (1994) Vulnerability and resilience to combat exposure: can stress have lifelong effects? *Psychol Aging* 9: 34-44.
- Pietrzak RH, Southwick SM (2011) Psychological resilience in OEF-OIF Veterans: application of a novel classification approach and examination of demographic and psychosocial correlates. *J Affect Disord* 133: 560-568.
- Zerach G, Solomon Z, Cohen A, Ein-Dor T (2013) PTSD, resilience and posttraumatic growth among ex-prisoners of war and combat veterans. *Isr J Psychiatry Relat Sci* 50: 91-99.
- Potter CM, Kaiser AP, King LA, King DW, Davison EH, et al. (2013) Distinguishing late-onset stress symptomatology from posttraumatic stress disorder in older combat veterans. *Aging Ment Health* 17: 173-179.
- King LA, King DW, Vickers K, Davison EH, Spiro A 3rd (2007) Assessing late-onset stress symptomatology among aging male combat veterans. *Aging Ment Health* 11: 175-191.
- Duan X, Kang E, Liu CY, Ming GL, Song H (2008) Development of neural stem cell in the adult brain. *Curr Opin Neurobiol* 18: 108-115.
- McKittrick CR, Magariños AM, Blanchard DC, Blanchard RJ, McEwen BS, et al. (2000) Chronic social stress reduces dendritic arbors in CA3 of hippocampus and decreases binding to serotonin transporter sites. *Synapse* 36: 85-94.
- Pham K, Nacher J, Hof PR, McEwen BS (2003) Repeated restraint stress suppresses neurogenesis and induces biphasic PSA-NCAM expression in the adult rat dentate gyrus. *Eur J Neurosci* 17: 879-886.
- McEwen BS, Sapolsky RM (1995) Stress and cognitive function. *Curr Opin Neurobiol* 5: 205-216.
- Sapolsky RM (1996) Why stress is bad for your brain. *Science* 273: 749-750.
- Gill JM, Saligan L, Woods S, Page G (2009) PTSD is associated with an excess of inflammatory immune activities. *Perspect Psychiatr Care* 45: 262-277.
- Schuff N, Neylan TC, Lenoci MA, Du AT, Weiss DS, et al. (2001) Decreased hippocampal N-acetylaspartate in the absence of atrophy in posttraumatic stress disorder. *Biol Psychiatry* 50: 952-959.
- Yehuda R, Schmeidler J, Labinsky E, Bell A, Morris A, et al. (2009) Ten-year follow-up study of PTSD diagnosis, symptom severity and psychosocial indices in aging holocaust survivors. *Acta Psychiatr Scand* 119: 25-34.
- Wu LF (2011) Group integrative reminiscence therapy on self-esteem, life satisfaction and depressive symptoms in institutionalised older veterans. *J Clin Nurs* 20: 2195-2203.
- Bohlmeijer E, Roemer M, Cuijpers P, Smit F (2007) The effects of reminiscence on psychological well-being in older adults: a meta-analysis. *Aging Ment Health* 11: 291-300.
- Cappeliez P, Guindon M, Robitaille A (2008) Functions of reminiscence and emotional regulation among older adults. *Journal of Aging Studies* 22: 266-272.
- Housden S (2009) The use of reminiscence in the prevention and treatment of depression in older people living in care homes: A literature review. *Group work* 19: 28-45.
- Zauszniewski JA, Eggenschwiler K, Reechawong S, Chung CW, Airey TF, et al. (2004) Focused reflection reminiscence group for elders: Implementation and evaluation. *Journal of Applied Gerontology* 23: 429-442.

30. Chao SY, Liu HY, Wu CY, Jin SF, Chu TL, et al. (2006) The effects of group reminiscence therapy on depression, self esteem, and life satisfaction of elderly nursing home residents. *J Nurs Res* 14: 36-45.
31. Yasuda K, Nakamura T, Kuwabara K (2005) Development of a personal reminiscence video for dementia and its evaluation for concentration. *Kouji NouKinou Kenkyu* 25: 69-70.