

The Effect of Sleep Stage and Position on Postoperative Polysomnography

Jeffrey J. Stanley*

Department of Neurology and Otolaryngology-Head and Neck Surgery, University of Michigan, USA

*Corresponding author: Stanley JJ, Department of Neurology and Otolaryngology-Head and Neck Surgery, Sleep Disorders Center, University of Michigan, Ann Arbor, MI, USA, 1500 E. Medical Center Drive, Ann Arbor, MI 48109, USA, Tel: 734-615-1737; Fax: 734-936-9625; E-mail: jjst@med.umich.edu

Received date: July 11, 2018; Accepted Date: July 14, 2018; Published date: July 18, 2018

Copyright: ©2018 Stanley JJ. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Editorial

Polysomnography is the gold standard for the diagnosis of obstructive sleep apnea (OSA), evaluation of its severity, and assessment of treatment response. Sleep apnea syndrome has been defined as an apnea hypopnea index (AHI) greater than 5 with the presence of associated symptoms such as excessive daytime sleepiness. Because the correlation between AHI and morbidity and mortality is well established, it is frequently used as the primary parameter to describe surgical treatment outcomes. However, recent reports suggest that polysomnographic indices such as AHI may be discordant with quality of life measures [1-4].

One potential explanation for this discrepancy is that the overall AHI is frequently reported as the primary or sole postoperative outcome. Without taking into account night to night variability in sleep position and percentages of various sleep stages, including rapid eye movement (REM) sleep, this assessment may lead to an inaccurate reflection of the true effect of surgery.

The “first night effect”, where patients typically sleep less soundly during their first night in a sleep laboratory may contribute to an underestimation of disease severity by reducing the percentage of REM sleep during initial diagnostic testing. It is also well established that OSA itself can alter sleep architecture by significantly decreasing rapid eye movement (REM) sleep. A preoperative AHI may appear spuriously low in a patient with REM predominant OSA if REM duration is significantly reduced [5-10].

Sleep position can also have a significant effect on the severity of disease in a subset of patients with OSA. AHI in the supine sleep position is generally higher than that in lateral or prone sleep due to effect of gravity on the oropharyngeal and hypopharyngeal soft tissues. Oskenberg coined the term “positional sleep apnea” for patients who have a supine AHI at least twice that of their non-supine AHI. Hence, position-dependent patients may have highly variable overall AHIs depending on the proportion of time spent in the supine position on any given night. More than half of patients with OSA have positional obstructive sleep apnea. Patients who have mild to moderate OSA, are non-obese and younger are more likely to have positional OSA. This description entails a large proportion of CPAP intolerant considered to be good surgical candidates [11-14].

Lee et al. published a report investigating the effect of sleep position on surgical outcomes in obstructive sleep apnea. In this retrospective review of 69 consecutive patients who underwent uvulopalatopharyngoplasty, 85% of those who failed to improve had positional sleep apnea. When a position-corrected AHI was calculated in order to eliminate the effect of sleep position, >20% of patients were reclassified to a different severity category (mild, moderate or severe).

Although improvement was noted in the postoperative supine AHI compared to the preoperative supine AHI, the overall AHI increased in half of the positional sleep apnea patients due to an increase in proportion of time spent in the supine position.

A sleep stage/position corrected AHI can be utilized to eliminate the confounding effects of varying sleep position and stage during preoperative and postoperative polysomnography. This allows for a more accurate assessment of the effect of surgery [14-18].

Corrected postoperative AHI = $([\text{Preoperative \%TST in supine REM} \times \text{postoperative supine REM AHI}] + [\text{preoperative \%TST in supine non-REM} \times \text{postoperative supine non-REM AHI}] + [\text{preoperative \%TST in non-supine REM} \times \text{postoperative non-supine REM AHI}] + [\text{preoperative \%TST in non-supine non-REM} \times \text{postoperative non-supine non-REM AHI}]) / 100$. Alternatively, one may individually compare the preoperative vs. postoperative supine REM AHI, supine non-REM AHI, non-supine REM AHI and non-supine non-REM AHI.

The following case illustrates the importance of detailed postoperative polysomnography reporting: A 55-year old woman with worsening of excessive daytime sleepiness and morning headaches underwent diagnostic polysomnography (PSG) which revealed an AHI of 11 (markedly worse in REM sleep, AHI=75), an oxygen saturation nadir of 81% and REM sleep limited to 2% of total sleep time (TST). Due to persistent CPAP intolerance, the patient decided to pursue surgical treatment of her sleep apnea and subsequently underwent uvulopalatopharyngoplasty with tonsillectomy. The postoperative PSG demonstrated an apparent worsening of her OSA with an overall AHI of 17. Despite this, the patient reported subjective improvement in her symptoms and sleep quality. Physical examination revealed normal postoperative healing. A detailed review of her postoperative PSG revealed a significant increase in %REM sleep to 21% of TST. The patient slept exclusively in the supine position during both PSGs. A modest postoperative improvement was noted compared to the preoperative REM AHI (51 vs. 75) and non-REM AHI (8 vs. 10) when assessed separately.

An apparent worsening of OSA severity following surgery should be interpreted carefully by assessing not only the overall AHI but also the sleep architecture and sleep position. To account for night to night variability in both sleep stage and position, a corrected AHI can be calculated to allow for a more accurate assessment of the effect of surgery on the underlying OSA. Although a postoperative increase of supine sleep time and percentage of REM sleep may account for an increase in postoperative AHI, it is unclear if this should be interpreted as the cause of apparent surgical failure or the consequence of surgical treatment itself. Further research is needed to confirm the validity and utility of a sleep stage/position-corrected AHI [19-22].

References

1. Fujita S, Conway W, Zorick F, Rpth T (1981) Surgical correction of anatomic abnormalities in obstructive sleep apnea syndrome: Uvulopalatopharyngoplasty. *Otolaryngol Head Neck Surg* 89: 923-934.
2. Janson C, Gislason T, Bengtsson H (1997) Long-term follow-up of patients with obstructive sleep apnea treated with uvulopalatopharyngoplasty. *Arch Otolaryngol Head Neck Surg* 123: 257-262.
3. Hicklin LA, Tostevin P, Dasan S (2000) Retrospective survey of long-term results and patient satisfaction with uvulopalatopharyngoplasty for snoring. *J Laryngol Otol* 114: 675-681.
4. Weaver EM, Woodson BT, Steward DL (2005) Polysomnography indexes are discordant with quality of life, symptoms, and reaction times in sleep apnea patients. *Otolaryngol Head Neck Surg* 132: 255-262.
5. Redline S, Budhiraja R, Kapur V (2007). The scoring of respiratory events in sleep: Reliability and validity. *J Clin Sleep Med* 3: 169-200.
6. Richard W, Kox D, den Herder C (2006) The role of sleep position in obstructive sleep apnea syndrome. *Eur Arch Otorhinolaryngol* 263: 946-950.
7. Cartwright RD, Samelson CF (1982) The effects of a nonsurgical treatment for obstructive sleep apnea. The tongue-retaining device. *JAMA* 248: 705-709.
8. Cartwright RD, Diaz F, Lloyd S (1991) The effects of sleep posture and sleep stage on apnea frequency. *Sleep* 14: 351-353.
9. Cartwright RD (1984) Effect of sleep position on sleep apnea severity. *Sleep* 7: 110-104.
10. Akita Y, Kawakatsu K, Hattori C (2003) Posture of patients with sleep apnea during sleep. *Acta Otolaryngol Suppl* 41-45.
11. Oksenberg A, Silverberg DS, Arons E (1997) Positional vs nonpositional obstructive sleep apnea patients: anthropomorphic, nocturnal polysomnographic, and multiple sleep latency test data. *Chest* 112: 629-639.
12. Indications and standards for cardiopulmonary sleep studies (1989) American thoracic society. Medical section of the American lung association. *Am Rev Respir Dis* 139: 559-568.
13. Sleep-related breathing disorders in adults (1999) Recommendations for syndrome definition and measurement techniques in clinical research. The report of an American Academy of sleep medicine task force. *Sleep* 22: 667-689.
14. Can M, Acikgoz S, Mungan G (2006) Serum cardiovascular risk factors in obstructive sleep apnea. *Chest* 129: 233-237.
15. Peker Y, Hedner J, Norum J (2002) Increased incidence of cardiovascular disease in middle-aged men with obstructive sleep apnea: A 7-year follow-up. *Am J Respir Crit Care Med* 166: 159-165.
16. Nishibayashi M, Miyamoto M, Miyamoto T (2008) Correlation between severity of obstructive sleep apnea and prevalence of silent cerebrovascular lesions. *J Clin Sleep Med* 4: 242-247.
17. Zhang XW, Li Y, Zhou F (2007) Association of body position with sleep architecture and respiratory disturbances in children with obstructive sleep apnea. *Acta Otolaryngol* 1-6.
18. Oksenberg A, Silverberg D, Offenbach D (2006) Positional therapy for obstructive sleep apnea patients: A 6-month follow-up study. *Laryngoscope* 116: 1995-2000.
19. Nakano H, Ikeda T, Hayashi M (2003) Effects of body position on snoring in apneic and nonapneic snorers. *Sleep* 26: 169-172.
20. Oksenberg A, Khamaysi I, Silverberg DS (2000) Association of body position with severity of apneic events in patients with severe nonpositional obstructive sleep apnea. *Chest* 118: 1018-1024.
21. Oksenberg A, Silverberg DS, Arons E (1997) Positional vs nonpositional obstructive sleep apnea patients: Anthropomorphic, nocturnal polysomnographic, and multiple sleep latency test data. *Chest* 112: 629-639.
22. Lee CH, Shin HW, Han DH (2009) The implication of sleep position in the evaluation of surgical outcome in obstructive sleep apnea. *Otolaryngol Head Neck Surg* 140: 531-535.