OPEN ACCESS Freely available online

Anatomy & Physiology: Current Research

Commentry

The Role of Cellular Physiology in Tissue Regeneration and Repair

Javier Morales*

Department of Functional Anatomy, University of Andalusian Health Sciences, Seville, Andalusia, Spain

DESCRIPTION

Tissue regeneration and repair are fundamental biological processes that enable the human body to recover from injury, infection and degenerative changes. Although some tissues possess remarkable regenerative capacity while others heal primarily through scar formation, successful recovery depends on highly coordinated cellular and physiological mechanisms that restore structural integrity and functional performance. Advances in cell biology, molecular physiology, regenerative medicine, and tissue engineering have significantly expanded our understanding of how cells communicate, proliferate, differentiate and remodel damaged tissues. These discoveries are transforming regenerative medicine by shifting therapeutic strategies from symptom management toward biological restoration of normal tissue function. Cellular physiology forms the foundation of tissue regeneration. Every tissue contains specialized cells responsible for maintaining normal structure and physiological function under healthy conditions. Following injury, these cells rapidly respond to biochemical signals generated within the damaged microenvironment. Growth factors, cytokines, chemokines, extracellular matrix proteins and mechanical stimuli collectively activate complex intracellular signaling pathways that regulate cell survival, proliferation, migration and differentiation. Rather than functioning independently, these processes occur in a precisely coordinated sequence that supports effective tissue repair while minimizing excessive inflammation or fibrosis.

The repair process generally begins with the inflammatory phase, which serves as an essential physiological response rather than merely a pathological event. Following tissue injury, platelets initiate clot formation while releasing growth factors that attract neutrophils, macrophages, and other immune cells to the affected region. These immune cells remove damaged cellular debris, eliminate pathogens and secrete cytokines that regulate subsequent stages of healing. In my opinion, the modern understanding of inflammation as a beneficial and carefully controlled component of regeneration has fundamentally changed approaches to tissue repair and regenerative medicine. Cell proliferation and migration represent the next critical stages of regeneration. Resident tissue cells, endothelial cells, fibroblasts, and stem cells become activated and migrate toward the injured site to replace damaged structures. Simultaneously, angiogenesis promotes the formation of new blood vessels that supply oxygen and nutrients while facilitating waste removal. Adequate vascularization is indispensable because newly regenerating tissues require continuous metabolic support to sustain cellular growth and functional recovery. This close relationship between cellular physiology and vascular development illustrates the integrated nature of tissue regeneration.

Stem cells play a particularly important role in regenerative physiology. Adult stem cells, including mesenchymal stem cells, hematopoietic stem cells and tissue specific progenitor cells, possess the ability to self renew and differentiate into specialized cell types required for tissue repair. Beyond direct cellular replacement, stem cells release numerous bioactive molecules, including growth factors, cytokines and extracellular vesicles, that modulate inflammation, stimulate angiogenesis and activate endogenous repair mechanisms. These paracrine effects have become increasingly recognized as major contributors to successful tissue regeneration and have expanded therapeutic interest in stem cell based regenerative strategies. The Extracellular Matrix (ECM) also plays a vital physiological role in tissue repair. Traditionally regarded as a passive structural scaffold, the ECM is now recognized as a dynamic biological environment that regulates cell adhesion, migration, proliferation and differentiation. During regeneration, matrix proteins undergo continuous remodeling through the coordinated activity of matrix metalloproteinases and their inhibitors. Proper ECM remodeling facilitates organized tissue reconstruction, whereas excessive collagen deposition contributes to fibrosis and impaired organ function. Understanding ECM physiology has therefore become central to the development of biomaterials and tissue engineering approaches. Cellular metabolism has emerged as another important determinant of regenerative capacity. Regenerating cells undergo profound metabolic reprogramming to meet the increased energy demands associated with proliferation, protein synthesis and extracellular matrix production.

Correspondence to: Javier Morales, Department of Functional Anatomy, University of Andalusian Health Sciences, Seville, Andalusia, Spain, E-mail: javier.morales@uahs.es

Received: 01-Sep-2025, Manuscript No. APCR-25-43608; **Editor assigned:** 03-Sep-2025, PreQC No. APCR-25-43608 (PQ); **Reviewed:** 17-Sep-2025, QC No. APCR-25-43608; **Revised:** 24-Sep-2025, Manuscript No. APCR-25-43608 (R); **Published:** 01-Oct-2025, DOI: 10.35248/2161-0940.25.15.563

Citation: Morales J (2025). The Role of Cellular Physiology in Tissue Regeneration and Repair. Anat Physiol. 15:563.

Copyright: © 2025 Morales J. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.