OPEN ACCESS Freely available online

Gynecology & Obstetrics

Review Article

The Role of Genetics in Early Pregnancy Loss

Maria Fernandez*

Department of Business Economics, Rey Juan Carlos University, Madrid, Spain

ABSTRACT

Early Pregnancy Loss (EPL), often referred to as miscarriage, is a common event, occurring in approximately 10–20% of clinically recognized pregnancies. A variety of factors, including genetic abnormalities, contribute to its incidence. Advances in genetic testing have significantly enhanced the understanding of the underlying causes of early pregnancy loss, revealing the role of chromosomal abnormalities, single-gene mutations, and other genetic factors. This paper explores the role of genetics in early pregnancy loss, highlighting the genetic causes, diagnostic tools, and clinical management strategies. Through genetic screening and counseling, healthcare providers can offer insights into the potential causes of miscarriage and provide guidance for future pregnancies. However, despite advances in genetics, many cases of early pregnancy loss remain unexplained, indicating the complexity of the condition. This review concludes with an emphasis on the importance of personalized medicine and ongoing research to better understand the genetic contributors to early pregnancy loss.

Keywords: Early pregnancy loss; Miscarriage; Genetics; Chromosomal abnormalities; Single-gene mutations; Genetic screening; Prenatal care

INTRODUCTION

Early Pregnancy Loss (EPL), also known as miscarriage, refers to the spontaneous loss of a pregnancy before the 20th week of gestation. This event is deeply distressing and emotionally challenging for women and their families. It is estimated that approximately 10–20% of clinically recognized pregnancies result in miscarriage, with the majority of these losses occurring in the first trimester. The emotional impact of early pregnancy loss can be profound, as it often brings about feelings of grief, loss, and emotional distress, and in some cases, it can also cause long-term psychological effects. However, while the emotional toll is widely recognized, the physical causes underlying miscarriage have been a focal point of scientific and medical research for decades. Understanding the causes of early pregnancy loss is crucial for improving prenatal care and enhancing outcomes for future pregnancies [1].

One of the key contributors to early pregnancy loss is genetic abnormalities, which can occur in both the mother and the fetus. Genetic factors are often responsible for the majority of early pregnancy losses, with chromosomal abnormalities being the most commonly identified cause. These abnormalities can arise due to errors during cell division, which may result in the fetus having an abnormal number of chromosomes or structural defects in the chromosomes. Genetic factors affecting the mother, such as inherited conditions or abnormalities, may also contribute to miscarriage, although fetal genetic abnormalities are more commonly implicated [2].

Advancements in genetic testing have revolutionized our understanding of the genetic underpinnings of early pregnancy loss. Techniques such as karyotyping, genetic screening, and next-generation sequencing have enabled clinicians to identify chromosomal abnormalities, such as aneuploidy, and single-gene mutations that can contribute to miscarriage. By identifying these genetic causes, healthcare providers can offer valuable insights into the potential reasons behind a miscarriage and provide guidance for future pregnancies. These advances in genetics have paved the way for a better understanding of the biological mechanisms involved in pregnancy loss, allowing for improved diagnostic accuracy and personalized interventions [3].

This review aims to explore the role of genetics in early pregnancy loss, providing an in-depth analysis of the genetic

Correspondence to: Maria Fernandez, Department of Business Economics, Rey Juan Carlos University, Madrid, Spain; E-mail: mariafernandez@gmail.com

Received: 29-Jan-2025, Manuscript No. GOCR-25-36917; **Editor assigned:** 01-Feb-2025, PreQC No. GOCR-25-36917 (PQ); **Reviewed:** 15-Feb-2025, QC No. GOCR-25-36917; **Revised:** 07-Mar-2025, Manuscript No. GOCR-25-36917 (R); **Published:** 14-Mar-2025, DOI: 10.35248/2161-0932.25.15.635

Citation: Fernandez M (2025) The Role of Genetics in Early Pregnancy Loss. Gynecol Obster. 15:635.

Copyright: © 2025 Fernandez M. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

factors that contribute to miscarriage. It will also examine the various diagnostic approaches that are used to identify genetic abnormalities in early pregnancy loss, as well as the potential management strategies available to couples who experience miscarriage. Through these efforts, the review will highlight how genetic testing and counseling can offer women and their partners critical information about their reproductive health, enabling more informed decisions and personalized care in subsequent pregnancies [4].

LITERATURE REVIEW

Genetic factors in early pregnancy loss

Genetics plays a critical role in early pregnancy loss, with chromosomal abnormalities being the most common cause. These abnormalities can occur as a result of errors during cell division, leading to an abnormal number of chromosomes or structural defects in the chromosomes. The most common genetic cause of miscarriage is chromosomal aneuploidy, particularly trisomy (the presence of an extra chromosome), with trisomy 16 being the most frequently observed abnormality. In many cases, these chromosomal abnormalities are not inherited but arise de novo during early fetal development [5].

Chromosomal abnormalities: A significant percentage of early pregnancy losses are attributed to chromosomal abnormalities, particularly aneuploidy. Chromosomal abnormalities, such as trisomy 13, 18, and 21, are commonly observed in early pregnancy losses. In fact, about 50% of first-trimester miscarriages are due to aneuploidy, with trisomy 16 being the most frequently encountered abnormality. Trisomy 16 is particularly notable because it is typically not compatible with life, leading to spontaneous miscarriage [6].

Other chromosomal abnormalities that can contribute to early pregnancy loss include structural abnormalities such as translocations, inversions, and deletions. These chromosomal rearrangements can disrupt fetal development, leading to miscarriage or congenital abnormalities if the pregnancy is carried to term. Balanced translocations, where portions of chromosomes are swapped without a loss of genetic material, can also increase the risk of miscarriage, as the rearranged chromosomes may lead to the production of unbalanced gametes that result in fetal abnormalities.

Single-gene mutations: In addition to chromosomal abnormalities, single-gene mutations have been implicated in some cases of early pregnancy loss. These mutations can affect critical genes involved in fetal development, leading to malformations or failure to implant. For example, mutations in genes responsible for DNA repair, cell division, and apoptosis may increase the risk of early pregnancy loss. While single-gene mutations are less common than chromosomal abnormalities, they still play a role in certain cases of miscarriage [7].

Inherited genetic disorders, such as cystic fibrosis, sickle cell anemia, and thalassemia, can also contribute to early pregnancy loss if the fetus inherits a mutation from both parents. Carrier screening for these disorders can provide valuable information for couples planning a pregnancy, allowing them to make informed decisions about reproductive options, such as preimplantation genetic testing or genetic counseling.

Mitochondrial disorders: Mitochondrial DNA is inherited exclusively from the mother, and mutations in the mitochondrial genome can lead to a variety of disorders, some of which may contribute to early pregnancy loss. Mitochondrial disorders can affect energy production within the cell, leading to fetal developmental defects and miscarriage. These conditions are relatively rare but are an important consideration in cases of recurrent pregnancy loss where other genetic causes have been ruled out [8].

DISCUSSION

Genetic testing in early pregnancy loss

Advancements in genetic testing have provided valuable insights into the genetic causes of early pregnancy loss. Genetic testing can help identify chromosomal abnormalities, single-gene mutations, and other genetic factors that may contribute to miscarriage. Several methods are available for testing genetic material, including karyotyping, Fluorescence *In situ* Hybridization (FISH), array Comparative Genomic Hybridization (aCGH), and Next-Generation Sequencing (NGS). These techniques allow for detailed analysis of chromosomal structure and the identification of genetic mutations.

Karyotyping: Karyotyping is a traditional method of examining chromosomes to detect structural abnormalities and aneuploidy. In cases of early pregnancy loss, karyotyping of fetal tissue obtained from products of conception can provide information about the chromosomal makeup of the fetus. However, karyotyping has limitations, particularly in detecting smaller genetic changes such as microdeletions or microduplications [9].

Fluorescence *In situ* Hybridization (FISH): FISH is a technique that uses fluorescent probes to detect specific chromosomal abnormalities. It is particularly useful for identifying aneuploidy and translocations in fetal tissue. FISH is more rapid than karyotyping and can be performed on samples obtained from the products of conception.

Array Comparative Genomic Hybridization (aCGH): aCGH is a high-resolution technique that allows for the detection of chromosomal microdeletions and microduplications. It is particularly useful in identifying submicroscopic genetic imbalances that may not be detectable by traditional karyotyping or FISH. This method has become a valuable tool in the diagnosis of recurrent pregnancy loss, as it can identify genetic causes that may have been missed by other testing methods.

Next-Generation Sequencing (NGS): NGS technology allows for the sequencing of large portions of the genome, providing detailed information about genetic mutations that may contribute to early pregnancy loss. NGS is particularly useful for identifying single-gene mutations and structural variations in the genome that may affect fetal development. This technique has the potential to revolutionize the understanding of genetic

causes of miscarriage, allowing for more personalized and targeted interventions.

Clinical management and counseling

The identification of genetic factors in early pregnancy loss has important implications for clinical management and counseling. Genetic counseling is a critical component of the care of women who experience miscarriage, particularly when a genetic cause is suspected. Genetic counselors can help couples understand the potential causes of pregnancy loss, discuss options for future pregnancies, and provide emotional support during a difficult time.

Recurrent Pregnancy Loss (RPL): Women who experience two or more consecutive early pregnancy losses are often referred for further evaluation to identify potential genetic causes. In these cases, genetic testing of both parents may be recommended to identify inherited genetic mutations or chromosomal rearrangements that could increase the risk of miscarriage. Carrier screening for common genetic disorders can also provide valuable information about the risk of inherited conditions in future pregnancies.

Preimplantation Genetic Testing (PGT): For couples with a known genetic risk of miscarriage, Preimplantation Genetic Testing (PGT) may be an option. PGT involves testing embryos created through *In vitro* Fertilization (IVF) for genetic abnormalities before implantation. This technique can help ensure that only embryos without genetic abnormalities are implanted, reducing the risk of miscarriage and improving the chances of a successful pregnancy.

Targeted therapies: In cases where a specific genetic cause of miscarriage is identified, targeted therapies may be available. For example, women with inherited thrombophilia may be prescribed anticoagulant medications to reduce the risk of miscarriage due to blood clotting abnormalities. Similarly, women with single-gene mutations may benefit from gene therapy or other interventions to correct the genetic defect.

CONCLUSION

Genetics plays a pivotal and multifaceted role in Early Pregnancy Loss (EPL), with chromosomal abnormalities and single-gene mutations being identified as the most common causes. Early pregnancy loss, often occurring in the first trimester, is frequently caused by errors in the genetic material of the developing fetus. These genetic issues can arise from a range of factors, including the abnormal number of chromosomes, known as aneuploidy, or structural defects in the chromosomes, which can lead to developmental abnormalities that are not compatible with life. Moreover, single-gene mutations, which can be inherited or arise spontaneously, may also contribute to pregnancy loss. These genetic factors are responsible for a significant proportion of early miscarriages and are often the first area of investigation when women experience recurrent miscarriages.

Over the years, advances in genetic testing technologies have provided invaluable insights into the genetic underpinnings of early pregnancy loss. The development of more sophisticated techniques, such as karyotyping, Fluorescence *In situ* Hybridization (FISH), and Next-Generation Sequencing (NGS), has enabled the identification of genetic abnormalities with greater accuracy and precision. These advancements allow for the identification of specific chromosomal abnormalities, such as trisomy, monosomy, and structural rearrangements, and single-gene mutations that may predispose to miscarriage. As a result, genetic testing has become an essential tool in the diagnostic work-up of women experiencing recurrent pregnancy loss, providing both clinicians and patients with more information about the potential genetic causes of the miscarriage and offering an opportunity for targeted management.

Despite the progress made in understanding the genetic causes of early pregnancy loss, a significant proportion of cases remain unexplained. This highlights the complexity and multifactorial nature of early pregnancy loss, suggesting that other genetic factors, as well as environmental, immunological, and hormonal influences, may also contribute to the occurrence of miscarriage. Some genetic causes may not yet be detectable with current technologies, and further research is needed to uncover additional genetic markers and mechanisms that may be involved. This ongoing research is essential for advancing the field and for improving clinical outcomes for women who experience miscarriage.

The role of genetic counseling has also become increasingly important in the management of early pregnancy loss. Genetic counseling provides affected couples with guidance, emotional support, and information about the genetic causes of miscarriage, as well as the implications for future pregnancies. In addition to genetic counseling, Preimplantation Genetic Testing (PGT) has emerged as a valuable tool in reducing the risk of miscarriage. PGT involves testing embryos for genetic abnormalities before implantation during *In vitro* Fertilization (IVF), allowing for the selection of embryos with normal chromosomal patterns, thus improving the chances of a successful pregnancy. Furthermore, the development of targeted therapies aimed at correcting or managing genetic issues in pregnancy is an exciting area of ongoing research, which may one day provide women with new options for addressing recurrent pregnancy loss.

In conclusion, while significant strides have been made in understanding the genetic factors involved in early pregnancy loss, there is still much to be learned about the complex interplay between genetics and miscarriage. Continued advancements in genetic testing, research into unexplained cases, and the integration of genetic counseling and preimplantation genetic testing into clinical practice hold great promise for improving the management of early pregnancy loss. By reducing the emotional and physical toll of miscarriage and providing couples with the tools to make informed decisions, these innovations could greatly improve the outcomes for women experiencing early pregnancy loss and help them achieve healthy pregnancies in the future.

REFERENCES

1. Romero ST, Geiersbach KB, Paxton CN, Rose NC, Schisterman EF, Branch DW, et al. Differentiation of genetic abnormalities in early pregnancy loss. Ultrasound Obstet Gynecol. 2015;45(1):89-94.
2. Qiao Y, Wen J, Tang F, Martell S, Shomer N, Leung PC, et al. Whole exome sequencing in recurrent early pregnancy loss. Mol Hum Reprod. 2016;22(5):364-372.
3. Kutteh WH, Papas RS, Maisenbacher MK, Dahdouh EM. Role of genetic analysis of products of conception and PGT in managing early pregnancy loss. Reprod Biomed Online. 2024;49(1):103738.
4. Li Q, Chen S, Dong X, Fu S, Zhang T, Zheng W, et al. The progress of research on genetic factors of recurrent pregnancy loss. Genet Res. 2023;2023(1):9164374.
5. Robbins SM, Thimm MA, Valle D, Jelin AC. Genetic diagnosis in first or second trimester pregnancy loss using exome sequencing: a systematic review of human essential genes. J Assist Reprod Genet. 2019;36(8):1539-1548.
6. Macklon NS, Geraedts JP, Fauser BC. Conception to ongoing pregnancy: the 'black box'of early pregnancy loss. Hum Reprod Update. 2002;8(4):333-343.
7. Page JM, Silver RM. Genetic causes of recurrent pregnancy loss. Clin Obstet Gynecol. 2016;59(3):498-508.
8. Muncey W, Scott M, Lathi RB, Eisenberg ML. The paternal role in pregnancy loss. Andrology. 2025;13(1):146-150.
9. Wang Y, Li Y, Chen Y, Zhou R, Sang Z, Meng L, et al. Systematic analysis of copy-number variations associated with early pregnancy loss. Ultrasound Obstet Gynecol. 2020;55(1):96-104.