Journal of Cancer Research and Immuno-Oncology

OPEN ACCESS Freely available online

Commentary

Tumor-Associated Macrophage Diversity and Its Clinical Relevance in Cancer Immuno-Oncology

Elian Foster*

Department of Oncology and Cellular Medicine, North Atlantic Medical University, Dublin, Ireland

DESCRIPTION

Cancer continues to present a significant clinical challenge despite continuous improvements in diagnosis, surgical procedures, radiation therapy, systemic treatment, and supportive care. During the past several decades, scientific attention has expanded beyond malignant cells alone to include the complex biological environment surrounding tumors. This environment contains numerous non-malignant cells, extracellular components, blood vessels, signaling molecules, and immune populations that collectively influence disease behavior. Among these cellular populations, macrophages have attracted considerable interest because they participate in tissue maintenance, immune regulation, repair processes, and inflammatory responses. Within malignant tissues, these cells are commonly referred to as tumor-associated macrophages and represent one of the most abundant immune cell populations found across many solid tumors.

Macrophages originate from circulating monocytes or tissue-resident precursor cells and normally contribute to the removal of damaged cells, elimination of infectious organisms, and maintenance of tissue equilibrium. In healthy organs, these cells respond rapidly to environmental signals and adapt their biological activities according to local conditions. Within malignant tissue, however, continuous exposure to altered signaling molecules, metabolic products, oxygen deprivation, and cellular debris changes their biological behavior. Rather than maintaining normal tissue function, these macrophages may gradually acquire characteristics that support malignant progression. Their activities differ according to cancer type, anatomical location, disease stage, and surrounding immune composition.

Current evidence indicates that macrophages cannot be accurately described using a simple binary classification. Earlier investigations frequently categorized these cells into inflammatory and anti-inflammatory groups, but more recent observations indicate that macrophage populations display a broad spectrum of biological characteristics. Individual macrophages may simultaneously express markers associated with several functional programs while continuously adapting to changing environmental conditions. This biological flexibility explains why macrophage behavior varies considerably between patients and even among different regions of the same tumor.

Within malignant tissues, cancer cells continuously release cytokines, chemokines, growth factors, and extracellular vesicles that influence macrophage recruitment and differentiation. Stromal fibroblasts, endothelial cells, adipocytes, and infiltrating immune cells also contribute signaling molecules that shape macrophage behavior. Oxygen limitation within rapidly growing tumors produces additional metabolic stress, leading to accumulation of lactate and altered nutrient availability. These local conditions modify macrophage metabolism, affecting cytokine production, antigen presentation, and interactions with neighboring immune cells.

One important characteristic of tumor-associated macrophages involves their communication with lymphocytes responsible for recognizing abnormal cells. Effective anti-tumor immunity depends upon successful activation of cytotoxic lymphocytes capable of identifying malignant antigens. Macrophages influence this process through antigen presentation, secretion of signaling molecules, and expression of immune regulatory proteins. Depending upon their functional state, macrophages may either support lymphocyte activation or reduce immune responsiveness. In several malignancies, increased macrophage infiltration has been associated with diminished cytotoxic lymphocyte activity and reduced immune-mediated elimination of cancer cells.

Blood vessel formation represents another important biological process influenced by macrophages. Rapidly growing tumors require continuous delivery of oxygen and nutrients to sustain expansion. Macrophages located near areas of oxygen limitation release multiple signaling molecules that stimulate endothelial cell migration and vascular development. Newly formed vessels within tumors often display abnormal architecture, resulting in irregular blood flow, impaired oxygen delivery, and variable drug distribution. These vascular abnormalities contribute to regional differences in therapeutic response and create additional biological diversity within malignant tissues.

Correspondence to: Elian Foster, Department of Oncology and Cellular Medicine, North Atlantic Medical University, Dublin, Ireland, E-mail: elian.foster@namu-ireland.edu

Received: 27-Feb-2026, Manuscript No. JCRIO-26-43977; **Editor assigned:** 02-Mar-2026, PreQC No. JCRIO-26-43977 (PQ); **Reviewed:** 16-Mar-2026, QC No. JCRIO-26-43977; **Revised:** 23-Mar-2026, Manuscript No. JCRIO-26-43977 (R); **Published:** 30-Mar-2026, DOI: 10.35248/2684-1266.26.12.276

Citation: Foster E (2026). Tumor-Associated Macrophage Diversity and Its Clinical Relevance in Cancer Immuno-Oncology. J Cancer Res Immunooncol. 11:276.

Copyright: © 2026 Foster E. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Spatial transcriptomic techniques provide additional information by preserving tissue architecture while analyzing molecular expression patterns. Rather than examining isolated cells alone, researchers can determine how macrophage populations are distributed throughout different tumor regions. This information has demonstrated that macrophage characteristics frequently differ according to their proximity to blood vessels, necrotic tissue, invasive margins, or lymphocyte-rich regions. These spatial relationships may influence therapeutic response and provide valuable information for future biomarker development.

Liquid biopsy technologies may also contribute to macrophage-related biomarker development. Analysis of circulating extracellular vesicles, cytokines, soluble receptors, and nucleic acids offers opportunities to monitor biological changes during disease progression and treatment. Although tissue biopsy remains essential for diagnosis, repeated sampling can be difficult in many patients. Blood-based biomarkers associated with macrophage activity may provide complementary information regarding treatment response, disease recurrence, or alterations within the tumor microenvironment over time.

CONCLUSION

Continued investigation into tumor-associated macrophages has expanded current understanding of cancer immuno-oncology by demonstrating that these cells participate in immune regulation, tissue remodeling, angiogenesis, metastatic dissemination, and therapeutic response. Their biological diversity reflects continuous adaptation to local environmental conditions, emphasizing the importance of studying cellular interactions rather than individual cell populations in isolation. As experimental technologies continue to improve, macrophage-directed research may contribute to more refined diagnostic biomarkers, improved patient stratification, and innovative therapeutic combinations that strengthen anti-tumor immune activity while supporting more effective clinical management across a wide range of malignancies.